

Understanding exosomes: Part 2—Emerging leaders in regenerative medicine

Richard J. Miron¹ | Nathan E. Estrin^{2,3} | Anton Sculean¹ | Yufeng Zhang⁴ 

¹Department of Periodontology,
University of Bern, Bern, Switzerland

²Advanced PRF Education, Venice, Florida,
USA

³School of Dental Medicine, Lake Erie
College of Osteopathic Medicine,
Bradenton, Florida, USA

⁴Department of Oral Implantology,
University of Wuhan, Wuhan, China

Correspondence

Richard J. Miron, Department of
Periodontology, University of Bern, Bern,
Switzerland.

Email: rick@themironlab.com

Abstract

Exosomes are the smallest subset of extracellular signaling vesicles secreted by most cells with the ability to communicate with other tissues and cell types over long distances. Their use in regenerative medicine has gained tremendous momentum recently due to their ability to be utilized as therapeutic options for a wide array of diseases/conditions. Over 5000 publications are currently being published yearly on this topic, and this number is only expected to dramatically increase as novel therapeutic strategies continue to be developed. Today exosomes have been applied in numerous contexts including neurodegenerative disorders (Alzheimer's disease, central nervous system, depression, multiple sclerosis, Parkinson's disease, post-traumatic stress disorders, traumatic brain injury, peripheral nerve injury), damaged organs (heart, kidney, liver, stroke, myocardial infarctions, myocardial infarctions, ovaries), degenerative processes (atherosclerosis, diabetes, hematology disorders, musculoskeletal degeneration, osteoradionecrosis, respiratory disease), infectious diseases (COVID-19, hepatitis), regenerative procedures (antiaging, bone regeneration, cartilage/joint regeneration, osteoarthritis, cutaneous wounds, dental regeneration, dermatology/skin regeneration, erectile dysfunction, hair regrowth, intervertebral disc repair, spinal cord injury, vascular regeneration), and cancer therapy (breast, colorectal, gastric cancer and osteosarcomas), immune function (allergy, autoimmune disorders, immune regulation, inflammatory diseases, lupus, rheumatoid arthritis). This scoping review is a first of its kind aimed at summarizing the extensive regenerative potential of exosomes over a broad range of diseases and disorders.

KEYWORDS

dermasomes, exosomes, immunosomes, regenerative medicine, ultra-centrifugation

1 | INTRODUCTION

Exosomes are the smallest extracellular vesicles ranging in size from 30 to 150 nm and are found in the majority of bodily fluids.¹ They can transport functional miRNAs, coupled with the inherent biological

functions of exosomes, which makes them a highly sought after and novel delivery platform.² Specifically, they have the characteristics of low immunogenicity and high transport efficiency, which can regulate inflammation and cross the blood-brain barrier (BBB).³⁻⁵ They transport signaling molecules that drive a number of biological

This is an open access article under the terms of the [Creative Commons Attribution](https://creativecommons.org/licenses/by/4.0/) License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2024 The Authors. *Periodontology 2000* published by John Wiley & Sons Ltd.

processes, including cell signaling, immunological responses, tumor metastasis, and other biological activities. Exosomes have now been shown in several investigations to play both diagnostic and therapeutic functions, in which their precise detection, separation, and quantification are crucial. Today, exosomes are one of the most highly researched topics in regenerative medicine with over 5000 publications being published on the topic yearly. Part 1 of this three-part series on exosomes focused on exosome biogenesis as well as their standard isolation techniques, including ultra-centrifugation, microfluidic, immunoaffinity, precipitation, size-exclusion chromatography, ultrafiltration technologies. This second article is focused on the therapeutic potential of exosomes in medicine. It is divided into eight categories highlighting the therapeutic potential of exosomes as follows:

- Neurodegenerative disorders: Alzheimer's disease, central nervous system, depression, multiple sclerosis, Parkinson's disease, post-traumatic stress disorders, traumatic brain injury, peripheral nerve injury.
- Treatment of damaged organs: heart, kidney, liver, stroke, myocardial infarctions, myocardial infarctions, ovaries.
- Degenerative processes: atherosclerosis, diabetes, hematology disorders, musculoskeletal degeneration, osteoradionecrosis, respiratory disease.
- Treatment of infectious diseases: COVID-19, hepatitis.
- Regenerative procedures: antiaging, bone regeneration, cartilage/joint regeneration, osteoarthritis, cutaneous wounds, dental regeneration, dermatology/skin regeneration, erectile dysfunction, hair regrowth, intervertebral disc repair, spinal cord injury, vascular regeneration.
- Cancer therapy: breast cancer, colorectal cancer, gastric cancer, and osteosarcomas.
- Improvements in immune function: allergy, autoimmune disorders, immune regulation, inflammatory diseases, lupus, rheumatoid arthritis.
- Treatment of random disorders: infertility, obesity, and sleep apnea.

The aim of this scoping review is to summarize the extensive regenerative potential of exosomes over a broad range of disorders and to elaborate on the potential clinical relevance of this novel treatment approach.

2 | EXOSOMES AND NEURODEGENERATIVE DISEASE

The medical community is persistently seeking appropriate treatment solutions for neurodegenerative illnesses due to the limitations of short-term symptomatic therapy and the dose-dependent adverse effects associated with pharmaceutical treatments. This pursuit is driven by the recognition of neurodegenerative disorders as a significant global health problem.⁶ The identification regarding

the therapeutic potential of stem cells for the treatment of neurodegenerative illnesses dates back to 1980 when Parkinson's disease (PD) was treated using fetal nerve tissue. Subsequently, a multitude of comprehensive investigations have been undertaken to formulate an effective therapeutic approach for the treatment of neurological disorders. Currently, there is a significant body of knowledge on the therapeutic potential of stem cells and their secreted factors in the context of treating neurodegenerative illnesses. This novel framework has shown distinct attributes pertaining to this therapeutic approach, including neuroprotective and neurodegenerative effects, remyelination capabilities, mitigation of brain inflammation, and restoration of functionality subsequent to produced damage.⁶ Nevertheless, the precise method by which stem cells facilitate the healing of nerve injury remains uncertain. One significant component of their secretory function, exosomes, has been proposed as a key contributor to these therapeutic benefits. A multitude of research conducted in recent decades has examined the therapeutic efficacy of exosomes in the management of many neurological disorders. The primary objective of this review article is to examine the potential of stem cell-derived exosome-based treatments in the context of treating and managing neurodegenerative disorders.⁶

Regrettably, the occurrence of diverse neurodegenerative diseases, such as amyotrophic lateral sclerosis (ALS), PD, multiple sclerosis (MS), Alzheimer's disease (AD), and Huntington's disease (HD), has been attributed to the progressive deterioration of neuronal function and structure in both the peripheral and central nervous systems.⁷ The process of aging is well recognized as the primary risk factor for several brain disorders.⁸ With the global elderly population seeing significant growth, it is anticipated that degenerative neurological illnesses will surpass cancer in the foreseeable future, therefore becoming the second most prevalent cause of mortality worldwide.⁹ Neurodegenerative disorders have a significant influence on several aspects of individual functioning, including but not limited to balance, motor skills, cognitive abilities, language proficiency, and respiratory function.¹⁰ While the etiology of several illnesses remains unclear, a confluence of hereditary and environmental variables might potentially contribute to their development.¹⁰ Most often, these disorders cause neurons to die thus affecting distinct parts of the brain, such as the striatal regions in PD and the cortical and hippocampal regions in AD.¹¹ These conditions may present with minor or severe symptoms, depending on the part of the brain that is impacted.¹² The presence of these diseases, characterized by their prolonged duration and significant treatment expenses, poses a substantial challenge for both patients and the healthcare community. Consequently, considerable efforts are being directed toward the development of efficacious treatments to address this global issue.

Currently, there is a lack of definitive cures for these diseases, with the majority of treatment approaches in Western Medicine focusing on symptom management, pain relief, symptom control, and enhancement of mobility.¹¹ During the 1970s, the first pharmacotherapy groups centered on target-based approaches emerged, serving as substitutes for second messenger modulators, direct

receptor agonists, enzyme inhibitors, releasing agents, and neurotransmitters. Among the first generation of these drugs are those used to replace dopamine, such as dopaminergic drugs,¹³ acetylcholinesterase inhibitors,¹⁴ analgesic drugs,¹⁵ and surgical treatments, like deep brain stimulation, for the management of many movement disorders.¹⁶ Cholinesterase inhibitors, another significant family of FDA-approved medications used to treat all stages of AD since 1978, and L-DOPA (L-3,4-dihydroxyphenylalanine), the first drug authorized for the clinical treatment of PD.¹⁷⁻²⁰ The goal of the second generation of these medications is to halt and reduce the disease's development. Examples include riluzole for Huntington's disease, cerebellar ataxia, and ALS²¹; nonsteroidal anti-inflammatory medications (NSAIDs) that lower the risk of AD with long-term usage²²; and CERE-120 (adeno-associated virus serotype 2-neurturin), which is in phase I study for people with idiopathic PD.²³⁻²⁵

The blood-brain barrier (BBB) is a specialized membrane that exhibits extreme selectivity in its permeability, effectively restricting the passage of big molecules and almost all small molecules from peripheral organs to the brain. Due to this rationale, a range of intrusive methodologies have been devised or are currently being explored to overcome the selective nature of the BBB. These procedures include neurosurgery, the biochemical manipulation of the BBB, and the use of diverse nanoparticle formulations. Nevertheless, these strategies are not without their limitations in the context of drug administration since they may encounter challenges related to the fast elimination of drugs by the mononuclear phagocyte system. Multiple studies have provided evidence suggesting that exosomes possess the ability to traverse the BBB and surmount the immune-privileged nature of the brain, resulting in a decrease in medication elimination. Significantly, exosomes possess the capability to cross the BBB and convey proteins and RNAs into the brain via various routes of administration, such as intranasal, intravenous, intraperitoneal, and intracranial methods. This observation underscores the considerable adaptability and suitability of exosome-mediated drug delivery for the treatment of central nervous system (CNS) disorders.

Therefore, exosomes have been used as a therapeutic intervention for many chronic degenerative neurological disorders. The conditions encompassed in this list are Alzheimer's disease, central nervous system diseases, depression, multiple sclerosis, neurodegenerative disease, Parkinson's disease, post-traumatic stress disorders, and traumatic brain injury. Numerous studies have indicated the clinical efficacy of these treatments, as will be discussed in the following sections.

2.1 | Alzheimer's disease

Alzheimer's disease is a common neurological illness that affects a significant number of people worldwide. Unfortunately, there are few therapeutic options available that have been proven efficacious. The primary strategy for therapeutic intervention in Alzheimer's disease involves the use of inhibitors targeting the enzyme BACE-1 in both neurons and glial cells. This technique effectively reduces the

levels of A-beta, the principal peptide associated with Alzheimer's pathology. The efficacy of drugs utilized for AD is often limiting since they must be taken following early diagnosis. The maintenance of homeostasis and normal brain function relies on cellular interactions within the brain. Exosomes have been identified as significant contributors to cell-to-cell communication among brain cells, facilitating the transfer of information from one cell type to another. It is noteworthy that extracellular vesicles released by mesenchymal stem cells have shown superior efficacy compared with the corresponding parent cells.²⁶ Exosomes possess several advantages upon their specific fusion with target cells. Firstly, they effectively exclude the heterogeneity present in the original cells through a precise selection process. Additionally, their engineering and drug accumulation capabilities enable them to exhibit molecular specificity. Exosomes are capable of inducing effective actions by leveraging variable concentrations of molecules and factors, as well as activating signaling cascades. The strength of their position is enhanced by their integration with a multitude of elements and activities.²⁶

Exosomes have been seen to exert their effects on neuronal and glial cells, facilitating the healing process after traumas and immunological responses.²⁶ The fusion of MSC-Exos has been found to be crucial in facilitating intercellular communication among brain cells, both in healthy conditions and in the context of different pathological states. Several protective effects have been documented, such as the facilitation of synaptic plasticity, provision of nutritional and metabolic assistance, promotion of neuron regeneration, modulation of inflammatory response, and removal of toxic elements.^{27,28} Both the brain and spinal cord may be subject to the induction of traumas and ischemia lesions. In several instances, it has been shown that the recuperation facilitated by modest dosages of MSC-Exos starts via the stimulation of a limited number of receptors. Subsequently, this leads to the phosphorylation of kinases (and other factors), as well as the activation of miRNAs.²⁸⁻³¹ Similarly, it has been shown that MSC-Exos had the ability to alleviate trained immune responses initiated by innate cells, resulting in specific localized tissue healing.³² Furthermore, MSC-Exos have been shown to regulate neuronal responses and mitigate the detrimental effects caused by glial cells, including astrocytes and microglia.^{33,34}

Preclinical research on MSC-Exos has been extensively conducted in murine models, with favorable results (Figure 1). The primary processes behind these effects include the release of secreted mesenchymal stem cell-derived extracellular vesicles (MSC-EVs) including their cargo, which contains crucial components such as nucleic acids, as well as other therapeutic agents. One of the significant impacts associated with AD is the downregulation of BACE-1 and A β levels, accompanied by an elevation in sphingosine-1-phosphate. These events are generated by the protective action of extracellular vesicles derived from MSC-Exos against neuronal damage caused by AD.³⁵⁻³⁷ As a result, the simple injection into the tail vein of animal models has been shown to produce positive effects on the mouse brain including ameliorating cognitive impairment and reducing A-beta aggregation as well as neuronal death.³⁸ Action potentials, dendritic spine augmentations, [Ca²⁺] oscillations, and mitochondrial




FIGURE 1 Therapeutic application of exosomes in AD. A, amyloid-beta; APP, amyloid precursor protein; BACE-1, beta-site amyloid precursor protein cleaving enzyme 1; BaX, Bcl-2-associated X protein; BBB, blood-brain barrier; BCL2, B-cell lymphoma 2; Cur, curcumin; ECVs, extracellular vesicles; MP-MSCs, multipotent mesenchymal stem cells; NEP, neprilysin; NF- κ B, nuclear factor kappa-light-chain enhancer of activated B cells; NFT, neurofibrillary tangles; NSCs, neural stem cells; QU, quercetin; STAT, signal transduction and activator of transcription protein. 1081×540 mm (118×118 DPI). Reprinted with permission from Soliman et al.⁴⁰

reactivations are among the effects that MSC-Exos restore.³⁸ Recent findings have shown MSC-Exos demonstrate great promise as a therapeutic agent and biomarker for AD in humans.^{39–41} The symptoms that were reported, such as the return of homeostatic levels, the preservation of synapses, and enhanced cognition, are comparable to those shown in mice.⁴² In actuality, MSC-Exos prevent neurodegeneration in AD and many other disorders, having favorable effects on brain regeneration and tissue repair.⁴³ A growing body of research indicates that exosomes facilitate cell-cell communication in the brain, which enables them to reduce neuroinflammation and enhance the control of synaptic function. The utilization of exosomes in AD as a therapeutic target or as a biomarker for early identification has been covered in a number of review articles over the years.^{26,40,44–49}

2.2 | Parkinson's disease

The use of exosomes as a therapeutic intervention for Parkinson's disease has garnered significant attention due to the escalating global prevalence of the condition. PD is a progressive movement condition and the second most prevalent neurological illness behind Alzheimer's that impacts around 2% of those aged over 65 years.⁵⁰ Resting tremors, stiffness, slowness, and inability to maintain balance are all examples of motor symptoms. As dopaminergic neurons of the substantia nigra pars compacta die off over time, surviving

neurons develop abnormal structures called Lewy bodies (LBs) and Lewy neurites (LNs) due to the accumulation of the synaptic protein alpha-synuclein (α -syn).⁵¹ The neurotransmitter dopamine (DA) is depleted as a result of these alterations, and an imbalance between acetylcholine (ACH) and DA occurs, resulting in motor difficulties.^{50,52}

RNA molecules, antioxidants, and dopamine agonists are some of the treatments now available to help reduce the motor symptoms associated with Parkinson's disease. Medication for PD nonmotor symptoms include secondary side effects such as sadness, exhaustion, sleeplessness, and dementia.^{53–55} Nevertheless, the efficacy of these medications in halting the ongoing dopaminergic damage is limited due to the reduced efficiency of drug transport caused by the BBB. Similar to several other neurological illnesses, the insufficient permeability of the BBB has been a challenge in the administration of therapeutic interventions for PD in the brain.

In recent years, exosomes have gained significant attention in the field of neuroscience, particularly in the context of neurological disorders such as stroke and brain tumors owing to their ability to cross the BBB. This heightened interest may be attributed to the remarkable biocompatibility shown by exosomes, their capacity to cross the BBB without inducing toxicity, and their ability to selectively target specific sites within the brain. The capacity of exosomes to readily cross the BBB without thrombotic concerns and integrating into neurons and glial cells has also provided the potential for utilizing nano-sized Exos as vehicles for therapeutic medicines or bioactive proteins.

Jarmalaviciute et al.⁵⁶ were one of the first to investigate the neuroprotective capabilities of extracellular vesicles (EVs) in an in vitro model using dopaminergic neurons subjected to 6-hydroxy-dopamine (6-OHDA) induced toxicity. Human exfoliated deciduous teeth were used to grow cultures of mesenchymal stem cells from the dental pulp. Dopaminergic neurons generated from neural stem cells were shielded against 6-OHDA when EVs were present in the growth medium. Thus, exosomes isolated from the tooth pulp prevented 6-OHDA-induced death of 80% of dopamine neurons.⁵⁶

Using a 6-OHDA model of PD in rats, in which 6-OHDA was injected into the medial forebrain bundle (MFB), a subsequent investigation showed that MSC-EVs delivered intranasally enhanced treatment success.⁵⁷ Eight days after unilaterally injecting 6-OHDA into MFB, animals who received daily Exo therapy for 15 days had substantial improvements in gait metrics and motor function. The results demonstrated that Exos from MSCs alleviated Parkinsonian symptoms in a rat model.⁵⁷

Similar outcomes were also seen in a different study when conditioned medium from human dental pulp stem cell cultures were administered in the PD animal model.⁵⁸ In a 6-OHDA rat model of Parkinson's disease, the impact of intravenous hUC-MSC-EV treatment (200 µg, every third day for 8 weeks) was investigated. Exos were shown to penetrate the BBB and integrate into the substantia nigra where they decreased apoptosis, thereby safeguarding dopaminergic neurons and elevating dopamine levels in the striatum. The Exo-treated animals also displayed diminished asymmetric rotation in response to apomorphine treatment at 8 weeks after transplantation.⁵⁹

Haney et al.⁶⁰ created EVs by transfecting macrophages demonstrating promise for the treatment of PD. By introducing a plasmid DNA (pDNA) containing the catalase gene into macrophages using electroporation, the resulting transfected macrophages released EVs that contained various components such as catalase genetic material (pDNA and mRNA), functional catalase, and NF-κB. These EVs were effectively transferred to neurons, leading to a notable decrease in inflammation and the activation of neuroprotective mechanisms in the animal model utilized for PD.⁶⁰

A notable study by Zhuang et al.⁵ observed that exosomes obtained from MSCs loaded with curcumin, a low molecular weight compound with anti-inflammatory properties exhibited a reduction in the activation of microglial cells involved in inflammation. Consequently, these exosomes provided protection against brain inflammation induced by lipopolysaccharides (LPS) in mice.⁵ Additionally, even MSC-Exos injected intraperitoneally were quickly absorbed by cells in the mice's brain, including microglial cells that were both active and in a resting state.⁵

In light of this, these combined results provides encouraging preclinical results regarding the use of exosomes as effective PD treatment alternatives with many review articles written on the topic⁶¹⁻⁶⁵ with Figures 2 and 3 summarizing these findings.

2.3 | Multiple sclerosis

Multiple sclerosis (MS) is an autoimmune neurodegenerative disease of the CNS that causes the brain and spinal cord to develop numerous demyelinating lesions. It affects the lives of an estimated 2.8 million individuals globally.⁶⁶ A diverse range of symptoms, including exhaustion, impairment of bladder and bowel function, visual deficits, difficulties with mobility and coordination, as well as sensory abnormalities. Individuals diagnosed with MS have significant cognitive and emotional alterations that have a profound impact on their overall quality of life. The occurrence of myelin loss or demyelination may result in neuronal disruption, perhaps accompanied by alterations in axonal structure, hence giving rise to a variety of neurological symptoms. In the relapsing-remitting type of MS, these symptoms may transiently manifest for extended periods of time (weeks) during disease exacerbation (known as relapse). Despite the identification of many genetic, metabolic, and environmental variables that influence the development of MS, there is a lack of comprehensive understanding of the specific processes involved in the damage to CNS tissue. Recent research has shown the substantial involvement of circulating extracellular vesicles in several pathological conditions. Exosomes have been identified as a crucial means of intercellular communication, facilitating the exchange of information between two distinct cell types, whether they are located in close proximity or in different regions of the body. In light of recent advancements in comprehending the pathophysiology of MS, investigations have unveiled diverse EVs and their respective functions in the progression of MS. These EVs hold potential as biomarkers for the purpose of monitoring the advancement of the condition. Moreover, their complete understanding will allow exosomes to be used as therapeutic options with the goal of potentially reversing the disease.

The immune-mediated damage occurring in the CNS of individuals with MS is a result of intricate interactions involving several types of immune cells, including NK cells, macrophages, dendritic cells, B cells, and T cells.⁶⁷⁻⁶⁹ Glial cells, including astrocytes and microglia, function as nonclassical immune cells in the pathophysiology of multiple sclerosis in addition to other immune cells.⁷⁰⁻⁷³ Exosomes are thought to be very effective therapeutic options for the treatment of human illnesses, such as MS. Recent research has shown that in animal models of progressive multiple sclerosis, MSC-Exos may aid in the recovery from demyelination.⁷⁴⁻⁷⁷ Notably, MSCs transfer has been studied as a cell treatment for multiple sclerosis for many years, with varying degrees of success. Studies have found that MSC-Exos increased microglial polarization and decreased cytokine production including IL-6, IL-12p70, IL-17AF, and IL-22.⁷⁴⁻⁷⁷

Recently, Casella et al. created a murine microglial cell line that generates EVs loaded with IL-4 that exhibit the endogenous "eat me" signal Lactadherin (Mfg-e8) on their surface. After only one injection of the modified EV, the mice with induced encephalitis showed less neuroinflammation and a lower EAE clinical score, which was linked to an increase in anti-inflammatory markers in phagocytic cells.⁷⁸




FIGURE 2 Graphical abstract highly the potential of exosomes to be utilized as both diagnostic markers as well as therapeutics for the treatment and management of Parkinson's disease. Research today has investigated the incorporation of various small biomolecules into exosomes owing to their ability to cross the blood–brain barrier. Reprinted with permission from Ouerdane et al.⁶⁵

Furthermore, in a recent study by the same research group, it was shown that naturally produced EVs from oligodendrocyte cultures containing myelin antigens found a decrease in the pathophysiology of experimental autoimmune encephalomyelitis (EAE).⁷⁹ In another study, myelin stem cell-derived EVs cocultured with microglia improved myelin healing by recruiting oligodendrocyte precursor cells favoring the recovery of damaged myelin.⁸⁰

A study performed by Williams et al. examined the role of EVs in an EAE model concluded that circulating exosomes exhibit heightened levels during pregnancy.⁸¹ Furthermore, exosomes have been shown to inhibit T-cell activation in an EAE mouse model and aided in the development and migration of oligodendrocyte precursor cells to sites of inflammation.⁸² Thus, currently commercially available exosomes derived from placental amniotic fluids may be an excellent avenue of future research for patients with MS.

2.4 | Depression

Depression is a very widespread mental condition that impacts a substantial global population of over 250 million individuals.⁸³ It may

be treated using a wide range of psychological and pharmaceutical techniques, the majority of which involve adjusting the central nervous system's levels of bioenergy amines.⁸⁴ Unfortunately, the effectiveness of currently available therapeutic techniques toward managing depression is limited mostly because of their inability to cross the BBB.⁸⁵ The pathophysiology of depression is limited by a combination of genetic susceptibility, disrupted monoamine production and function, and altered brain structure and function. Depression is linked to elevated oxidative stress as a result of reactive oxygen species (ROS) production and an imbalance in oxidant and antioxidant signaling.⁸⁶ Because of its increased oxygen use, greater lipid content, and more weakened antioxidative defense, the brain is more vulnerable to oxidative stress.⁸⁷

In a paper titled: "Exosomes: A Novel Therapeutic Paradigm for Treatment of Depression", Amanda Silva and colleagues highlighted the potential use of exosomes for the treatment of depressive disorders.⁸³ Numerous physiological processes, including nerve cell stress, cell-to-cell communication, synaptic plasticity, and neurogenesis, are often seen in the CNS. The ability for exosomes to cross the BBB and transport macromolecules to destination cells is one of their key and unique abilities. The use of exosomes as medicinal




FIGURE 3 Summary of the therapeutic approaches of EXOs in Parkinson's disease. Reprinted with permission from Ouerdane et al.⁶⁵

carriers has been thoroughly investigated in a variety of medical fields, including the treatment of autoimmune diseases and depression will significantly minimize the use of off-target drugs with harmful effects such as antidepressants.

2.5 | Traumatic brain injury

Traumatic brain injury (TBI) affects over 30 million people per year worldwide and caused by a variety of traumatic events including road accidents, falls, exposure to mechanical forces, interpersonal aggression, self-harm, sports injuries, and animal interactions. TBI carries the risk of abrupt or early death in addition to lifelong impairments. Even if the mechanical insult initially results in damage/healing, additional harmful effects are often caused by secondary injury brought on by dysregulated reactions after neuronal death and inflammation. The secretome of MSCs, particularly the exosomes, is primarily responsible for their functional characteristics which have been thoroughly studied as a potential treatment for traumatic brain injury. It has been shown that administering exosomes may improve TBI resulting in a fully recovered brain.⁸⁸ The ability to customize MSC-Exos to transport certain biomolecules of interest to improve their therapeutic efficacy is another benefit of recent technological advancements.

MSC therapy of traumatic brain injury in animal models has shown encouraging results. Zhang et al.⁸⁹ carried out the oldest known research in 2008 with a cohort of seven patients ranging in age from 6 to 55. In this study, bone marrow-derived MSCs (BMSCs) were administered by two different routes: either directly injected into the lesion or administered intravenously. 6 months later, it was discovered that six individuals had significantly improved their neurological function. Unfortunately, one of the patients had two seizure episodes in the first 2 months and had to go back for regular therapy.⁸⁹

Evidence has suggested that exosomes possess the same or better benefits/advantages when compared to their parent stem cells owing to their ability to cross the BBB, however without the dangers of cell therapies. Because of their anti-neuroinflammatory, angiogenic, neuroprotective, and neurogenesis qualities, exosomes play a major role in therapy of TBI (Figure 4).

2.6 | Post-traumatic stress disorder

In 1980, post-traumatic stress disorder (PTSD) was officially identified as a mental/anxiety disorder. Since then, a plethora of information on this disorder's symptoms, epidemiology, evaluation, subtypes, and therapy has surfaced.⁹¹ Particularly in military groups,




FIGURE 4 Schematic of mesenchymal stromal cell (MSC) exosome polarization of microglia from M1 to M2. MSC exosomes are internalized into trauma activated M1 microglia to promote polarization into M2 state, resulting in secretion of anti-inflammatory cytokines which promote brain repair. Adapted from Li et al.⁹⁰ and reprinted with permission from Mot et al.⁸⁸

PTSD is often encountered in conjunction with other mental disorders, traumatic brain injury, and other concomitant symptoms, including depression.^{92–94} Moreover, PTSD is associated with a host of other co-morbidities, including hypertension,⁹⁵ cardiovascular disease,⁹⁶ cardiometabolic diseases,⁹⁷ suicidal thoughts,⁹⁸ in addition to chronic pain.⁹⁹ These highly resemble the disbalance in macrophage polarization toward a pro-inflammatory M1 macrophage phenotype affecting many illnesses. For instance, C-reactive protein (CRP) may indicate PTSD in recently deployed Marines,¹⁰⁰ and cortisol levels in reaction to awakening can predict the outcome of PTSD therapy.¹⁰¹ While not widely studied in much efficacy, exosomes have been proposed as both an ability to serve as a biomarker in PTSD detection and, more importantly, as a therapeutic alternative to current drug-based therapies for anxiety and PTSD once more knowledge is acquired.⁹¹ Since exosomes are also able to lower inflammation, they may also be infused into the body to lower inflammation as a whole and improve macrophage polarization toward the pro-tissue healing/resolution M2 phenotype (Figure 4).

2.7 | Peripheral nerve injury

The most common type of traumatic damage to the nervous system is peripheral nerve injury (PNI) which affects more than a million individuals yearly.¹⁰² The majority of the time, these injuries lead to permanent impairment, which significantly lowers the patients' quality of life.¹⁰³ The peripheral nervous system has the intrinsic capacity to regenerate to a certain extent subsequent to PNI.^{104,105} The surgical technique includes end-to-end anastomosis and is widely regarded as the ideal method for repairing peripheral nerves.¹⁰⁶ It is now acknowledged that autologous nerve grafts (ANGs) are the “gold standard” therapy for PNI.¹⁰⁷ Nonetheless, there are still a number of serious

problems with the therapeutic use of ANGs, including a lack of donor nerve availability, morbidity at the donor site, the creation of painful neuromas, longer recovery durations, and mismatches in length or diameter.^{108,109} Moreover, less than half of the patients who had ANG transplants recovered fully and/or functionally.¹¹⁰

Interestingly, exosomes have been shown to have effects on axonal nerve regrowth. A study conducted by Bucan et al.¹¹¹ provided evidence that MSC-Exos could stimulate neurite extension in cultured dorsal root ganglion neurons. Furthermore, MSC-Exos have been shown to increase the regeneration process after crush damage to the sciatic nerve in vivo. Research conducted by Zhao et al.¹¹² showed that BMSC-Exos had a notable impact on enhancing neurite formation and increasing axon length.

Recent research has shown that exosomes may support and regulate the biological activities of Schwann cells (SCs).¹¹³ In comparison with the vehicle control group, an in vivo study demonstrated that the exosome-treated group that maximized the functioning of SCs had improved axonal regeneration, remyelination, and muscle repair.¹¹³ It was further shown by Zhang et al.¹¹⁴ that umbilical cord MSC-derived exosomes improved vascular regeneration by transporting Wnt4 to endothelial cells, where it activated the Wnt/ β -catenin pathway. Additionally, it has been shown that exosomes derived from bone marrow and adipose MSCs have pro-angiogenic effects during wound healing.^{115,116}

There is a growing body of data suggesting that exosomes have the ability to facilitate vascular regeneration favoring nerve regeneration post-injury. Zhang et al.¹¹⁷ provided evidence that exosomes derived from MSCs have a substantial impact on promoting angiogenesis in the lesion boundary zone.¹¹⁷ Consequently, the administration of Immunosomes effectively enhances the recovery of motor function in rats following traumatic brain injury. Figure 5 summarizes the benefits of exosomes for nerve regeneration.

3 | EXOSOMES AND AUTOIMMUNE DISORDERS

The immune system may lead to many illnesses such as heightened inflammation and autoimmune disorders when it becomes overactive. The field of immunomodulation is poised to emerge as a crucial therapeutic approach for the treatment of several disorders owing to the pivotal role played by the immune system in safeguarding the human body.^{119,120} Symptoms of an overactive immune system include autoimmune diseases and excess inflammation (Figures 6 and 7). While the body needs some degree of inflammation to defend itself, excessive inflammation may have harmful side effects and impede tissue function at the location of the illness.¹²¹ Autoimmune disorders arise due to the inadequate regulation of effector immune responses against the body's own tissues.¹²²

Tregs, or regulatory T cells, are essential for controlling the immune system.¹²² Defects in peripheral tolerance, allergy reactions, and eventual autoimmune disorders arise when Treg numbers or function decline. For instance, it has been demonstrated that an increase in TH1-like regulatory T cells is linked with autoimmune disorders.¹²² This suggests that in autoimmune disorders, maintaining the regulatory T cell's activity is crucial.¹²² Numerous studies have shown that Treg deficits are the root cause of autoimmune disorders such as type 1 diabetes, myasthenia gravis, multiple sclerosis, and rheumatoid arthritis.

Autoimmunity is not only caused by aberrant Treg function. Anti-inflammatory drugs that have the ability to alter inflammatory signals often used to treat inflammation such as nonsteroidal anti-inflammatory medicines (NSAIDs) which block the synthesis of prostaglandins.¹²³ Extensive research is being devoted toward the investigation of signaling pathway inhibitors,¹²⁴ stem cell transplantation,¹²⁵ and various other approaches.

3.1 | Exosomes and immune regulation

Prior to discussing the vast roles of exosomes and their therapeutic effects in the management of various autoimmune disorders, it is important to understand how exosomes affect immune cell regulation.¹²⁶ Exosomes produced from MSCs transport active signaling molecules and immunomodulatory effectors to control immune cell activity and hence mediate immunological suppression, particularly on T cells and macrophages (Tables 1 and 2). In a study titled: "Immunosuppressive Effects of Mesenchymal Stem Cells-derived Exosomes,"¹²⁷ MSC-Exos were clearly shown to regulate immune responses by interacting with immune effector cells. Both physical and chemical factors were shown to affect MSC-Exos.¹²⁷

Lipid molecules found in abundance in MSC exosome membranes may merge with target cells to convey cargoes.¹⁵⁰ The transfer and presentation of antigen peptides, the transport of miRNA, the DNA-driven cyclic GMP-AMP synthase stimulator of interferon genes in recipient cells, and other signaling pathways produced by surface-carrying ligands are possible causes of the immunoregulatory

actions of MSC-Exos.¹⁵¹ MSC-Exos suppress immune cell activation and encourage the production of anti-inflammatory molecules, which lessens inflammatory reactions.¹⁵²⁻¹⁵⁴

MSC-Exos have the ability to modulate immune cells when they are prompted to become active. For instance, when T cells are stimulated by antigen-presenting cells, MSC-Exos may convert T cells into Tregs.¹⁵⁴⁻¹⁵⁶ MSC-Exos, like their parent MSCs, also have an immunosuppressive effect. Crucially, pretreatment may regulate the specificity and immunosuppressive effects of MSC-Exos. For example, MSC-Exos that are treated with a high concentration of pro-inflammatory substances have higher immunosuppressive effects. MSC-Exos have the ability to migrate to the site of injury and impede the immune system. MSC-Exos are injected into the body, and their purpose is to reduce inflammation by migrating to the immune organs, inflammatory tissues, and injured areas. After being injected intravenously, MSC-Exos may be administered to the sites of damage, ingested by immune cells, and found in the spleen and CD260⁺ cells 24 h later.¹⁵⁷ In the past 2 years alone, there are dozens of systematic reviews that have investigated the role of exosomes in autoimmune disorders (Figure 8).¹⁵⁸⁻¹⁶⁷

3.2 | Allergy

An increasingly significant worldwide health and economic burden is caused by allergic reactions and disorders. The search for new therapeutic strategies is necessary since there are no disease-modifying treatments other than particular allergen immunotherapy (AIT), which is not available for all allergy types.¹⁶⁸ A lack of immunological tolerance and the expansion of TH2 cells, which activate B cells to generate IgE responses in response to benign antigens, characterize type I hypersensitivity.^{169,170} Because allergen-specific IgE binds to the high-affinity IgE receptor Fc ϵ RI, it sensitizes mast cells and basophils.¹⁷¹ Upon secondary contact with the allergen, those cells degrade and release inflammatory mediators.¹⁷² Symptomatic treatment options for allergies involve downregulation of the mediators released by mast cells or basophils (e.g., anti-histamine) or aim to downregulate IgE levels, such as the monoclonal anti-IgE antibody Omalizumab.¹⁷³

EVs significantly contribute to the shaping of immune responses in both physiology and disease states. While vesicles secreted by immune cells are often implicated in the allergic process, growing evidence indicates that EVs from nonimmune cells produced in the stroma or epithelia of the organs directly affected by inflammation may also play a significant role. An overview of the mechanisms of allergy to which those EVs contribute, with a particular focus on exosomes is presented in Figure 9.¹⁷⁴

Recent progress in the EV field determined that a thorough understanding of EV biology and function is pivotal for the comprehension of immune-driven diseases, including the pathogenesis of allergy. EVs contribute to asthma pathogenesis via various mechanisms related to both inflammation and pathological remodeling,¹⁷⁵ and there are interesting interdependencies that have been




FIGURE 5 Effects of exosomes on peripheral nerve regeneration. Exosomes can exert therapeutic effects via mediating axonal regrowth, Schwann cell activation, vascular regeneration, and inflammatory regulation, which contribute to providing a favorable microenvironment for peripheral nerve regeneration. Reprinted with permission from Yu et al.¹¹⁸




FIGURE 6 Schematic summary of immunoregulatory cells and cytokines. Reprinted with permission from Suh et al.¹²⁰




FIGURE 7 Schematic of the human body with inflammatory diseases. Reprinted with permission from Suh et al.¹²⁰

observed. Specifically, it has been shown that fibroblast-derived EVs secreted by cells obtained from severe asthmatics increase the proliferation of bronchial epithelial cells in comparison to those in healthy individuals due to a decrease in the TGF- β 2 content.¹⁷⁶ Vice versa, vesicular transfer between epithelial cells and fibroblasts which includes inositol polyphosphate 4-phosphatase type I A (INPP4A) cargo, may regulate inflammation and airway remodeling.¹⁷⁷ Furthermore, Gupta et al. demonstrated that sEV transfer between airway epithelial cells and human tracheobronchial cells promotes the expression of several proteins which may contribute to allergic inflammation and exacerbation of asthma symptoms, that

is, gel-forming mucins.¹⁷⁸ The addition of an allergen source (house dust mite) to the airway epithelial cells culture resulted in DC activation by secreted EVs in vitro and increased airway inflammation in a murine model.¹⁷⁹ While the goal of this overview review article is not to specifically provide major input on the role of exosomes in allergy, a recent review article¹⁷⁴ on this topic has greatly demonstrated all the preclinical animal studies (Table 3) and clinical studies (Table 4), currently underway highlighting the prolific role of small vesicles for the management/treatment of allergic diseases.

Furthermore, various groups have elicited various roles of exosomes in allergic sensitization and inflammation and further discuss

TABLE 1 miRNA involved in the immunosuppressive effects of mesenchymal stromal cell exosomes on macrophages cells.

miRNA	Mechanism	Effect	References
miR-let-7b	Negatively regulated TLR4 and p-p65 expression, suppressed STAT3 and AKT phosphorylation	Orchestrated macrophages plasticity	128
miR-let-7c	–	Downregulated MMP9, TGF- β 1	129
miR-145	Inhibited the expression of MRP1	Promoted phagocytic activity of macrophages	130,131
miR-146a	Inhibited the expression of TRAF6 and IRAK1, reduced phosphorylation of NF- κ B-p65 and I κ B α	Reduced the production of pro-inflammatory factors in macrophages	132
miR-146b	Lessened the release of pro-inflammatory factors and the pro-inflammatory response by NF- κ B	Improved the survival rate of sepsis mice	133
miR-17	Inhibited the activation of NLRP3 inflammatory bodies in macrophages by targeting TXNIP	Reduced the levels of pro-inflammatory factors IL-1 β and IL-18 and improved ALF	134
miR-181c	Reduced TLR4 expression	Reduced pro-inflammatory response in macrophages	135
miR-223	–	Inhibited the expression of TNF- α , IL-1 β , and IL-6 in sepsis-induced macrophages	136
miR-455-3p	Through combination of miR-455-3p and 3'UTR site of PIK3R1 gene	Reduced macrophages infiltration and the secretion of pro-inflammatory factors	137
lncRNA KLF3A51	Downregulated MMP13 expression	Inhibited chondrocyte apoptosis	138

Note: Reprinted with permission.¹²⁷

TABLE 2 miRNA involved in the immunosuppressive effects of mesenchymal stromal cell exosomes on T cells.

miRNA	Mechanism	Effect	References
miR-1246	Regulated the IL-6-gp 130-STAT3 pathway	Reduced the Th17/Tregs ratio in CD4 ⁺ T cells and improved liver dysfunction	139
miR-125a-3p	–	Inhibited the activation of CD3 ⁺ T cells and promoted the Tregs populations	140
miR-126	Reduced NLRP3 inflammatory body activity and NF- κ B-p65 levels	Improved retinal inflammation which caused by hyperglycemia	141
miR-1470	Targeted c-jun mRNA 3' region	Inhibited c-jun expression and promoted the upregulation of P27KIP1 and the ratio of CD4 ⁺ CD25 ⁺ Foxp3 ⁺ Tregs	142
miR-17	Inhibited the transactivation of Jak1	Reduced IL-7 signal transduction and the production of pro-inflammatory factors	143
miR-17-92	Negatively regulated PTEN expression and activated the PI3k/Akt/mTOR signaling pathway and inhibited GSK-3 β activity	–	144
miR-181a	Downregulated c-Fos protein expression the expression of TGF- β and IL-6 and the expression of Foxp3	Improved the area of cerebral infarction and lessened inflammatory infiltration and effectively reduced myocardial I/R damage	145
miR-221	Targeted TIMP2	Reduced the expression of PTEN and p27 proteins, accelerated the proliferation, migration, and invasiveness of GC cells	146
miR-223-3p	Negatively regulated STAT3	Inhibited the secretion of IL-17 by Th17 cells and reduced the expression of IL-1 β and IL-6, increased the ratio of Tregs/Th17	147
miR-29b-3p	Downregulated PTEN	Activated the Akt signaling pathway	148
miR-92a-3p	Inhibited Wnt	Regulated T cells development	149

Note: Reprinted with permission.¹²⁷

the mechanisms by which exosomes could potentially be used in immunotherapeutic approaches for the treatment of allergic diseases. For an excellent review article on the topic by Engeroff and Vogel,¹⁶⁸ the following four strategies have been proposed as potential for utilizing exosomes in the field of allergy:

- Exosomes generated from mast cells that express the IgE receptor Fc ϵ RI have the ability to absorb IgE and reduce systemic IgE levels.
- Tolerogenic exosomes have the potential to inhibit allergic reactions by stimulating regulatory T cells.










FIGURE 8 Therapeutic effects of extracellular vesicles (EV) on inflammatory diseases. Reprinted with permission from Suh et al.¹²⁰

EXTRACELLULAR VESICLES	EV SOURCE vs. EFFECT ON ALLERGIC INFLAMMATION			
				
	Asthma	Allergic rhinitis	Atopic dermatitis Contact dermatitis	Allergic GI disorders Food allergy
sEV 	+ HBECs → neutrophilic inflammation ↑ → monocyte chemotaxis ↑ → Th2 differentiation ↑ → DC maturation ↑ + BALF → proinflammatory response ↑ → leukotriens ↑ + FBs → HBECs proliferation ↑ + AECs → HTBEs: mucus production and inflammation ↑ + AECs → DC activation ↑ + AECs → Th2 bias via IL-33 ↑ - MSCs → ILC2 accumulation ↓ → immune infiltration ↓ → Th2 cytokines ↓ → mucus production ↓ - HBECs → airway remodelling ↓ - AECs → Treg induction ↑	+ NM → Th2 inflammation ↑ - NECs → IL-10+ monocytes ↑	+ KCs → DC activation ↑ - MSCs → IgE levels ↓ → blood eosinophils ↓ → mast cell infiltration ↓ → Th2/proinflammatory cytokines ↓ → epidermal barrier function ↑ → tissue damage ↓ - RBCs → delayed hypersensitivity ↓ → T cell activation ↓ - MSC → contact hypersensitivity ↓ → Tc, Th1 activation ↓ → Treg induction ↑	- IECs → Treg induction ↑ → IL-10 production ↑ → T cell proliferation ↑
m/IEV 	+ HBECs → macrophage inflammation ↑			

+ disease promoting effect; - disease alleviating effect; ↑ increase in a process; ↓ decrease in a process

FIGURE 9 Extracellular vesicles produced by nonimmune cells and their involvement in allergic diseases. Microvesicles and exosomes are the two types of extracellular vesicles, which have been implicated in the pathogenesis of allergic inflammation. There is significant predominance of the exosomal involvement, likely due to the phenotypic characteristics and physical properties of these vesicles, enabling more without damage and entering the circulation for long-distance delivery. Reprinted with permission by Hovhannisyan et al.¹⁷⁴

TABLE 3 Preclinical models using EVs for allergy treatment in animals.

Study title	Conditions	Outcomes	References
Exosomes from bronchoalveolar fluid of tolerized mice prevent allergic reaction	Allergy	BALF-derived exosomes induce tolerance and protection against allergic sensitization in mice	Prado et al. ¹⁸⁰
Pro-inflammatory role of epithelial cell-derived exosomes in allergic airway inflammation	Asthmatic inflammation	IL-13 treated epithelial cell-derived exosomes induce enhanced proliferation and chemotaxis of undifferentiated macrophages in the lungs during asthmatic inflammatory conditions	Kulshreshtha et al. ¹⁸¹
Selective release of miRNAs via extracellular vesicles is associated with house dust mite allergen-induced airway inflammation	Allergic airway inflammation	Selective sorting of Th2 inhibitory miRNAs into airway secreted EVs and increase release to the airway is involved in the pathogenesis of allergic airway inflammation	Gon et al. ¹⁸²
Exosomes derived from human adipose tissue-derived mesenchymal stem cells alleviate atopic dermatitis	Atopic dermatitis	Intravenously or subcutaneously injected human adipose tissue-derived MSC-Exos ameliorate AD in an in vivo mouse model	Cho et al. ¹⁸³
Extracellular vesicles from mesenchymal stem cells prevent contact hypersensitivity through the suppression of Tc1 and Th1 cells and expansion of regulatory T cells	Allergic contact dermatitis	Human umbilical cord-derived MSC-EVs prevent the pathology of contact hypersensitivity by inhibiting Tc1 and Th1 immune responses and inducing the Tregs phenotype in vivo and in vitro	Guo et al. ¹⁸⁴
Small extracellular vesicles derived from human mesenchymal stromal cells prevent group 2 innate lymphoid cell-dominant allergic airway inflammation through delivery of miR-146a-5p	Allergic rhinitis (patients) ILC2-dominant asthma (mouse model)	MSC-sEVs prevent ILC2-dominant allergic airway inflammation through miR-146a-5p	Fang et al. ¹⁸⁵
Exosomes from human adipose tissue-derived mesenchymal stem cells promote epidermal barrier repair by inducing de novo synthesis of ceramides in atopic dermatitis	Atopic dermatitis	Human adipose tissue-derived MSC-Exos effectively repair defective epidermal barrier functions in atopic dermatitis	Shin et al. ¹⁸⁶
Syngeneic red blood cell-induced extracellular vesicles suppress delayed-type hypersensitivity to self-antigens in mice	Delayed-type hypersensitivity contact hypersensitivity	Intravenous delivery of syngeneic mouse red blood cells that is mediated by EVs in a miRNA-150-dependent manner suppresses delayed-type hypersensitivity	Nazimek et al. ¹⁸⁷
Intranasal delivery of MSC-Exos attenuates allergic asthma via expanding IL-10 producing lung interstitial macrophages in mice	Allergic asthma	Intranasally delivered MSC-Exos inhibit allergic asthma in mice	Ren et al. ¹⁸⁸
Epithelial exosomal contactin-1 promotes monocyte-derived dendritic cell-dominant T-cell responses in asthma	Airway allergic models asthma	Epithelial contactin-1 in exosomes is a critical player in asthma pathology	Zhang et al. ¹⁷⁹

Note: Reprinted with permission by Hovhannisyan et al.¹⁷⁴

- Exosomes may encourage TH1-like reactions in response to an allergen.
- Exosomes may alter how IgE facilitates the presentation of antigens.¹⁶⁸

HBECs, human bronchial epithelial cells; BALF, bronchoalveolar lavage fluid; NM, nasal mucus; NECs, nasal epithelial cells; AECs, airway epithelial cells; HTBEs, human tracheobronchial cells; RBCs, red blood cells; IECs, intestinal epithelial cells; KCs, keratinocytes; FBs,

TABLE 4 Registered clinical trial investigating the feasibility of using EVs in allergic patients.

Study title	Conditions	Interventions	Locations	Identifier
Non-coding RNAs analysis of eosinophil subtypes in asthma	Allergic asthma, severe eosinophilic asthma	Biological: Dermatophagoides pteronyssinus allergen Procedure: Blood sampling, Procedure: Bronchial challenge with allergen	Lithuanian University of Health Sciences, Pulmonology Department Kaunas, Lithuania	NCT04542902
Effectiveness of Qufeng Shengshi Fang on treatment of allergic rhinitis	Rhinitis, allergic, perennial	Drug: Qufeng Shengshi Fang and Loratadine, Drug: Loratadine	Peking Union Medical College Hospital Traditional Chinese medicine department Beijing, Beijing, China	NCT02653339
Cohort study of the patterns of microvesicles in the serum of participants with Atopic and non-atopic asthma	Asthma, allergies	Biological: tumor derived microparticles, Drug: cisplatin	The Ohio State University Medical Center Columbus, Ohio, United States	NCT00700726
Influence on human bronchial epithelial cells smoker extracellular vesicles influence on human bronchial epithelial cells	Smokers human bronchial epithelial cells lung pathogenesis biomarkers	Diagnostic test: Bronchoalveolar lavages	Hôpital Saint-Philibert, Lomme, France	NCT03608293
Phase I/IIa study on chitin microparticles in subjects suffering from allergic rhinitis	Seasonal allergic rhinitis	Drug: Chitin microparticles by nasal route	Hammersmith Medicines Research, London, United Kingdom	NCT00443495
Exploratory study of the cutaneous penetration of biodegradable polymeric Microparticles in atopic dermatitis (MicroSkin)	Atopic dermatitis	Drug: Biodegradable and biocompatible polymeric microparticles containing a fluorochrome applied to the skin followed by a skin biopsy	Regional University Hospital Besançon, France	NCT02369432
Impact of narrowband UVB phototherapy on systemic inflammation in patients with atopic dermatitis	Atopic dermatitis	Other: Narrow band UVB treatment (NB-UVB)	The Rockefeller University New York, New York, United States	NCT03083730
Trial on vascular inflammation in atopic dermatitis	Atopic dermatitis vascular inflammation coronary atherosclerosis	Other: FDG-PET scan other: MDCT, other: biopsy and blood collection	Innovaderm Research Inc Montreal, Quebec, Canada	NCT02926807
Role of macrophage in immune modulation by mesenchymal stem cell derived exosome in asthma	Respiratory disease	Primary indicator: PD-L1, immunosuppression capacity of regulatory T cell	Sun Yat-Sen Memorial Hospital, Sun Yat-Sen University	ChiCTR2000031122

Note: Reprinted with permission by Hovhannisyan et al.¹⁷⁴

fibroblasts; MSCs, mesenchymal stem cells. ↑ increase in a process; ↓ decrease in a process; + disease promoting effect; – disease alleviating effect. Reprinted with permission.¹⁷⁴

3.3 | Atopic dermatitis and contact allergy

Atopic dermatitis (AD) is a chronic inflammatory skin disease that is characterized by severe pruritus, eczematous cutaneous lesions, and disruption of the epidermal barrier. The pathophysiology of AD is multifaceted, with Th2 cells and ILC2 being the primary mediators

of cytokine production.¹⁸⁹ Innate, Th17, and Th22 components are also involved.¹⁹⁰ However, delayed-type hypersensitivity to minor contact allergens, such as contact allergy and contact sensitization, is rather prevalent. An allergy to anything touched repeatedly or for extended periods of time is called contact dermatitis.^{191–193} The study conducted by Nazimek et al. demonstrates that the introduction of syngeneic mouse red blood cells through intravenous administration results in the generation of EVs. These EVs exhibit the ability to inhibit the immune response known as directed delayed-type hypersensitivity, and this effect was shown to be dependent on the presence of miRNA-150. More specifically, the EVs derived

from syngeneic mouse red blood cells reduce the activation of T cells and promote their programmed cell death.¹⁸⁷ Similarly, it has been shown that EVs generated from human umbilical cord MSCs may both prevent and treat the pathophysiology of touch hypersensitivity in mice.¹⁸⁴ In particular, these EVs suppressed the production of TNF- α and IFN- γ , induced Tregs, and decreased the amount of released IL-10 in both CD8⁺ cytotoxic cells and CD4⁺ Th1 cells.¹⁸⁴ While this field remains in its infancy in terms of clinical applicability, future understanding of exosomes, including their ability to modulate the immune system, should shed light on potential avenues of future uses of exosomes for the treatment of AD and contact allergy.

3.4 | Lupus

Systemic lupus erythematosus (SLE) is a common autoimmune connective tissue disease with unclear etiology and pathogenesis. It is characterized by the excessive synthesis of pathogenic autoantibodies and immunological complexes, aberrant immune cell activation, and organ-wide effects.¹⁹⁴ The immunological problems that

accompany the complicated etiology and pathophysiology of SLE include aberrant T cell, mononuclear macrophage cell, and B cell growth, differentiation, activation, and malfunction. Chronic inflammatory conditions and autoimmune diseases ultimately lead to harm to tissues and organs.^{195,196}

The most frequent and serious organ impairment in SLE is lupus nephritis.¹⁹⁷ Currently, SLE is conventionally treated with glucocorticoids and immunosuppressants. Nevertheless, a significant number of patients with refractory conditions continue to present challenges in achieving clinical remission, resulting in elevated death rates. SLE patients have significant economic and psychological challenges.¹⁹⁸ Therefore, there is still a pressing need to develop solutions to the issue for SLE patients.

In a paper by Yang et al. titled: "Immunomodulatory Effect of MSCs and MSCs-Evs in Systemic Lupus Erythematosus"¹⁹⁹ it has been shown that MSCs and the exosomes they produce are useful in controlling innate and adaptive immunity, which are engaged in a variety of pathological and physiological processes and support immunological homeostasis in sickle cell disease.¹⁹⁹ Figure 10 gives new insight into the pathophysiology of SLE and directs biological




FIGURE 10 Composition and mechanism of immunological tolerance of MSC-EVs in systemic lupus erythematosus. MSC-EVs are spheroidal shaped and two-layer lipid particles containing various types of protein, lipids, DNAs, non-coding RNAs, miRNAs, and mRNA, which cause genetic information exchange by various of signal pathway and reprogramming of the recipient cell. MSC-EVs can suppress the differentiation and proliferation of B cell by PI3K-AKT pathway and reduce production of IL-10. Similarly, T cells play the suspensive role on the proliferation and maturation, while reducing the production of Th17 and Th1 and improve function of Treg and Th2 through the TGF- β /NF- κ B pathway. EVs can suppress the proliferation and maturation of DCs and induce tolerable DCs with low expression of costimulatory makers. Macrophages can transform to anti-inflammatory M2 phenotype after treating by MSC-EVs through the PI3K/AKT pathway. EVs can suppress the proliferation, differentiation, and cytotoxicity of NK cells in a TGF- β -dependent manner. MSC-EVs play an important role in the pathogenesis of autoimmune diseases, including SLE, graft-versus-host disease, and experimental autoimmune encephalomyelitis. Reprinted with permission from Yang et al.¹⁹⁹

treatment by illuminating all the underlying mechanisms and immunomodulatory effects of MSC/MSC-EVs in SLE.¹⁹⁹

3.5 | Rheumatoid arthritis

Rheumatoid arthritis (RA) is a chronic inflammatory autoimmune disease that causes cartilage and bone damage with a considerably high social and economic impact. Severe systemic consequences, including cardiovascular events, as well as lifelong disability and function loss, are all possible outcomes of RA. The most frequent symptoms are related to symmetrical knee, foot, wrist, and hand discomfort and edema.²⁰⁰ Approximately 0.5%–1% of the global population is affected with RA, which may affect people of any age, but it is most common in those in their third, fourth, and fifth decades.²⁰¹ Even if disease remission may be achieved with strict management and treat-to-target tactics, as recommended by current guidelines, RA is still regarded as an incurable condition.²⁰¹ More than half of patients have disease flares, which significantly increase the risk of radiographic damage, worse quality of life, disability, use of health-care services, and expenses.^{202,203} At present, there are no markers for tailored treatment or predictors of therapeutic response. However, RA causes irreparable damage to joints or organs as well as persistent inflammation if treatment is not received. Therefore, the most prevalent symptoms in RA patients are increased disability and shorter lifetime.^{204,205}

Bone marrow mesenchymal stem cells (BMSCs) were tested originally and demonstrated significant improvements in symptoms of refractory patients in two clinical studies.^{206,207} These outcomes are in line with the majority of BMSC-based arthritic models. Consequently, BMSC-derived exosomes (BMSC-Exos) have been suggested as a potential treatment for RA because they have effects similar to those of their parent MSCs without compromising the immune system.²⁰⁸

The function of BMSC-Exos in RA models was first studied by Stella et al. who demonstrated that these Exos prevented T and B lymphocyte proliferation and generated Treg and IL-10-expressing regulatory B cells dose-dependently to mitigate experimental RA.²⁰⁸ The immunomodulatory properties of BMSC-EVs have shown positive outcomes in various animal models of osteoarthritis.^{209,210} Moreover, BMSC-EVs have been shown to contribute to angiogenesis and bone/cartilage regeneration.^{209,211}

The primary mechanism by which MSC-EVs affects the illness is by the transfer of miRNAs,²¹² thus inhibiting the cyclin I/ATM/ATR/p53 signaling pathway.²¹³ In addition, EVs were shown to overexpress a number of miRNAs that were previously downregulated in synovial tissue but were effective in treating inflammatory arthritis. One of them, miR-192-5p, regulates the immunological response by targeting Ras-related C3 botulinum toxin substrate 2, therefore delaying the inflammatory response in rat models of collagen-induced arthritis.²¹⁴ It was shown that miR-320, which is secreted from BMSCs via exosomes, selectively downregulates the chemokine ligand CXCL9, hence preventing RA-FLS activation, migration, and

invasion.²¹⁵ Matrix metalloproteinase 14 and vascular endothelial growth factor (VEGF) are two proteins involved in angiogenesis that are downregulated by exosomal miR-150-5p.^{216,217}

Although BMSC-EVs have a beneficial impact on RA, the mechanisms by which they enhance immune modulation and anti-inflammatory responses in RA remain poorly understood and hence need more study. Additionally, umbilical cord mesenchymal stem cells and EVs produced from AMSCs (AMSC-EVs)²¹⁸ have been proposed as a therapy for RA.^{219,220} Figure 11 highlights various mechanisms by which MSC-EVs impact RA.

3.6 | Osteoarthritis

Osteoarthritis (OA) is a long-term degenerative joint disease that is characterized by synovial inflammation, subchondral bone sclerosis, articular cartilage deterioration, and the growth of osteophytes.^{158,222} Joint pain, swelling, and abnormalities are the primary clinical signs. The loss of articular cartilage is the primary degenerative alteration.²²³ Joint pain is brought on by dysfunction, but it also has an impact on the patient's quality of life and ability to sleep.²²⁴

A recently filed clinical study (NCT05060107) aims to provide MSC-Exos intravenously for the purpose of managing OA. A total of 10 participants were enrolled in a Phase 1 clinical study and thereafter monitored for duration of up to 12 months in order to validate the safety of MSC-Exos as a therapeutic intervention for OA.

Many preclinical studies exist supporting the use of exosomes. A study conducted by Zhang et al.²²⁵ revealed that the use of chitosan hydrogels as carriers loaded with MSC-Exos had the potential to enhance the stability of proteins and miRNAs inside MSC-Exos. Liu et al. conducted a study in which they incorporated MSC-Exos into a hydrogel glue using the photoinduced imine crosslinking (PIC) technique. These results suggested that the MSC-Exos is a promising strategy for treating articular cartilage abnormalities since it may stimulate repair and regeneration of damaged cartilage.²²⁶

Additionally, anti-inflammatory molecules with low bioavailability such as curcumin may also be encapsulated using MSC-Exos as carriers. Consequently, curcumin has been loaded into MSC-Exos as a drug delivery system for curcumin at the nanoscale. The phosphorylation of p-38MAPK and Erk1/2, PI3K/Akt, caused by IL-1 β may be reduced by curcumin-encapsulated MSC-Exos, which in turn reduces the activity of pathways involving these kinases.²²⁷ Accordingly, the curcumin-encapsulated MSC-Exos may also cause target cells to express miR-126-3p more highly and lessen the IL-1 β -induced OA chondrocyte breakdown.²²⁷ While the therapeutic benefits of MSC-Exos have shown promise in both clinical and animal settings, it is essential to further increase research to elucidate the underlying mechanism and ensure the safety of MSC-Exos prior to their widespread clinical use.

The chondroprotective and anti-inflammatory effects of murine BMSC-Exos have been demonstrated in the treatment of a mouse model of collagenase-induced osteoarthritis (CIOA).¹⁵⁸ These effects were achieved by (1) restoring the homeostatic state of




FIGURE 11 Schematic view of the potential mechanisms of mesenchymal stem cell-derived EVs in the treatments of rodent models of RA. EVs from different sources of MSCs show efficacy in the treatment of RA models. These EVs mainly show the immunosuppressive function of inhibiting T-cell proliferation, downregulating Ig production and decreasing pro-inflammatory factors levels in vivo, thus attenuating clinical signs of paw swelling as well as histopathological indicators of bone and cartilage erosion and pannus formation. Several contents (including miR-34, TGF-β1, and IL-1ra) have been indicated to be associated with these functions. AMSC-EVs, adipose tissue mesenchymal stem cell-derived extracellular vesicles; BMSC-EVs, bone marrow mesenchymal stem cell-derived extracellular vesicles; CKs, cytokines; FLS, fibroblast-like synovial cells; FOXP3: fork-head box protein P3; Ig, immunoglobulin; IL, interleukin; IL-1ra, IL-1 receptor antagonist; miR-34, microRNA-34; MSC-EVs, mesenchymal stem cell-derived extracellular vesicles; PGE2, prostaglandin E2; RA, rheumatoid arthritis; ROR-γ, retinoic acid receptor-related orphan receptor γ; TGF-β: tumor growth factor beta; Th17, T helper 17; TNF-α, tumor necrosis factor alpha; Tr1, T regulatory type-1; Treg, regulatory T cells; UC-MSC-EVs, umbilical cord mesenchymal stem cell-derived extracellular vesicles. Reprinted with permission from Miao et al.²²¹

chondrocytes, (2) preventing chondrocyte apoptosis, and (3) promoting a shift in macrophages toward an anti-inflammatory phenotype.²¹⁰ Furthermore, by switching synovial macrophages from M1 to M2 and minimizing the breakdown of articular cartilage, a recent study showed that intra-articular injection of BMSC-Exos might reduce OA damage in rat models.²⁰⁹ In vitro TNF-α-stimulated OA, BMSC-EVs have also been suggested to enhance cartilage regeneration and exhibit anti-inflammatory properties.²²⁸ Zhu et al.²²⁹ also demonstrated that by promoting chondrocyte migration and proliferation, transplanting-induced pluripotent stem cell-derived MSC-Exos had a significant therapeutic benefit in the CIOA animal model when compared to MSC-Exos obtained from synovial membranes.

Overexpression of miR-140-5p in MSC-Exos has also been shown to promote cartilage tissue regeneration and reduce knee joint damage in a rat model of osteoarthritis.²³⁰ It has recently been shown that LPS-primed MSC-EVs promote knee cartilage healing in an OA animal model by increasing chondrocyte proliferation, migration, and decreasing apoptosis.²³¹ Wang et al.²³² showed that in a mouse model of OA, MSC-Exo-miR-155-5p increased chondrocyte proliferation and migration, decreased apoptosis, modulated production of cartilage ECM, that resulted in a final increase in cartilage regeneration. By restoring a healthy equilibrium between ECM production and degradation, intra-articular infusion of embryonic stem

cell (ESC)-MSC-Exos alleviated cartilage injury and decreased matrix degradation in a mouse model of medial meniscus instability.²³³ In addition, BMSC-Exos alleviated knee discomfort in arthritic rats and had regenerative effects on cartilage damage and ECM synthesis.²³⁴ Recent research by Jin et al.²³⁵ discovered that miR-26a-5p in particular, which is produced in BMSC-Exos, may reduce OA damage in vivo by inhibiting PTGS2. Polydactyly BMSC-Exos have also exhibited better capacity in promoting chondrocyte regeneration and attenuating OA in a mouse model via BMP4 signaling pathway.²³⁶ In another animal experiment by Liu et al.,²³⁷ MSC-Exos was administered into CIOA mice that led to an increase in chondrocyte proliferation and a decrease in chondrocyte apoptosis. Another study found that MSC-Exos reduced inflammation, restored matrix homeostasis, improved proliferation, increased matrix synthesis, and decreased apoptosis in an OA rat model of the temporomandibular joint (TMJ), therefore lowering pain and promoting TMJ regeneration.²³⁸ Using a CIOA-induced mouse model, Mao et al.¹⁴⁹ found that BMSC-Exo-derived miR-92a-3p inhibited cartilage degradation and maintained homeostasis via regulating WNT5A expression. Chen et al.²³⁹ implanted 3D-printed ECM/GelMA scaffolds with BMSC-Exos subcutaneously in rabbits with osteochondral defect. These scaffolds improved early OA treatment by rectifying cartilage mitochondrial dysfunction, increasing chondrocyte migration,

and polarizing synovial macrophages toward M2.²³⁹ Furthermore, Wu et al.²⁴⁰ demonstrated that intra-articular injections of MSC-Exos reduced cartilage damage in OA rats by inhibiting chondrocyte apoptosis, preserving cartilage homeostasis, and blocking the mTOR-autophagy pathway. Additionally, ADSC-Exos have been shown in an in vitro investigation to lessen inflammatory responses and oxidative stress in OA osteoblasts.²⁴¹ Tables 5 and 6 as well as Figure 12 display all recent publication highlighting the use of exosomes and EVs for the management of OA.²⁴²

3.7 | Inflammatory bowel disease

Inflammatory bowel disease (IBD) is an autoimmune disease including ulcerative colitis (UC) and Crohn's disease (CD) that significantly affects the quality of life in these patients.^{158,263,264} MSC-EVs have recently been investigated in various IBD models. Heidari et al. (2021) assessed the efficacy of ADSC-Exos in a mouse model of acute colitis caused by dextran sulfate sodium (DSS).²⁶⁵ In this animal model, it was demonstrated that an increase in Treg population and a decrease in inflammatory cytokines alleviated colitis.²⁶⁶ Additionally, in a DSS-induced IBD animal model, intravenous injection of ADSC-Exos enhanced epithelial regeneration, decreased inflammation, and supported functional recovery while preserving intestinal barrier integrity.²⁶⁶ Additionally, a recent study by Li et al.²⁶⁷ found that in a mouse model of DSS-induced colitis, ADSC-Exos as well as the parent ADSCs both exhibited comparable anti-inflammatory and immunosuppressive effects. Specifically, exosomes have been shown to regulate the mechanisms involved in inflammation, hence exhibiting potential therapeutic benefits in mice with DSS-induced colitis.²⁶⁸

Exosomes have also been shown to mitigate DSS-induced IBD in mice by means of ubiquitination²⁶⁹ and the control of IL-7 production in macrophages.²⁷⁰ Tolomeo et al.²⁷¹ conducted a comparative investigation including MSCs and MSC-EVs. The purpose of their research was to compare the effectiveness of two different therapy strategies for treating IBD in a DSS-induced colitis mouse model.²⁷¹ The results of their study demonstrated that cytokine-primed MSC-EVs exhibited a significant reduction in intestinal fibrosis and angiogenesis, while also enhancing the functionality of the intestinal epithelium. These effects were achieved by the modulation of macrophage polarization, specifically shifting from the M1 phenotype to the M2 phenotype, as well as an increase in the number of Treg cells.^{272,273} Moreover, rats exposed to 2,4,6 trinitrobenzenesulfonic acid (TNBS) developed colitis, but miR-146a-containing BMSC-EVs protected them by inhibiting IL-1 receptor-associated TNF receptor-associated factor 6 (TRAF6) and kinase 1 (IRAK1).¹³² In a study by Tian et al.,²⁷⁴ olfactory ecto-MSC-Exos dramatically reduced the degree of disease by downregulating Th17 and Th1 populations and upregulating Tregs in a mouse model of DSS-induced colitis. Furthermore, Duan et al.²⁷⁵ investigated the possibility that EVs derived from human placental MSCs might alleviate colitis produced by TNBS in mice by inhibiting inflammation and oxidative stress. Lastly,

overexpressing telomerase and hypoxia-inducible factor 1-alpha in MSCs primed with pro-inflammatory stimuli has been found to provide greater therapeutic advantages in mice with TNBS-induced colitis. According to their research, the EVs might reduce inflammation and fibrosis by producing more M2 macrophages.²⁷⁶ Table 7 highlights the use of exosomes in various autoimmune diseases including IBD.

3.8 | Type 1 diabetes

In type 1 diabetes mellitus (T1DM), T lymphocytes destroy pancreatic beta cells resulting in this complicated and hard to treat autoimmune disease.²⁷⁷ Environmental, immune system, and genetic variables all impact type 1 diabetes.^{278,279} A study by Nakano et al.²⁸⁰ assessed the efficacy of injecting rat BMSC-Exos intravenously in mice with diabetes treated with streptozotocin (STZ). In a second study utilizing the STZ-diabetic mouse model, ADSC-Exos was administered intraperitoneally to one group, whereas the other group received no treatment.²⁸¹ The autoimmune responses were significantly reduced in the treated group which led to improved regulation of IFN- γ , IL-17, IL-4, IL-10, and TGF- β . Furthermore, Shigemoto-kuroda et al. discovered that MSC-EVs could inhibit Th1 and Th17, thereby significantly increasing plasma insulin concentrations and effectively delaying the onset of T1DM in order to avoid islet inflammation.²⁸²

3.9 | Osteoporosis

Osteoporosis (OP) is a common complication in rheumatic diseases characterized by an imbalance between bone resorption and bone formation thereby contributing to the long-term deterioration of bone tissue including an elevated risk of bone fracture.^{158,283} MiR-150-3p from BMSC-Exos was investigated by Qiu et al.²⁸⁴ to reduce OP in rats. In 2020, Yang et al.²⁸⁵ demonstrated that the miR-1263 produced from huc-MSC-Exos exhibited anti-apoptotic properties in osteoporosis caused by hind limb unloading (HLU) via the Mob1/Hippo axis. In a mouse model, Huc-MSC-Exos have also been shown to play a role in osteoporosis inhibition and osteogenic induction.²⁸⁶ Furthermore, it was shown that ADSC-Exos relieved diabetic OP via inhibiting the NLRP3 inflammasome in rat osteoclasts.²⁸⁷ A 2020 publication showed that delivering miR-22-3p generated from BMSC-EVs was a useful strategy to enhance osteogenic differentiation in mice with ovariectomized-induced OP.²⁸⁸ The prevention of OP was also shown in association with cyclic mechanical stretch (CMS)-modified BMSC-Exos which function by lowering the activation of the NF- κ B signaling pathway.²⁸⁹ While the majority of studies to date focus on in vitro and in vivo work using exosomes for the management of osteoporosis,²⁹⁰⁻²⁹⁵ there remains great interest to translate these findings toward clinical practice. Tables 7 and 8 highlights additional studies investigating exosomes on autoimmune disorders.

TABLE 5 Therapeutic application of MSCs-derived EVs-miRNAs in OA treatment.

Cells	Isolation	Agent	Loading	Quantification	Animal OA model	Biological function	References
BMSCs	UC	miR-126	Curcumin	In vitro: Not available	None	In vitro: Reverse IL-1 β induced catabolic responses of chondrocytes	Li et al. ²²⁷
SMSCs	UC	circRNA3503	Melatonin/plasmid transfection	In vitro: Not available In vivo: 100 μ L sEVs (10^{11} sEV vesicles/mL) per week, 4 weeks	ACLT model with SD rats	In vitro: Rescue cells from the destructive effect of IL-1 β In vivo: Protect cartilage	Tao et al. ²⁴³
SMSCs	UC	miR-129	Mimic transfection	In vitro: Not available	None	In vitro: Reduce chondrocytes injury and ECM degradation	Qiu et al. ²⁴⁴
BMSCs	UF	miR-216a	Hypoxia/lentivirus transfection	In vitro: Not available In vivo: Not available	DMM model with SD rats	In vitro: Promote chondrocytes proliferation, migration, inhibit apoptosis In vivo: Promote cartilage regeneration	Rong et al. ²⁴⁵
SMSCs	UC	miR-31	Mimic transfection	In vitro: 10mg EVs for 24h In vivo: 5mL EV particles per mL, from the 5th to the 8th week after operation	ACLT model with C57 mice	In vitro: Promote proliferation and migration of chondrocytes In vivo: Alleviate cartilage damage and inflammation	Wang et al. ²⁴⁶
BMSCs	UF	miR-136	Agomir transfection	In vitro: Not available In vivo: 100 μ L of 10^{11} particles/mL for 1h	ACLT model with C57 mice	In vitro: Promote migration of chondrocytes In vivo: Reduce cartilage degeneration	Chen et al. ²⁴⁷
BMSCs	UC	miR-124 miR-143	Curcumin	In vitro: Not available In vivo: Not available	OA model with mouse	In vitro: Reduce apoptosis of chondrocytes In vivo: Attenuate OA	Qiu et al. ²⁴⁸
ADMSCs	UC	miRNAs	IFN- γ	In vitro: Not available	None	In vitro: Anti-inflammatory of inflamed chondrocytes and macrophages	Ragni et al. ²⁴⁹
ADMSCs	UC	miR-145 miR-221	None	In vitro: 400 μ g/mL exosomes for 48h	None	In vitro: Promote chondrogenic regeneration	Zhao et al. ²⁵⁰
UC-MSCs	UC	miR-381-3p	Kartogenin	In vitro: 20 μ g/mL for 48h In vivo: Not available	Full-thickness Cartilage defects with rabbit	In vitro: Enhance chondrogenesis In vivo: Promote cartilage repair	Jing et al. ²⁵¹
BMSCs	EQ	miR-26a-5p	Lentivirus transfection	In vitro: 2 μ g exosomes for 48h In vivo: 250ng/5 μ L EXO per week, 8 weeks	OA model with Wistar rats	In vitro/in vivo: Alleviate damage of synovial fibroblasts In vivo: Retard OA damage	Jin et al. ²³⁵
SHEDs	UC	miR-100	Mimic transfection	In vitro: Exosomes for 2h	None	In vitro: Anti-inflammatory in temporomandibular joint chondrocytes	Luo et al. ²⁵²

TABLE 5 (Continued)

Cells	Isolation	Agent	Loading	Quantification	Animal OA model	Biological function	References
IMSCs	EQ/UC	miR-100	Antagomir transfection	In vitro: exosomes (1, 5, or 10×10^8 particles/mL) for 24 h In vivo: $10 \mu\text{L}$ exosomes (10^{10} particles/mL) for 4 weeks or 6 weeks (twice a week)	DMM model with rat	In vitro: Inhibit the chondrocyte apoptosis and balance the anabolic and catabolic processes In vivo: Protect cartilage and ameliorate gait patterns	Wu et al. ²⁴⁰
BMSCs	EQ	miR-320c	Mimic transfection	In vitro: exosomes for 48/72 h	None	In vitro: Enhance chondrogenesis	Sun et al. ²⁵³
MSCs	SZC	miR-135b	TGF- β 1	In vitro: $10 \mu\text{g/mL}$ exosomes, 3 days In vivo: $100 \mu\text{L}$ exosomes (1×10^{11} particles/mL) per week for 12 weeks	In vivo: DMM model with rat	In vitro: Promote chondrocytes proliferation In vivo: Promote cartilage repairment	Wang et al. ²⁵⁴
SMSCs	SZC	miR-140	Mimic transfection	In vitro: 10×10^{11} particles/mL of Exos for 24 h In vivo: 100 mg exosomes per 100 mL per week until 12 weeks	In vivo: DMM model with rat	In vitro: Enhance proliferation and migration of chondrocytes In vivo: Prevent OA	Tao et al. ²³⁰

Note: Reprinted with permission from Shang et al.²⁴²

4 | TREATMENT OF DAMAGED ORGANS

This section is dedicated to the treatment of various organs that have benefited from exosome therapy. Today, a multitude of studies have tackled the effects of exosome therapy on the treatment and aid of various organs including heart (such as myocardial infarctions and stroke), kidney, liver, and ovaries.

4.1 | Heart

Cardiovascular disease (CVD) is a significant global health issue. Based on statistics from the US Center for Disease Control and Prevention, CVDs are the primary cause of mortality in the United States.³²¹ Furthermore, according to an epidemiological analysis in China, the prevalence of CVD continues to rise.^{322,323} As a result of societal and economic developments (especially within the aging population), the acceleration of urbanization, and changes in national living differences over the past few decades, unhealthy lifestyles are becoming increasingly common, and the effects of CVD risk factors on the health of residents are becoming more prominent/significant.

In the cardiovascular system, stem cells, vascular cells, progenitor cells, cardiomyocytes, and endothelial cells are linked and communicate with one another via exosomes. Exosomes have the ability to stimulate angiogenesis, suppress ventricular remodeling, enhance cardiac function, suppress local inflammation, and modulate immunological responses. Exosomes also have a significant impact on the growth, harm, and illness of the cardiovascular system. Nevertheless, being a relatively new field of medicine, clinical scientists are currently investigating the function of exosomes and their processes with the long-term goal of facilitating cardiac function and repair comprehensively.³²⁴

As a result, exosomes are crucial to the cardiovascular system and are involved in many different CVDs, including heart failure (HF), acute myocardial infarction (AMI), atherosclerosis (AS), and myocardial ischemia-reperfusion (I/R) damage as highlighted below.³²⁴

4.1.1 | Exosomes from different cell sources and cardiovascular signal transduction

Exosomes are effective disseminators of biological signals related to myocardial function (Table 9). In the heart, crosstalk between several cell types via EVs mediates local communication. EVs secreted by different heart cell types may also affect the same cells that release vesicles in an autocrine manner.³²⁴ Remote communication between the heart and other organs (e.g., the kidneys, brain, and bone marrow) is mediated by EVs secreted from the myocardium entering systemic circulation. Local and distant communication comprises endocytosis, membrane fusion, or gap junction-mediated transfer, and the exchange of nucleic acids, lipids, and proteins by exosomes.³²⁴ The mechanism by which exosomes work also includes receptor-mediated signal transduction, which regulates transcription and

TABLE 6 Therapeutic application of non-MSCs-derived EVs-miRNAs in OA treatment.

Cells	Isolation	Agent	Loading	Quantification	Animal model	Biological function	References
Chondrocytes	UC	miR-221	Mimic transfection	In vitro: 5×10^8 particles, 2 days or 2 weeks	None	In vitro: Inhibit bone formation	Shang et al. ²⁵⁵
Macrophage	UC	miR-1246	Lipopolysaccharide	In vitro: 1×10^9 p/mL of sEVs for 48 h In vivo: 50 µg/µL, one time	CFA-induced TMJOA model with SD rats	In vitro: Promote inflammation In vivo: Promote TMJ inflammation	Peng et al. ²⁵⁶
Fibroblast-like synoviocytes	UC	miR-126	Mimic transfection	In vitro: Not available In vivo: 40 µL of 500 µg/mL per week, until 10 weeks	ACLT + MMx OA model with SD rats	In vitro: Suppress chondrocytes inflammation and apoptosis In vivo: Maintain subchondral bone structure and suppress synovial inflammation-mediated cartilage degeneration	Zhou et al. ²⁵⁷
Osteoblasts	UC	miR-210	Mimic transfection	In vitro: 20 µg/mL exosomes	None	In vitro: Promote cartilage degeneration	Wu et al. ²⁵⁸
Osteoclasts	UC	let-7a	None	In vitro: Not available	None	In vitro: Promote chondrocytes hypertrophy	Dai et al. ²⁵⁹
Chondrocytes	UC	miR-8485	Mimic transfection	In vitro: Not available	None	In vitro: Promote chondrogenic differentiation of BMSCs	Li et al. ²⁶⁰
Chondrocytes	UC/UF	miR-449a	IL-1β	In vitro: Not available In vivo: 10^9 particles in 5 µL vehicle per week, 8 weeks	DMM OA model with mice	In vitro: Inhibition autophagy In vivo: Aggravated synovitis and cartilage erosion	Ni et al. ²⁶¹
Chondrocytes	UC	miR-95	Mimic transfection	In vitro: Not available	None	In vitro: regulate cartilage development and homeostasis	Mao et al. ²⁶²

Note: Reprinted with permission from Shang et al.²⁴²




FIGURE 12 Schematic diagram of potential factors that affect the management of EVs-miRNAs, which might further affect its clinical application in human OA. Reprinted with permission from Shang et al.²⁴²

post-transcriptional processes in target cells. Because of these abilities, exosomes have proven to be effective disseminators of biological signals, which are involved in multiple processes that regulate cardiac function under normal physiological and pathological cardiac conditions.³²⁴

4.1.2 | Myocardial infarctions and injury

The restricted capacity of adult mammalian cardiomyocytes to undergo cell division in response to cardiac injuries is a significant factor toward the development of detrimental fibrotic and myocardial remodeling, eventually leading to heart failure. The persistent prevalence of heart failure-related illness and mortality necessitates the focused efforts by researchers globally to develop effective treatments for cardiac restoration.³²⁵ Following heart damage, this lack of proliferation leads to pathological healing processes and fibrotic scarring.^{326–328}

The majority of studies on stem cell therapies has demonstrated enhanced cardiac function and vascularization along with decreased infarct size despite the transplanted cells dying quickly in the damaged myocardium. This suggests that paracrine mechanisms play the most significant role in cardiac healing. It has been noted that all progenitor and stem cells used in cell-based therapies secrete substances that affect nearby cells in a paracrine manner.

These exosomes are grouped around pro-angiogenic, pro-survival, proliferative, and immunogenic factors.^{329–335} Noteworthy, many miRNAs have now been discovered as either being in pro-repair or pro-pathological state, as shown in Figure 13.³³⁶

Exosomes derived from various origins have the ability to mitigate cardiac ischemia/reperfusion (I/R) damage through the modulation of inflammation, autophagy, and apoptosis. Chen et al.³²³ discovered that exosomes obtained from BMSCs containing miR-125b enhanced cell survival and decreased the rate of apoptosis by specifically targeting SIRT7. As a result, this reduces the size of myocardial infarction and inhibits harm caused by myocardial ischemia/reperfusion.³²³ In another study, Zhao et al.³³⁷ discovered that MSC-Exos employed miR-182 to alter the polarization of macrophages and suppress cardiac inflammation, therefore mitigating myocardial ischemia-reperfusion (I/R) damage in mice. Youn et al.^{338–340} discovered that administering exosomes produced from cardiac progenitor cells (CPCs) directly into the animal heart may effectively inhibit cell death in acute ischemia/reperfusion damage models. Zhang and Zhang³⁴¹ discovered that serum exosomes have the potential to enhance hemodynamics and decrease apoptosis by stimulating the PI3K/AKT signaling pathway, hence reducing I/R damage. Additionally, it was observed that serum exosomes may also increase the production of inflammatory factors. The study conducted by Dai et al.³⁴² showed that exosomes produced from M2 macrophage exosomes mitigated myocardial I/R damage by suppressing TXNIP and deactivating the

TABLE 7 Therapeutic potential of mesenchymal stem cell-derived extracellular vesicles (MSC-EVs) in autoimmune and rheumatic diseases.

Injury	Animal model	Infusion method	Origin of EVs	Dose of injection	Outcome	References
Inflammatory bowel disease (IBD)	Rat	Intravenous injection	BMSC-EV-derived miR-146a	100 µg	Attenuate experimental colitis by targeting TRAF6 and IRAK1	132
IBD	Mice	Intraperitoneal injection	BMSC-EVs	50 µL	Attenuate dextran sodium sulfate-induced ulcerative colitis by promoting M2 macrophage polarization	272
IBD	Mice	Intraperitoneal injection	AD-MSC-EVs	-	Have immunosuppressive and anti-inflammatory functions	267
IBD	Mice	Intravenous injection	Huc-MSC-Exo-derived miR-378a-5p	1 mg	Attenuate colitis by regulating macrophage pyroptosis via the NLRP3 pathway	296
IBD	Mice	Intravenous injection	AD-MSC-Exos	300 µg	Protect mice from DSS-Induced IBD by promoting intestinal-stem-cell and epithelial regeneration	266
IBD	Mice	Intraperitoneal injection	AD-MSC-Exos	100 µg	Alleviate acute colitis by Treg cell induction and inflammatory cytokine reduction	265
IBD	Mice	Intravenous injection	Huc-MSC-Exo-derived miR-326	1 mg	Inhibit neddylation to relieve IBD	297
IBD	Mice	Intraperitoneal injection	Huc-MSC-Exo-derived TSG-6	200 µg	Protect against IBD through restoring mucosal barrier repair and intestinal immune homeostasis	298
IBD	Mice	Intravenous injection	MSC-Exos	-	Reduce murine colonic inflammation via a macrophage-dependent mechanism	273
IBD	Mice	Intravenous injection	Olfactory ecto-MSC-Exos	60 µg	Ameliorate colitis via modulating Th1/Th17 and Treg cells	274
IBD	Mice	Intravenous injection	Huc-MSC-Exos	400 µg	Alleviate IBD through the modulation of IL-7 expression in macrophages	270
IBD	Mice	-	MSC-EVs	-	Resulted in polarization of intestinal macrophages toward and anti-inflammatory phenotype	271
IBD	Mice	-	Huc-MSC-Exos	200 µg	Possess immunosuppressive effect in vitro and exhibit a therapeutic capability in vivo through suppressing inflammation mechanism	268
IBD	Mice	-	Placental MSC-EVs	200 µg	Alleviate experimental colitis in mice by inhibiting inflammation and oxidative stress	275
IBD	Mice	Intraperitoneal injection	Human dental pulp-MSC-EVs	50 µg	Decrease fibrosis and inflammation by increasing the level of M2 macrophages	276
Type 1 diabetes mellitus (T1DM)	Mice	Intravenous injection	BMSC-Exos	0.5 µg	Improve the cognitive impairments of by repairing damaged neurons and astrocytes	280
T1DM	Mice	Intraperitoneal injection	AD-MSC-Exos	50 µg	Exert ameliorative effects through increasing Tregs, IL-4, IL-10, TGF-β and reduction in IL-17 and IFN-γ	281

TABLE 7 (Continued)

Injury	Animal model	Infusion method	Origin of EVs	Dose of injection	Outcome	References
T1DM	Mice	Intravenous injection	MSC-Exos	30 µg	Attenuate immune responses	282
Osteoarthritis (OA)	Mice	Articular injection	Synovial MSC-Exo-derived miR-155-5p	30 µL	Prevent OA via enhancing proliferation and migration, attenuating apoptosis, and modulating extracellular matrix secretion in chondrocyte	232
OA	Mice	Articular injection	BMSC-Exos	250 ng	Protect cartilage and bone from degradation in osteoarthritis	210
OA	Rat	Articular injection	BMSC-Exos	10 ¹⁰ /mL	Prevent OA by regulating synovial macrophage polarization	209
OA	Mice	Articular injection	iMSC-Exos and SM-MSC-Exos	10 ¹⁰ /mL	iMSC-Exos exerted a stronger stimulatory effect on chondrocyte migration and proliferation than did SM-MSC-Exos	229
OA	Rat	Articular injection	SMSC-Exo-derived miR-140-5p	100 µg	Enhance cartilage tissue regeneration and prevent OA of the knee	230
OA	Mice	Articular injection	Polydactyl BMSC-Exos	-	Alleviate OA by promoting chondrocyte proliferation	236
OA	Mice	Articular injection	LPS-primed SMSC-ECVs	10 ¹¹ /mL	Inhibit ECM degradation and prevent OA	231
OA	Mice	Articular injection	Embryonic-MSC-Exos	10 ⁶ /mL	Alleviate OA through balancing synthesis and degradation of cartilage extracellular matrix	233
OA	Rat	Joint cavity injection	BMSC-Exos	40 µg	Promote cartilage repair and extracellular matrix synthesis, as well as alleviate knee pain	234
OA	Rat	Joint injection	Huc-MSC-Exo-derived miR-26a-5p	250 ng	Alleviate OA via downregulation of PTGS2	235
OA	Mice	-	MSC-Exo-derived KLF3- AS1	-	Elevate chondrocyte proliferation and inhibit apoptosis through miR-206/GIT1 axis	237
OA	Rat	Articular injection	Embryonic stem cell-derived MSC-Exos	100 µg	Reduce pain and repair osteoarthritic TMJs by attenuating inflammation and restoring matrix homeostasis	238
OA	Mice	-	BMSC-Exo-derived miR-92a-3p	500 µg	Enhance chondrogenesis and suppress cartilage degradation via targeting WNT5A	149
OA	Rabbit	Subcutaneous injection	ECM/GelMA/BMSC-Exos	-	Restored chondrocyte mitochondrial dysfunction, enhanced chondrocyte migration, and polarized the synovial macrophage response toward M2	239
OA	Mice	Articular injection	IPFP-MSCs-Exo-derived miR-100-5p	10 ¹⁰ /mL	Protect articular cartilage and ameliorate gait abnormalities via inhibition of mTOR	240
OA	Rat	Articular injection	AD-MSC-Exos	100 µg	Facilitate cartilage injury repair and alleviate OA	299

(Continues)

TABLE 7 (Continued)

Injury	Animal model	Infusion method	Origin of EVs	Dose of injection	Outcome	References
Osteoporosis (OP)	Rat	-	BMSC-Exo-derived miR-150-3p	-	Promote osteoblast proliferation and differentiation in osteoporosis	284
OP	Rat	Intravenous injection	GPNCB-modified BMSC-EVs	100 µg	Attenuate bone loss in an ovariectomized rat model of OP	300
OP	Mice	Intraperitoneal injection	Huc-MSC-Exos	-	Induce osteogenesis and prevent OP	286
OP	Mice	Intravenous injection	BMSC-Exos	0.1 mL	Inhibit RANKL-induced osteoclastogenesis through the NF-κB signaling pathway	289
OP	Rat	Intravenous injection	Epimedium-Preconditioned BMSC-EVs derived miR-27a-5p	100 ng	Stimulate osteogenesis by targeting Atg4B-mediated autophagy	290
OP	Rat	Intravenous injection	BMSC-Exos derived miR-186	10 ¹³ /mL	Promote osteogenesis through hippo signaling pathway in OP	292
OP	Rat	-	BMSC-EVs derived miR-20a	-	Promote the osteointegration of porous titanium alloy by enhancing osteogenesis via targeting BAMBI	293
OP	Rat	-	Huc-MSC-Exos derived miR-1263	-	Prevent apoptosis in disuse osteoporosis	285
Multiple sclerosis (MS)	Mice	Intravenous injection	AD-MSC-EVs	25 µg	Attenuate motor deficits through immunomodulatory actions, diminishing brain atrophy and promoting remyelination	74
MS	-	-	Huc-MSC-Exos	-	Suppress proliferation of activated PBMCs	301
MS	Mice	Intravenous injection	MSC-Exos armed with high-affinity aptamer	200 µg	Produced synergistic immunomodulatory properties and remyelination effect	302
MS	Rat	Intravenous injection	BMSC-EVs	100 and 400 µg	Attenuate inflammation and demyelination of the central nervous system in EAE rats by regulating the polarization of microglia	76
MS	Mice	Intravenous injection	Placental MSC-EVs	10 ⁷ or 10 ¹⁰ /mL	Promote myelin regeneration	80
Amiotrophic lateral sclerosis (ALS)	Mice	Intravenous or intraperitoneal injection	IFN-γ-primed MSC-EVs	-	Affect neuroinflammation possibly through specific immunomodulatory miRNAs acting on microglia	303
Rheumatoid arthritis (RA)	Mice	Intraperitoneal injection	BMSC-Exo-derived miRNA-150-5p	50 µg	Decrease migration and invasion and inhibiting angiogenesis in vitro and alleviating the symptoms of RA by the modulation of MMP14 and VEGF	216
RA	Rat	-	BMSC-Exo-derived miR-192-5p	50 mg	Delays inflammatory response	214
RA	-	-	BMSC-Exo-derived miR-320a	-	Regulate RA-FLS activation by suppressing CXCL9	215

TABLE 7 (Continued)

Injury	Animal model	Infusion method	Origin of EVs	Dose of injection	Outcome	References
RA	Rat	Intravenous injection	BMSC-EV-derived miR-34a	75 µg	Reduce RA inflammation via the cyclin I/ATM/ATR/p53 axis	213
RA	Mice	Intravenous injection	BMSC-EVs	-	Exert an anti-inflammatory role on T and B lymphocytes independently of MSCs priming	208
Sjögren's syndrome (SS)	Mice	-	Labial gland-MSC-Exos	50 µg	Ameliorate SS by modulating the balance of Treg and Th17 cells	304
SS	Mice	Intravenous injection	Olfactory ecto-MSC-Exos	100 µg	Ameliorate SS by modulating the function of myeloid-derived suppressor cells	305
Systemic lupus erythematosus (SLE)	Mice	Intravenous injection	Huc-MSC-Exos	100 µg	Regulate macrophage polarization to attenuate SLE-associated diffuse alveolar hemorrhage	306
SLE	-	-	MSC-Exo-derived tsRNA-21109	-	Alleviate SLE by inhibiting macrophage M1 polarization	307
Systemic Sclerosis (SSc)	Mice	Intravenous injection	IFN-γ-Primed-MSC-EVs	250 ng	Improved lung fibrosis by modulating anti-inflammatory and antifibrotic markers	308
SSc	Mice	Intravenous injection	MSC-EV-derived miR-29a-3p	100 µg	Downregulate the expression of several pro-fibrotic, remodeling, and anti-apoptotic factors as well as methylases	309
SSc	Mice	Subcutaneous injection	MSC-Exo-derived miR-196b-5p	20 µg	Suppress skin fibrosis	310
SSc	Mice	-	BMSC-EVs	15 µg	Alleviate SSc pathogenic changes by regulating the WNT and TGF-β signaling pathways	311
SSc	Mice	-	MSC-Exos	-	Ameliorate dermal fibrosis in bleomycin-induced scleroderma	312
Chronic graft-versus-host disease (cGVHD)	Mice	Intraperitoneal injection	Huc-MSC-EVs	100 µg	Prevent skin fibrosis by suppressing the activation of macrophages and B cells immune response	313

Note: Reprinted with permission.¹²⁷

TABLE 8 Role of MSC-Exos in the treatment of autoimmune diseases (all sources).

Disease types	Exosome source	Reactive molecules	Target cells/tissue	Mechanism of action	Effect	References
Experimental autoimmune encephalomyelitis	BMSCs	-	Microglia	Regulated the polarization of microglia	Attenuated inflammation and demyelination of the CNS	76
	HPDLSCs	-	Spinal cord	Modulated NF- κ B level and inhibited NALP3 inflammasome activation	Reduced pro-inflammatory infiltration	314
	MSCs	HSP70 Galectin 1 PD-L1	Spinal cord T/B lymphocyte	Promoted the differentiation of Tregs Reduced pro-inflammatory cytokines	Reduced demyelination and decreased neuroinflammation	75
Collagen-induced arthritis	BMSCs	-	T/B lymphocyte	Regulated the differentiation of lymphocyte anti-inflammatory subtype and the release of cytokine	Decreased clinical signs of inflammation	208
	BMSCs	miR-192-5p	HFLS-RA	Upregulated RAC2	Delayed the pro-inflammatory response in RA	214
Rheumatoid arthritis	MSCs	miR-150-5p	FLS	Targeted MMP14 and VEGF to decrease migration and invasion in RA FLS	Inhibited synoviocyte hyperplasia and angiogenesis	216
	HMSCs	miR-124a	HFLS-RA	-	Inhibited the proliferation and migration of fibroblast-like synoviocyte cell line	315
Osteoarthritis	HMSCs	lncRNA-KLF3-AS1 miR-206/GIT1	Chondrocytes	Promoted chondrocyte proliferation, migration, and inhibited apoptosis	Facilitated cartilage repair	138,237
Type 1 diabetes	AD-MSCs	-	T/B lymphocyte	Promoted the differentiation of Tregs and the release of cytokine	Improve islet function, and decrease blood glucose	281
	MSCs	Pdx-1	Islet	-	Induced the β cells mass regeneration	316
	ADSCs	miR-126 miR-130a miR-132 miR-let7b miR-let7c	Penis	Promoted angiogenesis Inhibited the cavernous fibrosis	Ameliorated erectile function	317
	MSCs	miR-151-5p	Recipient bone marrow MSCs	Inhibited IL-4R α expression and downregulated mTOR pathway activation	Reduced the damage of recipient bone marrow mesenchymal stem cells to improve osteoporosis	318
Experimental autoimmune uveoretinitis	MSCs	-	Retina	Inhibited the chemo-attractive effects of CCL2 and CCL2 I on inflammatory cells	Reduced the infiltration of T cells subsets and other pro-inflammatory cells	319
Autoimmune hepatitis	BMSCs	miR-223-3p	Macrophages	Regulated the expression of STAT3 gene and inflammatory cytokine	Attenuated pro-inflammatory response	147
	BMSCs	miR-223	Liver	Regulated NLRP3 and caspase-1	Protected liver injury	320

Note: Reprinted with permission.¹²⁷

TABLE 9 Exosomes from different cell sources and cardiovascular signal transduction.

Cells	Inclusions	Factor	Function
Cardiomyocytes	miR-222/143	HIF-1 α , TNF- α	Induce angiogenesis
	miR-217	IL-6, CCL2/6/7	Cardiac hypertrophy, cardiac fibrosis (enhance the proliferation of fibroblasts)
Cardiac fibroblasts	miR-21-3p	Ang II	Induce cardiomyocyte hypertrophy
Cardiac endothelial cells	miR-214	16K PRL	Inhibit the senescence of endothelial cells, induce angiogenesis
	miR-146a	ErbB4, Notch1, Irak1	Reduces the metabolic activity of cardiomyocytes
	miR-143/145	KLF2	Induced atherosclerotic protective phenotype
	miR-92a-3p	THBS1 dependent mechanism	Regulating the angiogenesis
	-	ERK1/2 MAPK signaling pathway	Resistance to simulated I/R injury
	Mst1	-	Inhibiting autophagy, promoting cell apoptosis, and inhibiting glucose metabolism in cardiomyocytes
	VCAM1	-	Regulate the local inflammatory response
Smooth muscle cells	-	coagulation protein prothrombin	Regulate blood coagulation and calcification
Immune cells (dendritic cells)	-	CD4(+) T cells	Promote wound healing after MI
Immune cells (macrophages)	mir-155	-	Inhibited the proliferation of fibroblasts and enhanced inflammation.
Other heart-derived	miR-21-5p	PTEN/Akt pathway	Promotes angiogenesis and cardiomyocyte survival
Circulating exosomes	AT1Rs	Ang II	Improve blood pressure responsiveness
	miR-939-5p	iNOS-NO pathway	Promote angiogenesis
	miR-342-5p	-	Protect the heart from myocardial I/R injury
	myo-miRs	CXCR4	Systemic responses to cardiac injury

Note: Reprinted with permission from Cui et al.³²⁴

Abbreviations: 16K PRL, 16-kDa N-terminal prolactin fragment; Akt, Protein kinase B, also known as PKB or Rac; Ang II, Angiotensin II; AT1Rs, Ang II receptor type 1; CCL2/6/7, CC-motif chemokine ligands 2/6/7; CD4, cluster of differentiation 4; CXCR4, C-X-C motif chemokine receptor type 4; ERK1/2, extracellular signal-regulated kinase1/2; HIF-1 α , hypoxia-inducible factor-1 α ; I/R, ischemia-reperfusion; IL-6, interleukin-6; iNOS, inducible nitric oxide synthase; KLF2, Kruppel-like factor 2; MAPK, mitogen-activated protein kinase; MI, myocardial infarction; NO, nitric oxide; PTEN, gene of phosphate and tension homology deleted on chromosome 10; THBS1, thrombospondin 1; TNF- α , tumor necrosis factor- α ; VCAM1, vascular cell adhesion 1.

TLR4/NF- κ B/NLRP3 inflammasome signaling pathways. Several other studies also found positive outcomes when investigating exosomes on I/R injury.³⁴³⁻³⁴⁵

Zhu et al.³⁴⁶ showed that exosomes produced from ADSCs enhanced the formation of new blood vessels (angiogenesis) in the ischemic hind limbs and heart of mice. This effect was mediated by the miRNA-31/FIH1/HIF-1 α pathway, leading to an improvement in MI damage. Pan et al.³⁴⁷ extracted exosomes from ADSCs that were either unmodified or modified with miR-146a. They then assessed the therapeutic impact of these exosomes in a rat model of acute myocardial infarction (AMI) by using hypoxia-induced H9c2 cells. The study revealed that exosomes containing miR-146a had a more pronounced effect compared with the group treated with unmodified exosomes in suppressing AMI-induced apoptosis, inflammation, and fibrosis.³⁴⁷ The study further discovered that exosomal miR-146a diminishes myocardial damage caused by AMI by suppressing the EGR1-mediated TLR4/NF- κ B signaling pathway.³⁴⁷

Studies in pigs of both acute and chronic MI models demonstrated that the administration of exosomes isolated from

cardiosphere-derived cells improved cardiac healing by enhancing cardiac function and reducing infarct size.³⁴⁸ Moreover, it was recognized that cardiac progenitor cells (CPCs) control heart repair and protection. Significantly, the exosomes obtained from CPCs also enhanced the viability and reproduction of H9C2 cells by promoting the expression of Akt and activating the Akt/mTOR pathway.³⁴⁹ A pediatric investigation has shown that exosomes obtained from newborn CPCs enhanced heart function and repair. In contrast, exosomes produced from CPCs of older children need hypoxic preconditioning in order to manifest cardioprotective advantages. The observed outcomes included enhanced formation of new blood vessels (angiogenesis) and decreased formation of scar tissue (fibrosis), leading to enhanced functioning of the heart after a heart attack (infarction).³⁵⁰

In addition, exosomes produced from CD34⁺ hematopoietic stem cells (HSCs) exhibited a significant expression of pro-angiogenic miRNAs (miRNAs), such as miR-126 and miR-130, which enhance the development of blood vessels in the damaged heart.³⁵¹ A study of miRNA sequences revealed a comparable miRNA profile between




FIGURE 13 miRNAs target myocardium in pathological process after MI. miRNAs are involved in myocardial cell apoptosis, myocardial fibrosis, and myocardial hypertrophy acting on myocardial targets after MI. Upregulation of red miRNAs promoted the development of pathology, while upregulation of green miRNAs inhibited or even reversed the pathological process. Reprinted with permission from Wang and Zheng.³³⁶

MSCs and exosomes formed from MSCs, indicating a mechanistic commonality in the way MSCs and MSC-Exos contribute to cardiac healing. This highlights the enhanced therapeutic significance of exosomes produced from MSCs compared to MSCs themselves in the context of heart repair and has reported as being much safer.³⁵² Additional research has shown that exosomes obtained from MSCs that have been preconditioned under hypoxic conditions exhibit higher levels of several miRNAs associated with better cell survival, angiogenesis, and decreased fibrosis. As a result, these exosomes produced from hypoxic MSCs are more effective in promoting cardiac repair compared to exosomes derived from MSCs under normal oxygen conditions.³⁵³⁻³⁵⁶ Additional examination shown that pre-treating CSCs with exosomes produced from MSCs also leads to enhanced viability and angiogenic capability of CSCs.³⁵⁷

4.1.3 | Hypertension

Hypertension is a prevalent global health issue that significantly contributes to the occurrence and death rates of CVDs. An estimated 1.13 billion individuals worldwide were affected by this silent illness, and the number continues to rise.³⁵⁸⁻³⁶⁰ According to the American Heart Association, about 45% of individuals in the United States have uncontrolled hypertension. Vascular smooth muscle cell (VSMC) migration is crucial for the restructuring of hypertensive blood arteries, whereas adventitia fibroblasts (AFs) have a

significant impact on the maintenance of vascular structures. Tong et al.³⁶¹ cultured primary VSMCs and AFs from the aorta of spontaneously hypertensive rats and Wistar-Kyoto (WKY) rats and found that AF exosomes from hypertensive rats convert Ang-converting enzyme transfer to VSMCs, thereby increasing the level of Ang II and activating the Ang II type 1 receptor (ATR) in VSMCs, thereby promoting VSMC migration.

Due to their unique properties, EVs have attracted attention as a possible prognostic and therapeutic option for the treatment of hypertension. First, EVs serve as prognostic biomarkers and may be collected using minimally invasive (circulating EVs) or noninvasive (urinary EVs) procedures. They also provide information to the clinician on the potential etiology of hypertension since EVs include components from their parent cells. It has been proposed that EVs may serve as surrogate indicators for vascular damage, endothelial dysfunction, and elevated renal sodium transporter/exchanger activity in hypertension.³⁶²⁻³⁶⁵ Additionally, EVs may be used as indicators of enhanced vascular endothelial function.

During a 12-week course of treatment with aliskiren (a direct renin inhibitor), hypertensive patients receiving hemodialysis had increased levels of platelet-derived EVs and flow-mediated dilatation.³⁶⁶ The goal of further research was to examine EVs as potential functional biomolecules as potential biomarkers.³⁶⁷ These studies were employed to track the vascular health of individuals including the expression of circulating EVs and their favorable correlation with the endothelial activation marker E-selectin.³⁶⁸

Equally relevant, EVs have significant benefits as therapeutic agents. Due to their minimal immunogenicity, EV-based therapeutics have the potential to overcome many of the challenges associated with cell-based therapies.^{369–371} EVs are currently in the early stages of development as a therapeutic option for hypertension. Research conducted *in vivo* has revealed that circulating EVs from WKY and SHR with low plasma levels may modify the vasoreactivity of isolated mesenteric arteries in a distinct way.³⁷² In a preclinical model, plasma EVs had the ability to control systemic blood pressure and have a positive impact on end-organ damage caused by hypertension. While still early in the process, EV-mediated therapies for blood pressure management should be available in the near future offering much hope for the management of hypertension in a more natural way. Additionally, disease progress and resolution can be monitored by using exosomes as biomarkers throughout the disease process.

4.1.4 | Atrial fibrillation

Atrial fibrillation (AF) is the most prevalent type of cardiac arrhythmia worldwide, has a heavy socioeconomic burden, high morbidity, and death rate, as well as poses major consequences including heart failure and stroke.^{373,374} It is predicted that the incidence of AF will more than double over the next 40 years, impacting 1%–1.5% of the global population. The condition's frequency is directly correlated with growing age.^{375,376} An established therapy for AF, particularly for paroxysmal AF (PAF), is catheter ablation. However, since the operation is often accompanied by hazards and other clinical problems, the success rate for persistent AF (PsAF) is not optimal. Additionally, the AF lacks efficient upstream management.^{377–379}

At present, there are relatively few studies on exosomes and AF, and the studies that have been conducted have mainly focused on the diagnosis of AF. Fibroblast activation of myofibroblasts is a crucial step in the pathophysiology of fibrosis. Wei et al.³⁸⁰ demonstrated that the expression of miRNA extracted from the plasma of individuals with AF and normal sinus rhythm differs. Research by Mun et al.³⁸¹ established a connection between atrial shape and function, oxidative stress, and fibrosis pathways through serum exosomal miRNA and the genes it targets. Based on these findings, exosomal miRNAs show some promise as potential biomarkers for tracking AF development.

4.1.5 | Heart failure

Heart failure (HF) is a clinical state that may be caused by several disorders affecting the myocardium, cardiac valves, pericardium, or vasculature. Idiopathic dilated cardiomyopathy is one such ailment. Ischemic heart disease is another. The main symptoms of HF are usually exhaustion, edema, and dyspnea. HF is mostly caused by coronary artery disease, valve disease, hypertension, and dilated

cardiomyopathy in the Western world.³⁸² HF is an expensive and possibly fatal illness.³⁸³ While the lifetime risk of heart failure is already quite high (20%–45%),³⁸⁴ projections indicate that the prevalence of the condition is expected to increase even further, and the spending on treatment/therapy is expected to increase by 127% by the year 2030.³⁸⁵

Chen et al.³⁸⁶ discovered that exosomes obtained from BMSCs had a substantial protective effect on the myocardium against cardiac hypertrophy, reducing myocardial cell death and fibrosis, and preserving cardiac function under pressure overload conditions. Additionally, studies have demonstrated that exosomes generated from MSCs also inhibit cell hypertrophy induced by Ang II. Exosomes also stimulate the early recovery of aging of myofibroblasts in a laboratory setting, suggesting that they have beneficial effects in reducing fibrosis during cardiac remodeling. In summary, exosomes offer cardiomyocytes protection from pathological hypertrophy and hold great potential as a therapeutic option for HF.

4.1.6 | Atherosclerosis

Atherosclerosis (AS) is an inflammatory condition affecting blood vessels characterized by the buildup of lipids in the vessel walls. Accumulation results in the deposition of plaques and the narrowing of the channel's inner opening.^{387,388} Smoking, obesity, diabetes, and vascular damage are significant contributors to the accumulation of fatty substances and cholesterol in the bloodstream, which is a key element in the development of the disease.^{389–392} Atherosclerosis involves alterations in the characteristics of vascular systems, such as impaired function of endothelial cells (EC), increased proliferation and movement of vascular smooth muscle cells (VSMC), calcification of blood vessels, inflammation, infiltration of macrophages into the plaque, and polarization of macrophages.^{393–395}

Multiple studies have shown that exosomes have a significant impact on different phases of the onset and progression of AS. They have a crucial function during inflammation, oxidative stress, and apoptosis.^{396–403} Research has shown that exosomes derived from many origins may contribute to the onset and progression of AS by engaging in the NF- κ B signaling pathway.^{396–398} Yao et al.³⁹⁶ found that the overexpression of peripheral platelet exosomal (PLT-exosomes) miR-25-3p inhibited coronary vascular endothelial cell inflammation. Further studies found that the inhibitory effect of PLT-exosomes carrying miR-25-3p is related to the NF- κ B signaling pathway.³⁹⁶ Zhong et al.³⁹⁷ verified that exosomes obtained from fully developed dendritic cells (mDC-exosomes) have a role in endothelial inflammation and atherosclerosis (AS). The process involves the transfer of miR-146 from mDC-exosomes to HUVECs. The activation of the NF- κ B signaling pathway has been shown to enhance the production of adhesion molecules in endothelial cells and shields HUVECs from further stimulation by suppressing IL-1 receptor-associated kinases. This suggests that mDC-exosomes play a key role in the feedback mechanism that regulates inflammation in a negative manner.

Oxidized low-density lipoprotein (ox-LDL) also has a significant impact on the initiation and progression of AS by activating macrophages and endothelial cells. After discovering that exosomal miR-146a, which are derived from ox-LDL-treated macrophages, Zhang et al.⁴⁰⁰ discovered that AS in mice was greatly worsened after intravenous injection of exosomes originating from ox-LDL-treated THP-1 cells. Chen et al.⁴⁰¹ found that exosomal miR-505 from ox-LDL-treated vascular endothelial cells intensified AS by triggering the production of NETs.

Another study by Li et al.³⁹⁹ showed that the miR-let7/HMGA2/NF- κ B pathway was responsible for the improvement of AS in ApoE mice and the promotion of M2 macrophage polarization in plaques. Furthermore, exosomes produced from MSCs prevented macrophage recruitment in plaques via the miR-let7/IGF2BP1/PTEN pathway. This finding offers novel methods to reduce atherosclerotic plaque inflammation and enhances our knowledge of how MSC-Exos impact this condition. According to Lin et al.,⁴⁰² miR-203-3p was discovered to decrease cathepsin S production and AS-related characteristics in BM-derived macrophages when transferred from dendritic cell-derived exosomes (DEXs). A prospective therapeutic target for AS, the p38/MAPK signaling pathway, has emerged as a result of studies aiming to slow the progression of the disease in

mice. The combined results demonstrated that exosomes mitigate the development of macrophage foam cells by modulating the expression of cholesterol transporters.⁴⁰³ This discovery unveils a novel method by which PVAT safeguards the vasculature against atherosclerosis.

Figure 14 summarizes the number of findings derived from extensive studies on exosomes as therapeutic agents for the treatment of various heart conditions. These studies have shown that exosomes promote angiogenesis, encourage the polarization of M2 macrophages, reduce the presence of pro-inflammatory immune cells, and mitigate the antifibrotic properties of the injury.⁴⁰⁴

4.2 | Stroke

Ischemic strokes are a major contributor to chronic disability worldwide, and there are few therapies that are successful. Mounting data indicate that exosomes have a role in ischemic disease and have beneficial therapeutic benefits by facilitating cell-to-cell contact in individuals with significant impairment and substantial economic constraints.^{405,406}




FIGURE 14 Mesenchymal stromal cell (MSC)-derived exosomes for cardiac repair and function. Exosomes isolated from different sources of MSCs carry and deliver proteins, nucleic acids (DNA, miRNAs, mRNAs, and other RNAs), and metabolites to the damage heart tissue, consequently promoting cardioprotective effects. Reprinted with permission from Joladarashi and Kishore.⁴⁰⁴

The inflammatory response may be seen as both beneficial and detrimental in the progression of cerebral ischemia.^{407–409} The narrow treatment window (<4.5h) and methodological limitations of ischemic strokes have led to a focus on innovative techniques in clinical research. Inflammation may cause the immune system to become overactive in the early stages of a stroke as inflammatory factors or biological components produced by dying cells stimulates resident microglia and invades peripheral immune cells. Following this, these cells release a range of inflammatory mediators, chemokines, adhesion molecules, tissue-degrading enzymes, and complement system activators.^{410–412} Together, they worsen brain injury and accelerate BBB degradation in a vicious cycle.⁴¹¹

4.2.1 | MSC exosomes for stroke therapy

Clinical investigations and animal models have shown that MSC transplantation may facilitate recovery after a stroke (Table 10).⁴¹³ Increasing data supports the concept that that stem cells primarily exercise their therapeutic benefits via paracrine processes, with a particular focus on the release of exosomes.^{414,415} MSC-derived exosomes have been shown to effectively enhance stroke recovery in cases of ischemia.

Findings have most commonly utilized intravenous delivery of exosomes. It was demonstrated in a rat model that intravenous delivery of MSC exosomes improved function and recovery in the transient middle cerebral artery occlusion (tMCAO) paradigm by stimulating neurogenesis, neurite remodeling, and angiogenesis.⁴¹⁷ The results demonstrated that MSC exosomes and MSCs themselves had almost identical effects on neuroprotection, angiogenesis, and immunomodulation.^{418,419} Exosomes produced from brain extract-treated or oxygen–glucose deprivation (OGD)-treated MSCs also demonstrated superior therapeutic benefits. This may be based on the fact that their exosomes were enriched with certain functional proteins.^{420,426}

Interestingly, MSC-Exos were also shown to be beneficial in a stroke model in primates. The recovery of fine hand motor function was aided by intravenous injection of MSC-Exos 24h and 14 days after the damage in the macaca mulatta cortical hand motor brain injury model.⁴³² Additional mechanistic research revealed that in old macaca mulatta, MSC-Exos not only decreased neuroinflammation but also changed the roles of microglia into restorative ones.⁴³⁰ Furthermore, it impeded hyperexcitability caused by damage and reinstated the equilibrium between excitatory and inhibitory processes.⁴³¹ These effects may be explained by the transfer of functional miRNAs or proteins and the subsequent activation of downstream signaling pathways. Prior to ischemia, intranasal administration of MSC-Exos decreased neuronal mortality, encouraged the formation of oligodendroglia, and suppressed microglia-mediated neuroinflammation, perhaps via the Toll-like receptor 4/CD14/NF- κ B signaling pathway.^{427,433}

MSC-Exos further improved the hippocampus spatial learning and memory impairments in global ischemia, perhaps via

cyclo-oxygenase-2 expression regulation.⁴²⁵ MSC-Exos facilitated axonal sprouting and encouraged the restoration of white matter in cases of subcortical ischemia.⁴²⁴ In vitro, the enhancement of axonal development was facilitated by the transfer of miR-17-92 cluster via exosomes, which then activated the PTEN/mTOR signal pathway.⁴³⁴ Exosomal miR-134 targeted caspase-8 to prevent oligodendrocytes from undergoing apoptosis.⁴³⁵ Another research group that utilized lentiviruses in both knockdowns and knock in miR-133b in MSCs revealed that exosomal transfer of miR-133b to astrocytes and neurons was a partial mechanism by which MSCs mediated their effects.⁴³⁶ MSC-Exos prevented neurons in OGD from dying by transferring let-7-5 p and then suppressing the production of caspase-3.⁴³⁷

Lately, there has been a growing interest in developing techniques to optimize the therapeutic potential of exosomes using engineering methodologies (Figure 15).^{438,439} For example, MSC-Exos that were enhanced with the miR-17-92 cluster showed more significant enhancements in oligodendrogenesis, neurogenesis, and neurite remodeling when compared to the control group.¹⁴⁴ Researchers in a mouse photothrombosis model enhanced the properties of MSC-Exos by including rabies virus glycoprotein (RVG), a peptide that targets neurons. This was achieved by fusing RVG with exosomal protein lysosome-associated membrane glycoprotein 2b (Lamp2b). Subsequently, the exosomes were loaded with miR-124-mimics using electroporation. The engineered exosomes demonstrated effective transportation of miRNA-124 to the area affected by ischemia, resulting in the improvement of brain injury through facilitation of neural progenitor differentiation.⁴²³

4.2.2 | Neural stem cell-derived exosomes for stroke therapy

Since 2018, there has been major interest in studies on exosomes produced from neural stem cells (NSCs) in stroke patients (Table 11). In a thromboembolic stroke model, the effects of exosomes from NSCs and MSCs—both of which were derived from the same pluripotent stem cell line—were compared. It was found that NSC exosomes improved the function and reduced infarct volume more than MSC-Exos, which were linked to a more potent effect in polarizing macrophages toward an M2 phenotype and reducing inflammation.⁴⁴¹ Furthermore, the maintenance of astrocyte activity may be a contributing factor to the efficacy of NSC exosomes in lowering infarct volume.⁴⁴² In a senior stroke rat model, NSC exosomes showed encouraging therapeutic benefits.⁴⁴¹ Significantly, NSC exosome therapy improved white matter integrity and function recovery in a porcine stroke model while also reducing infarct volume and brain edema.⁴⁴³

4.2.3 | Adipose-derived stem cells derived exosomes for stroke therapy

Adipose-derived stem cells (ADSCs) have also shown promise as an ischemic therapy option and are readily acquired from surgically

TABLE 10 Published studies of mesenchymal stem cell-derived exosomes in ischemic stroke.

Animals	Stroke model	Time of treatment	Routes of exosome delivery	Exosome modifications	Proposed mechanisms	References
Rat	2 h tMCAO	24 h after ischemia	Tail vein	No	Promote neurite remodeling, neurogenesis, and angiogenesis	417
Mouse	30 min tMCAO	1, 3 and 5 days after ischemia	Femoral vein	No	Immunomodulation	418
Rat	30 min tMCAO	1, 3 and 5 days after ischemia	Femoral vein	No	Immunomodulation	419
Rat	pMCAO	48 h after ischemia	Common carotid artery	No	Promote angiogenesis, neurogenesis, anti-inflammation	420
Ovine fetuses	25 min global hypoxia-ischemia	1 h and 4 days after ischemia	Intravenously	No	Preserve baroreceptor reflex sensitivity and reduce white matter injury	421
Rat	2 h tMCAO	24 h after ischemia	Intravenously	Enriched in miR-17-92 cluster	Enhance oligodendrogenesis, neurogenesis, and neurite remodeling	144
Mouse	30 min tMCAO	Once every other day for 14 days after ischemia	Tail vein	RGD-exosomes loaded with miR-210	Promote VEGF expression and angiogenesis	422
Mous	Photothrombosis	24 h after ischemia	Tail vein	RVG-exosomes loaded with miR-124	Promote neurogenesis	423
Rat	Endothelin-1 subcortical stroke	24 h after ischemia	Tail vein	No	Promote white matter repair	424
Mouse	5 min bilateral common carotid arteries occlusion	Right before ischemia	Intracerebroventricular injection	No	Inhibition of Cox-2 expression	425
Rat	2 h tMCAO	Immediately after ischemia	Tail vein	No	Reduce oedema	426
Rat	Perinatal brain injury	Before ischemia	Intranasally	No	Neuroprotection, anti-inflammation	427
Rat	2 h tMCAO	24 h after ischemia	Intra-arterially	Enriched with miR-134b	Promote neurite remodeling	428
Macaca mulatta	Cortical hand cerebral injury	24 h and 14 days post-injury	Intravenously	No	Not studied	429
Aged macaca mulatta	Cortical hand cerebral injury	24 h and 14 days post-injury	Intravenously	No	Reduce inflammation, shift microglia toward restorative functions	430
Macaca mulatta	Cortical hand cerebral injury	24 h and 14 days post-injury	Intravenously	No	Dampen injury-related hyperexcitability restores excitatory-inhibitory balance	431

Note: Modified with permission Li et al.⁴¹⁶

Abbreviations: RVG, rabies virus glycoprotein; tMCAO, transient middle cerebral artery occlusion; VEGF, vascular endothelial growth factor.




FIGURE 15 MiRNAs in exosomes secreted by mesenchymal stem cells stimulate cell and structure growth. miRNA-133b can downregulate the expression of CTGF in astrocytes, reduce the formation of glial scar, and promote the remodeling of myelin sheath. miRNA-17-92 can downregulate the expression of PTEN, thereby activating the PIK3-Akt pathway and inactivating GSK-3b, promoting the growth of neuronal axons. miRNA-124 promotes neurogenesis in SVZ and striatum regions. miRNA-22-3p can inhibit KDM6B-mediated KBMP2/BMP pathway and play a neurotrophic role. Akt, protein kinase B; BMP, bone morphogenetic protein; CTGF, connective tissue growth factor; GSK-3b, glycogen synthase kinase-3b; KBMP2, neurotrophic factor; KDM6B, lysine(K)-specific demethylase 6B; PIK3, phosphatidylinositol 3-kinase; PTEN, phosphatase and tensin homolog; SVZ, subventricular zone. Reprinted with permission from Xiong et al.⁴⁴⁰

removed adipose waste tissue. Exosomes from ADSCs are becoming increasingly popular as a replacement treatment (Table 12).⁴⁴⁴ While miR-181b-5p was customized, ADSC exosomes enhanced brain micro-vessel endothelial cell migration and tube formation in vitro.⁴⁴⁵ In rats with ischemic conditions, systemic injection of ADSC exosomes overexpressing miRNA-126 has the potential to efficiently suppress neuroinflammation, decrease neuronal death, stimulate neurogenesis, and aid in functional recovery when compared to normal and miRNA-126 knockdown ADSC exosomes.⁴⁴⁶ ADSC-Exos enriched in miR-30d-5p protected brain damage like the normal and knockdown groups via enhancing M2 microglia polarization. A mechanistic investigation on beclin-1 and autophagy-related genes found that MiR-30d-5p inhibited M1 microglia polarization.⁴⁴⁷ Intraventricular injection of ADSC exosomes modified with a multi-functional protein pigment epithelium-derived factor inhibited neuronal death by upregulating the expression of autophagy-associated proteins in a rat ischemia model.⁴⁴⁸

4.2.4 | Other cell-derived exosomes for stroke therapy

Other cell-derived exosomes, including microglia, endothelial cells, and astrocytes, have shown therapeutic benefits in stroke patients

(Table 13). Exosomes generated from astrocytes have been shown to control autophagy, decrease infarct volume, and prevent neuron death.⁴⁵⁰ By releasing miR-92b-3p, exosomes from OGD-preconditioned astrocytes reduced OGD-induced neuronal death.⁴⁵¹ Sema-3A inhibitors may decrease astrocyte activation and increase axonal elongation in rats with persistent ischemia via activating the GTPase-1/R-Ras/Akt/GSK-3β signaling pathway. Additionally, the authors confirmed that exosomes from astrocytes treated with OGD Sema-3A inhibitors enhanced neuron axonal outgrowth more than exosomes from both normal and OGD astrocytes.⁴⁵²

Exosomes derived from normal microvascular endothelial cells were shown to prevent astrocyte death in vitro, lessened BBB disruption, decreased infarct volume, and aided in the restoration of neurological function in vivo.⁴⁵³ In contrast, exosomes from OGD endothelial cells had the opposite effects.⁴⁵³ These inconsistencies may be explained by variations in the exosomal protein and miRNA contents that were separated from the same cells under differing cell culture conditions.⁴⁵⁷

To improve cell migration and invasion during OGD, SH-SY5Y cells were exposed to exosomes produced by OGD human umbilical vein endothelial cells (HUVEC) resulting in cell death reduction.⁴⁵⁸ Furthermore, exosomes from endothelial cells inhibited the activation of macrophages and inflammation by transferring miR-10a and blocking the NF-κB signaling pathway.⁴⁵⁹

TABLE 11 Published studies of neural stem cell-derived exosomes in ischemic stroke.

Animals	Stroke model	Time of treatment	Routes of exosome delivery	Exosome modifications	Proposed mechanisms	References
Mouse	TE-MCAO	2, 14, 38/6, 24, 48 h (in aged mice) after ischemia	Tail vein	No	Immunomodulation, inhibit inflammation	441
Mouse	1-h tMCAO	2 h after ischemia	Internal jugular vein	No	Preserve astrocyte function	442
Pig	pMCAO	2, 14, 24 h after ischemia	Intravenously	No	Protect the integrity of BBB and WM	443

Note: Modified with permission Li et al.⁴¹⁶

Abbreviations: BBB, blood–brain barrier; pMCAO, permanent middle cerebral artery occlusion; TE-MCAO, thromboembolic middle cerebral artery occlusion; WM, white matter.

TABLE 12 Published studies of adipose-derived stem cell-derived exosome in ischemic stroke.

Animals	Stroke model	Time of treatment	Routes of exosome delivery	Exosome modifications	Proposed mechanisms	References
Rat	50-min tMCAO	3 h after ischemia	Intravenously	No	Anti-inflammation, anti-apoptosis	449
Rat	tMCAO	Immediately after ischemia	Tail vein	Enriched with miR-30d-5p	Reduce autophagy and inflammation, and promote microglia M2 polarization	447
Rat	tMCAO	Not mentioned	Intravenously	Enriched with miR-126	Promote neurogenesis, angiogenesis, anti-inflammation	446
Rat	1-h tMCAO	3 days before ischemia	Lateral cerebral ventricle injection	Loaded with pigment epithelium-derived factor	Promote autophagy	448

Note: Modified with permission Li et al.⁴¹⁶

Abbreviation: tMCAO, transient middle cerebral artery occlusion.

Interestingly, a research model using diabetic mice were given brain endothelial cell exosomes intravenously 3 days after ischemia was induced. Their neurological and cognitive results improved, and the findings demonstrated an increased myelin density and axon outgrowth, stimulated angiogenesis, and more M2 macrophage polarization.⁴⁵⁴ These positive benefits were significantly diminished when endothelial cell exosomes with reduced levels of miR-126 were utilized.⁴⁵⁴

Lastly, exosomes derived from BV2 microglia cells that were activated with IL-4 enhanced the process of HUVEC tube formation.⁴⁶⁰ Systemic delivery of M2 microglial exosomes decreased the size of the infarct, encouraged the restoration of neurological function, and prevented the death of neurons, potentially via transferring miR-124 and modifying ubiquitin-specific protease 14 in neurons.⁴⁵⁶ The culmination of these findings suggest the importance of exosomes in intercellular communication and their potential therapeutic use for stroke management (Table 14).

4.2.5 | Routes of exosome administration for stroke therapy

Exosome administration pathways in stroke may be categorized into two main groups: systemic and local administration. Subcutaneous,

intraperitoneal, intranasal, and intravenous injections into the tail vein, femoral vein, or internal jugular vein are all examples of systemic administration methods. The majority of exosomes administered to rats with ischemic stroke models enter the bloodstream via the tail vein.^{417,423,424,441,447,453,456} When indium-111-labeled exosomes were administered 1 h after ischemia, single photon emission CT (SPECT) imaging revealed that the exosomes first arrived at the infarcted region and were mostly removed from the brain by the next day.⁴⁴¹ A further investigation showed that the process of exosomes being eliminated from the bloodstream started promptly after 1 h and continued progressively for a duration ranging from 1.5 to 6 h after injection.⁴⁷⁶

Using a rat model of prenatal brain damage, a thorough investigation of intranasally administered exosomes before the occurrence of ischemia showed that exosomes were seen in the frontal cortex as soon as 30 min after administration and were equally distributed throughout the whole brain 3 h after being administered.⁴²⁷ The analysis of biodistribution after cerebral ischemia demonstrate that the quantity of exosomes supplied intranasally and labeled with gold nanoparticles was more than twice as high in the brain compared to those administered intravenously. Following intranasal distribution, a significant quantity of exosomes remained in the brain, while there was an almost minimal amount following intravenous delivery 24 h

TABLE 13 Published studies of other cell-derived exosomes in ischemic stroke.

Animals	Stroke model	Time of treatment	Source of exosomes	Routes of exosome delivery	Exosome modifications	Proposed mechanisms	References
Mouse	pMCAO	1 h after ischemia	Astrocyte	Tail vein	No	Anti-apoptosis	450
Mouse	90 min tMCAO	30 min after ischemia	Brain microvascular ECs	Tail vein	No	Protect BBB integrity, inhibit astrocyte activation	453
Mouse	Photothrombosis	3 days after ischemia	Brain endothelial cells	Intravenously	Enriched with miR-126	Promote angiogenesis, neurogenesis and M2 macrophage polarization	454
Rat	2 h tMCAO	Immediately after ischemia	Macrophage cell line	Tail vein	No	Promote M2 microglial polarization	455
Mouse	90 min tMCAO	1, 2, 3 days after ischemia	Microglia cell line	Tail vein	No	Promote neuronal survival	456

Note: Modified with permission Li et al.⁴¹⁶

Abbreviations: pMCAO, permanent middle cerebral artery occlusion; tMCAO, transient middle cerebral artery occlusion.

after injection. Significantly, a considerable proportion of exosomes supplied by the nasal route also gathered in the lungs, spleen, and kidney.⁴⁷⁷ The findings indicated that delivering exosomes via the nose was a more effective and promising noninvasive treatment approach for ischemic stroke.

The typical method of administering local anesthesia involves injecting it into the brain ventricle.^{425,448} Intracerebroventricular injection of MSC-Exos prevented cyclo-oxygenase-2 expression, enhanced spatial learning and memory, and restored hippocampal synaptic transmission impairments in global ischemia.⁴²⁵ Furthermore, infarct volume and cell apoptosis were improved 3 days after tMCAO by ADSC exosomes administered via the lateral cerebral ventricle for approximately 3 days before ischaemia.⁴⁴⁸

4.2.6 | Exosome clinical trials in stroke

Exosomes were shown in clinical pilot studies to be adequate in promoting neurovascular remodeling and functional recovery after ischemic stroke. To encourage the translation of exosomes into the clinic, well-designed clinical trials are necessary since the clinical studies of exosome-based treatment for stroke have just recently started. To date, documented clinical studies (NCT03384433) are presently being investigated in patients on the use of exosomes to treat stroke with IRB approval. Clinical investigators are still debating the best route of MSC-Exo administration. Patients in this clinical study received 200mg of MSC exosomes 1 month following the commencement of ischemia. 12 months after the delivery of exosomes, the modified ranking scale and adverse treatment-related events were measured.

Remarkably, the majority of clinical studies concentrate on the functions that miRNAs play in stroke in terms of prognosis, diagnosis, and prediction. For instance, the analysis of blood miRNAs from 1523 controls aged 40–69 years and 173 cases of cerebrovascular illness aged above 69 years is being investigated to determine serum miRNAs for the prediction of stroke risk. They discovered 10 miRNAs that connected with a projected risk of stroke (miR-1268b, miR-4433b-3p, and miR-6803-5p).⁴⁷⁸ The purpose of clinical trials NCT04175691 and NCT04230785 is to use next-generation sequencing to examine patients' expression patterns of circular RNA, microRNAs, and long non-coding RNA. The trials will then investigate and differentiate related biomarkers that can be utilized to detect and predict the outcome of acute ischemic stroke, as well as track the progression and prognosis of acute ischemic stroke patients undergoing endovascular treatment. A final clinical study (NCT03577093) aims to explore the molecular pathways via which microRNA-494 regulates the cell cycle after cerebral ischemia. These investigators intend to collect peripheral blood DNA samples from individuals aged 18–80 who had a stroke within 6 h. Over the coming years, additional exosomal signals and pathways will be discovered including their ability to aid in neuroprotection (Figure 16).

4.3 | Liver

A number of articles have now been published on the use of exosomes for the management or treatment of various illnesses/diseases involving the liver. These include acute liver failure, drug-induced liver injury, hepatocellular carcinoma, and viral and alcohol-induced hepatitis and cirrhosis.

TABLE 14 Therapeutic effects of exosome on stroke by targeting microglia polarization.

Exosome	Objectives	Contents	Significance	Mechanism	References
Exosomes from serum of young rats	Aged ischemic rats	More CD46, less C1q, C3a, C3b	Improved short and long-term functional outcomes after ischemic stroke and reduced synaptic loss	Reducing Iba1 ⁺ CD86 ⁺ microglia but increasing Iba1 ⁺ CD206 ⁺ microglia	462
EVs from serum	AIS patients	hsa-miR-124-3p	Reduced serum pro-inflammatory cytokines and the NIHSS score	Reversing the LPS-induced inflammatory effect in BV2 microglia by inhibiting the expression of GRB2 and AKT3 gene involved in pro-inflammatory signaling pathways	463
Exosome from ADSCs	AIS rats	miR-30d-5p	Reduced cerebral injury	Suppressing autophagy and promoting M2 microglia polarization	460
Exosomes from serum	Endotoxemia mice	miR-15a, miR-15b, miR-21, miR-27b, miR-125a, miR-146a, and miR-155	Increased systemic pro-inflammatory cytokine production, and elevated CNS expression of pro-inflammatory cytokine mRNA and the inflammation-associated miR-155	Inducing the expression of Iba-1 and microglial uptake of exosomes derived from serum-containing inflammation-related miRNAs	464
Exosomes from BMSCs	ICH rats	miR-146a-5p	Improved neurological function and reduced neuronal apoptosis	Inhibiting microglial M1 polarization by downregulating the expression of IRAK1 and NFAT5	465
Exosomes from hUMSCs	Ischemic mice	miR-146a-5p	Improved recovery of function, attenuated microglia-mediated inflammation	Decreasing IBA-1 ⁺ CD16 ⁺ cells and increasing IBA-1 ⁺ CD206 ⁺ cells by suppressing IRAK1/TRAF6 signaling pathway	466
EVs from neural progenitor cell	MCAO mice	let-7g-5p, miR-99a-5p, let-7i-5p, miR-139-5p, miR-98-5p, miR-21-5p, and let-7b-5p	Suppressed inflammation response	Inhibiting the expression of Iba-1 and MAPK of an inflammation-related pathway	467
Exosome from BMSCs	MCAO rats	NR	Attenuated cerebral ischemia-reperfusion injury-induced neuroinflammation and pyroptosis	Shifting M1-polarized microglia shifted toward M2-polarized microglia	468
Exosomes from macrophage	pMCAO rats	Edaravone	Enhanced neuroprotection	Promoting the polarization of microglia from M1 to M2	469
Exosomes from plasma	pMCAO rats	Melatonin	Decreased infarct volume, improved recovery of function, and reduced microglia pyroptosis	Inhibiting TLR4/NF- κ B pathway-mediated microglial inflammation and NLRP3-mediated microglia pyroptosis	470
Exosomes from hUMSCs	tMCAO rats	CCR2	Enhanced oligodendrogenesis and remyelination	Decreasing CD16 and IL-1 β mRNA expression and increasing CD206 and Arg-1 mRNA expression	471
Exosomes from LPS stimulated macrophage	MCAO/R rats	NR	Increased neuroprotection and functional improvement	Enhancing the microglial polarization from M1 phenotype to M2 phenotype	472
Exosomes from ADSCs	MCAO/R rats	miR-126	Improved neurogenesis and functional recovery	Inhibited microglial activation and the expression of inflammatory factors	473

TABLE 14 (Continued)

Exosome	Objectives	Contents	Significance	Mechanism	References
Exosomes from MSCs	MCAO/R rats	miR-223-3p	Decreased cerebral infarct volume, improved neurological deficits, learning and memorizing abilities	Inhibiting microglial M1 polarization mediated pro-inflammatory response	474
Exosomes from human embryonic kidney cells	Photothrombotic ischemic mice	NGF and NGF mRNA	Reduced ischemic injury via reducing inflammation and cell death	Reducing CD16 ⁺ M1 microglia but increasing CD206 ⁺ M2 microglia	475

Note: Reprinted with permission from Wan et al.⁴⁶¹
Abbreviations: ADSCs, adipose-derived stem cells; AIS, acute ischemic stroke; BMSCs, bone marrow mesenchymal stem cells; C1q, complement component 1q; CCR2, C-C chemokine receptor type 2; CNS, central nervous system; CysLT2R, cysteinyl leukotriene receptor 2; EVs, extracellular vesicles; HUC-MSCs, human umbilical cord mesenchymal stem cells; ICH, intracerebral hemorrhage; IRAK1, interleukin 1 receptor-associated kinase 1; LPS, lipopolysaccharide; MCAO/R, middle cerebral artery occlusion and reperfusion; MSCs, mesenchymal stem cells; NFAT5, nuclear factor of activated T cells 5; NF-κB, nuclear factor-κB; NGF, nerve growth factor; NIHSS, National Institutes of Health Stroke Scale; NLRP3, NLR family pyrin domain-containing 3; NR, not reported; OGD, oxygen–glucose deprivation; pMCAO, permanent middle cerebral artery occlusion; STAT3, signal transducer and activator of Transcription 3; TLR, toll-like receptor; tMCAO, transient middle cerebral artery occlusion; TRAF6, TNF receptor-associated factor 6; USP14, ubiquitin-specific protease 14.

4.3.1 | Acute liver failure

Acute liver failure (ALF) manifests 24 weeks after the beginning of the first symptoms⁴⁸⁰ and are categorized depending on the time and intensity of their clinical presentation: hyperacute, acute, and subacute.⁴⁸¹ Fulminant hepatic failure is present in both the hyperacute and acute forms, although subacute variants are also known as subfulminant.⁴⁸² Remarkably, the fatality rate is lower among individuals who develop hepatic encephalopathy 8 weeks after the first symptoms (fulminant hepatic failure) when compared to those with a more progressive progression.⁴⁸³ Multiorgan failure (MOF) is the primary reason (>50%) for death in cases of ALF, with intracranial hypertension (ICH) and infection being the other significant causes of mortality in this group of patients.⁴⁸⁴

Shokravi and colleagues conducted a thorough review paper examining the use of stem cells and their exosomes in the context of acute liver failure (ALF).⁴⁸⁵ It has been shown that the usage of MSCs affects how cells differentiate into hepatocytes, reduce inflammation, possess antifibrotic effects, and help release and absorb antioxidants.⁴⁸⁵ Exosomes have been shown to be much safer with their effects examined in Table 15. In summary, exosomes have been shown to reduce oxidative stress, enhance angiogenesis, enhance liver survival and function, and lessen inflammation in the liver overall.

4.3.2 | Drug-induced liver injury

Drug-induced liver injury (DILI) is a prominent factor in cases of acute liver injury (ALI), leading to black box warnings and the removal of medications from the market. This condition may be attributed to over 1000 pharmaceuticals that are presently accessible.^{499–501} Nonsteroidal anti-inflammatory, anti-infectious, psychotropic, and hypolipidemic medications were the most frequently implicated pharmaceuticals.^{502,503}

Recent findings indicate that exosomes in circulation and their contents may serve as noninvasive sources of prospective molecular biomarkers for their early identification, tracking, and assessment of DILI. Additionally, it was shown that exosomes present in urine contained RNAs or proteins suggestive of DILI. Exosomes derived from hepatocytes or mesenchymal stem cells are also thought to be promising therapeutic agents since they may reduce hepatocyte apoptosis, regulate the inflammatory response, and encourage liver regeneration. In a mouse model of chronic liver damage caused by CCl4, exosomes were suggested to mitigate hepatic inflammation and collagen deposition.⁵⁰⁴ Additionally, in mice models of CCl4-induced acute liver damage, exosomes generated hepatoprotective effects by increasing hepatocyte proliferation through enhanced proliferating cell nuclear antigen and high cell survivability. EVs also prevented hepatocyte apoptosis caused by APAP and H₂O₂ by up-regulating the expression of the Bcl-xL protein.⁵⁰⁵

Human umbilical cord-derived MSCs (huc-MSCs) are a valuable asset in the field of regenerative medicine for treating liver




FIGURE 16 Neuroprotective mechanism of exosomal miRNA. Reprinted with permission from Xu et al.⁴⁷⁹

damage. These cells have been extensively studied in phase I and II clinical trials with the aim of enhancing liver function. Therefore, huc-MSC exosomes could be the best option for DILI. Apoptosis and reactive oxygen species were considerably reduced, as well as pro-inflammatory cytokines such as G-CSF, IL-1 α , and MCP-1 after a single systemic injection of human MSC exosomes was given to patients with acute liver failure brought on by CCl₄. Additional research revealed that huc-MSC exosomes prevented oxidative stress-induced apoptosis and stimulated ERK1/2 activation and Bcl2 expression. It is interesting to note that GPX1 knockdown hindered the function of huc-MSC exosome-induced liver recovery and for this reason, GPX1 is crucial for the antioxidant activity and liver protection provided by huc-MSC exosomes.⁴⁸⁶ In comparison with the Bifendate group, which is often employed in clinical therapy to mitigate liver damage caused by CCl₄, the huc-MSC exosomes group exhibited better integrated hepatic tissue structure and a reduced

loss of hepatic lobules.⁵⁰⁶ Thus, the current research showed that huc-MSC exosomes had a hepatoprotective effect against the development of liver damage.

4.3.3 | Therapeutic intervention for hepatocellular carcinoma

The sixth most prevalent cancer, and the one with the highest death rate, is hepatocellular carcinoma (HCC). EVs have been utilized in various trials to treat HCC cells. Hepatic stellate cells, stem cells, HCC cells, hepatocytes, and fat-free milk from cows are a few examples of several sources of EVs. The bulk of research involved loading miRNA, or other therapeutic compounds, into EVs by using an endogenous loading technique. Before EV shedding, the target donor cells are altered using the endogenous loading technique, also

TABLE 15 Mesenchymal stromal cells (MSCs) derived molecules (e.g., exosome) in liver failure preclinical models, especially acute liver failure (ALF).

Sources	Model	Intervention	Result (reference)
Umbilical cord	Mice	MSCs exosome	GPX1-enriched exosomes diminished oxidative stress and also apoptosis ⁴⁸⁶
Placenta	Rat	MSCs exosome	CRP-enriched exosome provoked angiogenesis by upregulation of Wnt signaling axis ⁴⁸⁷
Bone marrow	Rat	MSCs exosome	Stimulation of hepatoprotective impacts by exosome-rich fractionated secretome ⁴⁸⁸
Bone marrow	Mice	MSCs exosome	Suppression of NLRP3 in macrophages and thereby reducing ALF by TNF- α pretreated exosome ⁴⁸⁹
Menstrual blood	Mice	MSCs exosome	Liver function recovery, improved survival rates, and suppressed hepatocellular apoptosis ⁴⁹⁰
Umbilical cord	Mice	MSCs extracellular vesicles	Inhibition of T-cell activation in liver tissue following reserve of CD154 expression ⁴⁹¹
Bone marrow	Mice	MSCs-conditioned medium	Promoting hepatocyte proliferation, inhibition of their apoptosis, hindrance of the infiltration of macrophages, improving Th2/Th1 ratio, and enabling hepatic stellate cell (HSC) loss ⁴⁹²
Bone marrow	Rat	MSCs-conditioned medium	Marked attenuation of panlobular immune cells infiltrates and also hepatocellular apoptosis ⁴⁹³
ESCs-MSCs	Mice	MSCs-conditioned medium	Supporting hepatocyte growth by VEGF enriched conditioned medium ⁴⁹⁴
Bone marrow	Mice	MSCs-exosome	Attenuation of liver inflammation by exosomal miR-20a-5p/ intracellular CXCL8 axis ⁴⁹⁵
Bone marrow	Rat	MSCs-conditioned medium	Reduced hepatocyte apoptosis ⁴⁹⁶
Bone marrow	Rat	MSCs-conditioned medium	Improving the hepatoprotective impacts of the conditioned medium by SMGO potentially elicited through inhibition of inflammation and loss of hepatocytes ⁴⁹⁷
Amniotic fluid	Mice	MSCs-conditioned medium	Hepatic progenitor-like (HPL)-CM showed superiority over amniotic fluid-MSCs in terms of liver recovery ⁴⁹⁸

Note: Reprinted with permission from Shokravi et al.⁴⁸⁵

Abbreviations: CM, conditioned medium; CRP, C-reactive protein; ESCs, embryonic stem cells; GPX1, glutathione peroxidase1; IL-8 or CXCL8, interleukin 8; NLRP3, NLR family pyrin domain-containing 3; SMGO, silica magnetic graphene oxide; Th1/2, T helper 1/2; TNF- α , tumor necrosis factor- α ; VEGF, vascular endothelial growth factor.

known as preloading.^{507,508} Following the loading of therapeutic compounds into the donor cells, EV transference into the recipient HCC cell is made possible by coculturing the donor and recipient HCC cells. Every study so far has shown that their therapeutic payload can be delivered to target tumor cells and promote apoptosis in those cells,⁵⁰⁹ lessen the chemoresistance,⁵¹⁰ inhibit cellular proliferation,^{509,511-513} and decrease the movement of cells.⁵¹¹

In another study, recipient HCC cells were cocultured with donor HCC cells that had been transfected with genes encoding the sodium iodide symporter (NIS).⁵¹⁴ The NIS protein increased the sensitivity of transplanted HCC to I-131 ablation by increasing the toxicity of I-131 in those cells.⁵¹⁴

In various animal models, HCC xenograft tumors have been subcutaneously implanted into various animals in immunocompromised mouse models. The EVs were loaded with antitumoral molecules and intravenously injected via tail veins or via direct intratumoral injections. The therapeutic EVs resulted in reduced tumor diameters,^{509-511,515-519} enhanced apoptosis of tumor cells,^{509-511,515} as well as enhanced chemoresistance.^{510,516}

Apart from their function as therapeutic cargo, EVs have also been found to have an indirect application in the treatment of HCC. It has been shown that intraperitoneal injections of propofol stimulated tumor-associated macrophages to create miR-142-3p EVs, which is linked to a reduction in tumor development.⁵¹⁸ In one research study, mice with xenograft HCC were treated with an exosome-based tumor vaccination (dendritic cells triggered by tumor EVs).⁵¹⁹ The outcome showed that the xenograft tumor volume was successfully reduced by intravenous injection of an exosome-based tumor vaccination.⁵¹⁹ Because EVs may transfer therapeutic compounds into xenograft HCC, reducing tumor development, all existing in vivo investigations displayed the potential benefits of EVs as a therapeutic modality for HCC. EVs can also be utilized as a therapeutic cargo or as a tumor vaccination.

An article by Nimitrungtaewee et al.⁵²⁰ concluded that increasing data demonstrates the ability of HCC cells to communicate with one another using EVs in order to stimulate cell proliferation. Clinical research has previously shown that RNA cargo from HCC-derived EVs may serve as prospective serum biomarkers to aid in the diagnosis of

HCC. A portion of this EV-RNA is also linked to tumor size, suggesting that it might be used as a prognostic marker. Furthermore, data from research conducted in vivo and in vitro suggested that EVs may be used to reduce tumor size and development. A summary of the potential roles of EVs is shown in Figure 17.

4.3.4 | Viral and alcohol-induced hepatitis and cirrhosis

Liver illnesses, including viral hepatitis, alcoholic hepatitis and cirrhosis, nonalcoholic steatohepatitis, and hepatocellular carcinoma, impose a significant burden on many individuals globally. Worldwide, liver illnesses claim the lives of over 2 million individuals each year. The main causes of these deaths are hepatocellular carcinoma (HCC), complications from liver cirrhosis, and viral hepatitis (VH).⁵²¹

A study by Zhou et al.⁵²¹ focused on immune cell exosomes and their effects on liver disease treatments examining both the pathogenic and possible therapeutic functions of these exosomes in liver illnesses. Although the authors discussed the potential use of exosomes produced from different immune cell sources to treat liver

disease, further study is necessary, particularly in human subjects. Nevertheless, many benefits associated with exosomes have been found across a wide range of liver-related illnesses.

4.4 | Kidneys

Acute kidney injury (AKI) is a severe global issue that causes abrupt loss of renal function as a result of tissue destruction and then progresses to chronic kidney disease. It has a high incidence of morbidity and death. Numerous assaults, including nephrotoxic drugs, environmental toxins, ischemia, systemic inflammation, nephritis, and urinary tract obstruction, may cause AKI.⁵²² While research into viable treatments for AKI is ongoing, at this time, only simple supportive therapy is available. Regenerative medicine for kidney illnesses is not yet possible due to the multiple etiologies of renal damage and the significant complexity of the kidney structure. Nonetheless, modern stem cell treatments have been utilized in several preclinical models and clinical studies over the last few decades due to the immense promise that stem cell technology has demonstrated.^{523–525} More significantly, EVs made from stem cells have attracted much more attention as a novel




FIGURE 17 Roles of HCC-derived EVs, HCC-derived EVs as a biomarker, and the evidence of therapeutic EVs from currently available reports. HCC cells secrete EVs that can lead to increased tumor cell proliferation, migration, chemoresistance, and decreased tumor cell apoptosis. They can also affect tumor microenvironments such as increased angiogenesis. Some of these HCC-derived EVs can be detected in circulation, making them available for use as a diagnostic biomarker. Moreover, it is possible for EVs to be used as a therapeutic cargo to transfer therapeutic molecules into tumor cells. Several in vitro and in vivo reports have demonstrated antitumoral effects using this method. ANGPT2, angiotensinogen; Circ, circular RNA; linc, long interceding/intergenic non-coding RNA; miR, microRNA; siRNA, signal interference RNA; TUC, tumor ultra-conserved RNA. Reprinted with permission from Nimitruntawee et al.⁵²⁰

therapeutic alternative for improved healing from many kinds of kidney damage without requiring allogeneic stem cells.⁵²⁶

In addition to their therapeutic possibilities, exosomes have also been studied for their potential as biomarkers for the identification of AKI.⁵²⁷ A research paper titled: "Unconscious Cells and Sudden Kidney Damage: Potential Therapeutic Avenue for Renal Repair and Regeneration"⁵²⁸ highlighted the effects of exosomes and how they affect multiple pathways of tissue regeneration including their role in (1) attenuating inflammation and improving immune modulation, (2) aiding in cell proliferation, (3) improving oxidative stress, (4) autophagy, and (5) minimizing cell death. Multiple lines of evidence substantiate the renoprotective impact of EVs derived from various sources in mitigating renal injury in diverse experimental models of AKI as presented in Table 16.⁵²⁸

4.4.1 | Bone marrow MSC-derived EVs for kidney therapy

A number of AKI models have shown the protective function of EVs produced from BMSCs. Bruno et al.⁵²⁹ demonstrated that by transporting certain cellular mRNAs, human BMSC-derived EVs enhanced the growth and minimized apoptosis of renal proximal tubular cells and helped mice recover from glycerol-induced AKI. Additionally, by reducing inflammation and apoptosis, a single injection of BMSC-EVs soon after an ischemia-reperfusion (I/R) damage-induced acute kidney injury promoted kidney recovery and enhanced function.^{530,531} Collino et al.⁵³⁴ showed that miRNAs transported by BMSC-EVs played a significant role in the recovery process after AKI. Specifically, miR-199a-3p from BMSC-EVs inhibits the Akt and Erk1/2 signaling pathways to prevent renal I/R damage and apoptosis.⁵³⁵ BMSC-EVs also improved renal function and morphological abnormalities in AKI produced by gentamicin or cisplatin.^{532,533} Importantly, Gregorini et al.⁵⁵⁹ showed that by modifying the expression of genes involved in membrane transport and cell energy metabolism, the immediate pre-transplantation incubation of donated rat kidneys with BMSC-derived EVs decreased ischemia damage.⁵⁵⁹

4.4.2 | Umbilical cord MSC-derived EVs for kidney therapy

In many AKI models, the administration of EVs generated from umbilical cord MSC (UC-MSC) was useful in the treatment of kidney injury. In rats, UC-MSC-EVs injected underneath the renal capsule prevented cisplatin-induced AKI.⁵³⁶ In vitro and in vivo cell proliferation was promoted by injecting UC-MSC-EVs, which also reduced oxidative stress and apoptosis.⁵³⁶ Additionally, UC-MSC-EVs enhanced kidney function in vivo and reduced renal NRK-52E cell production of caspase 3 and the p38/MAPK pathway activation in vitro.⁵³⁶ UC-MSC-derived EVs also provided protection against kidney damage, inflammation, and apoptosis caused by cisplatin.⁵⁴³ In addition to enhancing renal function, UC-MSC-derived EVs

demonstrated protection against sepsis-induced AKI by reducing tubular cell death and inflammation. In particular, NF- κ B inhibition and miR-146b overexpression in renal tubular cells mediated the protective effects of EVs.¹³³ Interestingly, UC-MSC-EVs also prevented inflammation, apoptosis, and kidney damage in rats after I/R by suppressing CX3CL1 and reducing macrophage accumulation,⁵³⁷ and also demonstrated antioxidative qualities.⁵⁴⁰ Evidence from recent research suggests that UC-MSC-derived EVs control kidney angiogenesis in a way that is independent of HIF-1. This suggests that these EVs enhance kidney function after unilateral I/R by decreasing apoptosis and increasing proliferation and angiogenesis.⁵⁴² The pro-angiogenic payload of EVs, which includes RNAs and VEGF, may account for their capacity for regeneration.⁵⁴² Additionally, UC-MSC-EVs accelerate tubular cell dedifferentiation and proliferation by upregulating HGF expression. This is probably due to the EVs' stimulation of Erk1/2 signaling while transferring RNA to damaged tubular cells.⁵³⁸

Through the use of miR-30b/c/d, human UC-MSC-derived EVs prevented I/R-induced kidney damage by preventing mitochondrial fragmentation and lowering apoptosis,⁵³⁹ as well as miR-125b-5p/p53,⁵⁴⁵ stimulation of the Nrf2/ARE system,⁵⁴¹ and OCT-4-mediated Snail pathway recruitment in RPTECs.⁵⁴⁴ According to in vivo imaging data, EVs preferred to settle near renal proximal tubular cells in ischemia-injured kidneys.⁵⁴⁵

4.4.3 | Placental tissue MSC-derived EVs for kidney therapy

Liu et al.⁵⁴⁸ examined the function of EVs produced from human placental MSC (hP-MSC) in AKI and demonstrated that they had a greater impact when encapsulated in a collagen matrix prior to intrarenal delivery. The collagen matrix promoted renal tubular cell proliferation and angiogenesis while preventing apoptosis and endoplasmic reticulum stress. It also enhanced the retention and therapeutic efficiency of hP-MSC-EVs in I/R-induced AKI.⁵⁴⁸ Similarly, Zhang et al.⁵⁴⁹ created hydrogels known as RGD (Arg-Gly-Asp) to control the stability and retention of the EVs. RGD bound to integrins on the surface of the MSC-EV membrane mediated the interaction between EVs and hydrogel. In I/R-induced AKI, EV-RGD hydrogels effectively prevented renal damage and reduced their morphological damage by encouraging autophagy and proliferation in tubular epithelial cells.⁵⁴⁹

4.4.4 | Adipose tissue MSC-derived EVs for kidney therapy

Adipose MSC-derived EVs (ADSC-EVs) have also been studied in relation to AKI recovery and regeneration. ADSC-EVs significantly reduced oxidative stress, apoptosis, and inflammation while also promoting renal angiogenesis thereby improving kidney dysfunction from I/R damage. An even better result than with EVs alone

TABLE 16 Application of EVs as therapeutic agents in acute kidney injury.

EV source	AKI model	EV cargo	Signaling pathway	Mechanism	Administration	References
BMSCs	Glycerol	mRNA	n/i	Proliferation, apoptosis	Intravenous	529
	I/R injury	RNA	n/i	Proliferation, apoptosis	Intravenous	530
	I/R injury	CCR2	NF-κB p65	Inflammation	Intravenous	531
	Cisplatin	n/i	n/i	Proliferation, apoptosis	Intravenous	532
	Gentamicin	RNA	n/i	Proliferation, apoptosis	Intravenous	533
	Glycerol	miRNA	n/i	Inflammation	Intravenous	534
	I/R injury	miR-199a-3p	Akt, Erk1/2	Apoptosis	Intravenous	535
UC-MSCs	Cisplatin	n/i	p38/MAPK, Erk1/2	Oxidative stress, apoptosis, proliferation	Renal capsule	536
	I/R injury	n/i	CX3CL1	Apoptosis, inflammation	Intravenous	537
	I/R injury	HGF/RNA	Erk1/2	Proliferation, apoptosis	Intravenous	538
	I/R injury	miR-30b/c/d	n/i	Apoptosis	Intravenous	539
	I/R injury	n/i	NOX2/gp91	Oxidative stress, apoptosis, proliferation	Intravenous	540
	I/R injury	n/i	Nrf2/ARE	Oxidative stress, apoptosis	Intravenous	541
	I/R injury	VEGF, RNAs	n/i	Apoptosis, proliferation, angiogenesis	Intravenous	542
	Cisplatin	n/i	n/i	Inflammation, apoptosis, autophagy	Renal capsule	543
	I/R injury	Oct-4	Snail	Apoptosis, proliferation	Intravenous	544
	Sepsis	miR-146b	NF-κB	Apoptosis, inflammation	Intravenous	133
	I/R injury	miR125b-5p	p53	Apoptosis, proliferation	Intravenous	545
AD-MSCs	I/R injury	n/i	n/s	Inflammation, apoptosis, oxidative stress, angiogenesis	Intravenous	546
	Sepsis	n/s	SIRT1	Apoptosis, inflammation	Intravenous	547
P-MSCs	I/R injury	n/i	n/i	Proliferation, angiogenesis, apoptosis	Intrarenal	548
	I/R injury	Let-7a-5p	n/i	Proliferation, apoptosis, autophagy	Intrarenal	549
K-MSCs	I/R injury	mRNA	n/i	Proliferation, angiogenesis	Intravenous	550
	I/R injury	miRNAs	n/i	Proliferation	Intravenous	551
L-MSCs	Glycerol	n/i	n/i	Proliferation, apoptosis	Intravenous	552
u-EVs	Glycerol	miRNA, Klotho	n/i	Proliferation, inflammation	Intravenous	553
TECs	I/R injury	CD26	p53, p21	Proliferation, inflammation	Intravenous	554
USCs	I/R injury	miR-146a-5p	NF-κB	Apoptosis, inflammation	Intravenous	555
Mac	I/R injury	IL-10	mTOR	Inflammation, autophagy	Intravenous	556
EPCs	I/R injury	miRNAs	n/i	Proliferation, apoptosis	Intravenous	557
	Sepsis	miR-93-5p	H3K27me3/TNF-α	Inflammation, apoptosis	Intravenous	558

Note: Reprinted with permission from Kosanovic et al. [528](#)

Abbreviations: AD-MSCs, adipose tissue MSCs; AKI, acute kidney injury; ARE, antioxidant response element; BMSCs, bone marrow MSCs; EPCs, endothelial progenitor cells; EVs, extracellular vesicles; I/R, ischemia-reperfusion; K-MSCs, kidney resident MSCs; L-MSCs, liver resident MSCs; Mac, macrophages; MAPK, mitogen-activated protein kinase; MSCs, mesenchymal stem cells; mTOR, mammalian target of rapamycin; n/i, not investigated; n/s, not specified; P-MSCs, placental MSCs; TECs, tubular epithelial cells; UC-MSCs, umbilical cord MSCs; uEVs, renal derived EVs isolated from urine; USCs, urine-derived stem cells; VEGF, vascular endothelial growth factor.

was obtained when ADSCs and ADSC-EVs were used in combination.⁵⁴⁶ Through the SIRT1 signaling pathway, ADSC-derived EVs reduced renal inflammation and apoptosis in a sepsis-induced AKI model.⁵⁴⁷

4.4.5 | EVs derived from other sources for kidney therapy

There is growing evidence that EVs produced from macrophages, kidney MSCs, liver MSCs, tubular epithelial cells, or endothelial progenitor cells (EPCs) are advantageous in experimental AKI.

In I/R-induced AKI, kidney-derived MSC-EVs were shown to enhance kidney function and structural alterations by transferring mRNA with pro-angiogenic characteristics, which in turn promoted the proliferation of peritubular capillary endothelial cells and prevented the loss of peritubular microvessels.⁵⁵⁰ Glomerular-MSC-EVs activated cell proliferation and transferred miRNA cargo to protect against I/R-induced AKI.⁵⁵¹ Additionally, CD26-containing tubular epithelial cell-derived EVs reduced inflammation and promoted cell proliferation by lowering p53 and p21, therefore mitigating I/R-induced AKI.⁵⁵⁴

It is interesting to note that EVs produced from human urine stem cells (USCs) prevented I/R-induced AKI and reduced inflammation and apoptosis by transferring miR-146a-5p and consequently reducing NF- κ B activation.⁵⁵⁵ Similarly, EVs generated from liver MSCs ameliorated mice's glycerol-induced AKI.⁵⁵² Normal urine-derived EVs from kidneys (uEVs) were shown to enhance the recovery from glycerol-induced AKI by promoting tubular cell proliferation, re-establishing endogenous Klotho levels, and reducing inflammation via miRNA cargo and Klotho transfer to resident kidney cells.⁵⁵³

EPC-secreted EVs stimulated renal regeneration in I/R-induced AKI by delivering miRNA cargo to local tubular epithelial cells and thus triggering regenerative mechanisms.⁵⁵⁷ In addition, EVs released by EPCs containing miR-93-5p were shown to provide protection against sepsis-induced AKI by reducing vascular leakage, inflammation, and apoptosis. This protective effect was achieved via the modulation of the H3K27me3/TNF-axis.⁵⁵⁸

Ultimately, when EVs obtained from macrophages were loaded with IL-10, it led to a successful delivery to renal tubular cells and macrophages in damaged kidneys and improved durability of vesicles. This process helped alleviate kidney injury, suppress inflammation, and stimulate mitophagy by inhibiting mTOR signaling.⁵⁵⁶

At present, a list of clinical trials is found on www.clinicaltrials.gov investigating the use of exosomes for treating various kidney related illnesses. Exosomes and EVs generated from umbilical cord MSCs were shown in a phase II/III clinical trial to be effective in treating chronic kidney disease and slowing disease progression. Serum creatinine, blood urea, and urine albumin creatinine ratio were significantly improved in patients with stage III and IV chronic renal disease who received eGFR-EVs made from cord tissue MSC.⁵⁶⁰

4.4.6 | MSC-Exos for chronic kidney disease

Chronic kidney disease (CKD) affects 10% of the population globally with many patients unaware of their condition.⁵⁶¹ While the etiology of CKD may differ, diabetes and hypertension remain the predominant factors.⁵⁶¹ Regardless of the many factors that contribute to the initial kidney damage, the development of renal fibrosis is a shared characteristic of all types of CKD.⁵⁶²

In experimental CKD, MSCs have shown encouraging effectiveness in reducing kidney damage.^{563,564} MSC-Exos, which share repair capabilities with MSCs, have been extensively used in the treatment of CKD, including DKD and renal fibrosis. The preclinical experiments included numerous CKD animal models, varied doses and timings, and several delivery techniques (tail infusion, organ perfusion, or direct application in the kidney), yet each study showed promising outcomes when utilizing EVs/exosomes.

4.4.7 | MSC-Exos for diabetic kidney disease

Chronic kidney disease (CKD), a consequence of diabetes mellitus (DM) affecting small blood vessels, is the most prevalent type of CKD and is expected to rise significantly worldwide.⁵⁶¹ Microalbuminuria is a common early sign in DKD, but it may not always occur, and it consequently indicates a higher likelihood of developing kidney damage over time. Diabetic individuals with renal disease have a higher risk of death compared to those without renal impairment. Given the projected rise in the worldwide adult population affected by DM, from 8.8% in 2015 to 10.4% in 2040, the significance of DKD and need for therapeutic options to manage or treat the disease is eminently needed.⁵⁶¹

Hyperglycemia induces the activation of various inflammatory pathways through several mechanisms, including the direct generation of reactive oxygen species, oxidative stress, stimulation of the renin-angiotensin-aldosterone system (RAAS), secretion of profibrotic cytokines like TGF- β , and formation of advanced glycation end-products.⁵⁶⁵ All of these result in albuminuria, podocyte and tubular damage, and apoptosis. Progressive fibrosis results from the accumulation of ECM proteins, including collagen and fibronectin, in the tubulointerstitium and renal mesangium due to an increase in matrix protein synthesis and a reduction in protein breakdown.^{565,566} The potential therapeutic effects of MSC-Exos in STZ-DKD has been quite frequently investigated using in vivo models of the disease (mice or rats) as well as in vitro podocyte, tubular epithelial cell (TEC), and glomerular endothelial cells. The effectiveness of MSC-Exos in alleviating DKD in preclinical models is highlighted in [Table 17](#).

4.4.8 | MSC-Exos for kidney fibrosis

Commonly, kidney fibrosis, namely tubulointerstitial fibrosis, is the ultimate result of almost all progressive forms of chronic kidney

TABLE 17 Summary of therapeutic effects of MSC-Exos from various sources in preclinical models of DKD.

MSC source	Model	Dose	Administration	Effects	Mechanism of action	References
Rat bone marrow	In vivo: STZ-induced DKD	Single: 5.3×10^7	Renal Subcapsular	↓ Renal tubule expansion, vacuolation, tubular atrophy ↓ Degeneration	↓ TGF- β ZO-1 was maintained ↓ Inflammatory cell Infiltration	566
	In vitro: Primary TECs	Not stated	Co-culture	↓ Degeneration ↓ Apoptosis		
Rat bone marrow	STZ-induced DKD	100 μ g/kg once per day \times 4 weeks	Intravenous (tail vein)	↑ Autophagy: ↑ LC3-II/LC-I, p62, Beclin-1 ↓ BUN, Scr, Glu, proteinuria at 10 and 12 weeks ↓ Fibrosis	↓ mTOR, S6K1, p62 ↓ Collagen, FN ↓ TGF- β	568
Mouse adipose	In vivo: Spontaneous diabetes	Single: not stated, 12-week therapy	Intravenous (tail vein)	↓ BUN, creatinine, proteinuria ↑ Autophagy ↓ Podocyte apoptosis	↑ miR-486 ↓ Smad1/mTOR activation ↓ Cleaved caspase 3	569
	In vitro: HG-treated MPC5	25 μ g/mL for 48 h	Co-culture	↑ Cell viability ↓ Apoptosis		
Mouse adipose	In vitro: HG-treated MPC5	Not stated	Co-culture	↓ Podocyte EMT ↑ miR-215-5p, -879-5p, -3066-5p, -7a-5p	↓ ZEB2 transcription	570
Adipose	In vivo: STZ-induced DKD	50 μ g twice weekly \times 3	Intravenous (caudal vein)	↓ Glu, Scr, UACR, kidney/body weight ↓ Mesangial hyperplasia ↓ Kidney fibrosis	Delivery of miR-125a ↓ HDAC1, ET-1	571
	In vitro: HG-treated rat GMC	Not stated	Co-culture	↓ IL-6, Col I and FN ↑ Bcl-2 and Bax		
Human umbilical cord	In vitro: HG-treated HK2, NRK-52E and hrGECs	25, 50, and 100 μ g/mL for 24 h	Co-culture	↓ TGF- β , IL-6, IL-1 β , and TNF- α	Secretion of EGF, FGF, HGF, and VEGF	572
Human urine	In vivo: STZ-induced DKD	Multiple: 100 μ g weekly \times 12	Intravenous (tail vein)	↓ Urine volume, albuminuria ↓ Apoptosis of podocyte and tubular cells ↑ Glomerular endothelial cell proliferation ↑ Angiogenesis	↓ Caspase-3 Delivery of VEGF, TGF- β 1, angiogenin, BMP-7	573
	In vitro: HG-treated immortalized human podocytes	5, 10, and 50 μ g/mL for 72 h	Co-culture	↓ Podocyte apoptosis		

TABLE 17 (Continued)

MSC source	Model	Dose	Administration	Effects	Mechanism of action	References
Human urine	In vivo: STZ-induced DKD	100 µg once weekly × 12	Intravenous (tail vein)	↓ Glu, KW, BUN, Scr, Ucr ↓ Podocyte injury ↓ Apoptosis	↓ VEGFA, MCP-1, TGF-β1 and TNF-α ↓ Bax and Caspase-3	574
	In vitro: HG-treated human podocytes		Co-culture	↑ Cell viability ↓ Apoptosis		

Note: Reprinted with permission from Cao et al.⁵⁶⁷
Abbreviations: AD-MSCs, adipose-derived mesenchymal stem cells; Bax, Bcl-2-associated X protein; Bcl-2, B-cell lymphoma 2; BMP-7, bone morphogenetic protein-7; BUN, blood urea nitrogen; CK1d, casein kinase 1d; COL1, collagen-1; DKD, diabetic kidney disease; EMT, epithelial–mesenchymal transition; ET-1, endothelin-1; FGF, fibroblast growth factors; FN, fibronectin; Glu, blood glucose; GMC, glomerular mesangial cell; GSH, glutathione; HDAC1, histone deacetylase 1; HG, high glucose; HGF, hepatocyte growth factor; IL-6, interleukin-6; KW, kidney weight; LC3, microtubule-associated protein light chain 3; MCP-1, monocyte chemoattractant protein-1; miR, microRNA; MSC, mesenchymal stem cells; mTOR, mammalian target of rapamycin; ROS, reactive oxygen species; S6K1, ribosomal protein S6 kinase beta-1; Scr, serum creatinine; STZ, streptozotocin; TGF-β, transforming growth factor-β; TECs, tubular epithelial cells; TGF-βR1, transforming growth factor-β type 1 receptor; TNF-α, tumor necrosis factor-α; Ucr, urine creatinine; VEGF, vascular endothelial growth factor; ZO-1, tight junction protein 1; 2K-1C, 2 kidneys, 1 clip model.

disease.⁵⁶² Tubulointerstitial fibrosis histology is distinguished by the accumulation of extracellular matrix in the interstitium. This accumulation is associated with damage to tubular cells, activation and proliferation of fibroblasts, and a decrease in the density of blood vessels surrounding the tubules.⁵⁶² Through a variety of pathways, TGF-beta has been shown to be a key profibrotic factor in several investigations.⁵⁷⁵
Recently, the antifibrotic efficacy of MSC-Exos has been assessed using a variety of rodent models of CKD, including diabetic and hypertensive CKD models, and ischemia-reperfusion injury (Table 18). Unilateral ureteral obstruction causes significant renal damage, which manifests as decreased glomerular filtration rate and renal blood flow within 24 h. Interstitial inflammation peaks 2–3 days later, and tubular dilatation, tubular atrophy, and fibrosis appear a week later. It results in severe interstitial renal fibrosis with excessive ECM buildup, tubular cell death, resident renal cell phenotypic change, and macrophage infiltration into the interstitial space.⁵⁷⁶ Table 18 summarizes the use of MSC-Exos from various sources on antifibrotic effects in kidney fibrosis.

4.5 | Ovaries

While not as commonly investigated as some of the other organs, ovaries have also been a target of exosome-related therapies. In a study titled: “Mesenchymal stem cells therapy: An auspicious approach for the management of uterine scarring and early ovarian failure,” Gao and colleagues investigated stem cells from various sources and their impact on ovarian regeneration and improvements in scarring.⁵⁸³ While most of the research to date has been on stem cells, few studies have investigated exosomes with the authors hinting at their low immunogenicity as a key advantage. According to a study by Yang et al., exosomes produced by BMSCs considerably improved the estrous cycle, augmented the quantity of basal and sinus follicles, and modulated the levels of reproductive hormones in rats suffering from premature ovarian failure. The enhancement was linked to the suppression of cell death in granulosa cells by specifically targeting phosphatase and tensin homologs.⁵⁸⁴ Moreover, the existence of miRNAs in BMSC-derived Exos might play a significant role in their ability to resist fibrosis while repairing intrauterine adhesions in the endometrium.^{585,586} Exosomes have the potential to cure a variety of ovarian illnesses, albeit research in this area of medicine remains in its early stages.⁵⁸³

5 | DEGENERATIVE PROCESSES

While many of the upcoming sections could certainly be attributed to the degeneration of tissues, this section focuses specifically on long-term degeneration tissues and their common diseases. Exosomes have played an important role in their improvements. These sections include primarily the effects of type 2 diabetes on the body as well as the number of issues related to respiratory disease/degeneration

TABLE 18 Summary of antifibrotic effects of MSC-Exos from various sources in preclinical models of kidney fibrosis.

MSC source	Model	Dose	Administration	Effects	Mechanism of action	References
Human umbilical cord	In vivo: UUO In vitro: NRK52E incubated with TGF- β	Single: 200 μ g Not stated	Left renal artery Co-incubation with isolated exosome	↑Renal function (↓ Scr, BUN) ↓ Tubular injury ↓ Tubulointerstitial fibrosis ↓ Apoptosis ↑proliferation ↓ Oxidative stress ↓ Apoptosis ↑Proliferation ↓ Oxidative stress	↓ ROS-mediated p38 MAPK/ERK signaling pathway ↓ Bax, cleaved caspase-3 ↓ ROS, MDA ↑ antioxidants: GSH	577
Human bone marrow	In vivo: UUO In vitro: NRK52E incubated with TGF- β	Single: Released from 1×10^6 MSCs Not stated	Intravenous Co-incubation with isolated exosome	Exosomes home to injured kidneys ↓ Fibrosis ↓ Fibrosis	Delivery of miR-let7c ↓ Collagen, MMP-9, α -SMA, TGF- β R1	559
Human bone marrow (transfected with anti-let-7i-5p)	In vivo: UUO In vitro: NRK52E incubated with TGF- β	Single: 1 mg/kg Not stated	Intravenous MSC on Transwell with NRK52E grown on the lower chamber	↑Renal function (↓ BUN, ↓ Scr, ↓ Ucr, ↑eGFR) ↓ Fibrosis ↓ TGF- β 1-induced fibrogenic responses ↓ EMT ↓ Tubulointerstitial fibrosis	↓ Let-7i-5p ↓ Collagen, FN, α -SMA, ↑TSC1 ↓ Phosphorylation of mTORC1, p70S6K and 4E-BP1 Exosomes delivered CK1 δ and β -TRCP to degrade YAP	578
Human umbilical cord	In vivo: UUO	Single: 200 μ g	Intravenous	↓ PTC rarefaction ↓ Tubulointerstitial fibrosis ↑ Endothelial function, angiogenesis ↓ HUVEC injury ↓ Apoptosis ↑ Endothelial angiogenesis	↑SIRT1/p-eNOS ↓ α -SMA ↑ VEGF, ↓ HIF-1 α	579
Adipose (transfected with GDNF)	In vivo: UUO In vitro: HUVEC against H/SD	Single: 200 μ g Single: 100 μ g/mL	Caudal vein Co-incubation with isolated exosome	↓ PTC rarefaction ↓ Tubulointerstitial fibrosis ↑ Endothelial function, angiogenesis ↓ HUVEC injury ↓ Apoptosis ↑ Endothelial angiogenesis	↑SIRT1/p-eNOS ↓ α -SMA ↑ VEGF, ↓ HIF-1 α	580
Adipose	In vivo: IRI In vitro: primary TECs stimulated with TGF- β	Single: 100 μ g Not stated	Caudal vein Co-incubation with isolated exosome	↑ Tubular proliferation, regeneration ↓ Interstitial fibrosis ↓ Inflammation ↓ TGF- β 1-induced transformation of TECs to pro-fibrotic phenotype	↑Sox9 ↓ α -SMA, PDGFR- β	581
Adipose	2K-1C Unilateral renal artery stenosis	Single: 100 μ g	Caudal vein	↓ HIF-1 α Stabilized systolic blood pressure ↑ Natriuresis ↓ Fibrosis ↓ Inflammation	↓ Collagen, TGF- β ↓ IL-10	582
Pluripotent stem cell	In vivo: UUO In vitro: NRK-52E	Single: 10^{11} particles/mL $10^6/10^7/10^8$ particles/m	Tail vein Co-incubation with isolated exosome	↓ Fibrosis ↓ Inflammation ↓ Col-1, α -SMA ↑E-cadherin	↑SIRT6 ↓ β -catenin	544

Note: Reprinted with permission from Cao et al.⁵⁶⁷

Abbreviations: AD-MSCs, adipose-derived mesenchymal stem cells; Bax, Bcl-2-associated X protein; Bcl-2, B-cell lymphoma-2; BMP-7, bone morphogenetic protein-7; BUN, blood urea nitrogen; CK1 δ , casein kinase 1 δ ; COL-1, collagen-1; DKD, diabetic kidney disease; EMT, epithelial-mesenchymal transition; ET-1, endothelin-1; FGF, fibroblast growth factors; FN, fibronectin; GDNF, glial-derived neurotrophic factor; GMC, glomerular mesangial cell; GSH, glutathione; HDAC1, histone deacetylase 1; HG, high glucose; HGF, hepatocyte growth factor; H/SD, hypoxia/serum deprivation; HUVECs, human umbilical vein endothelial cells; IL-6, interleukin-6; IRI, ischemia/reperfusion injury; LC3, microtubule-associated protein light chain 3; miR, microRNA; KW, kidney weight; MCP-1, monocyte chemoattractant protein-1; MDA, malondialdehyde; MSC, mesenchymal stem cells; mTOR, mammalian target of rapamycin; mTORC1, mammalian target of rapamycin complex 1; PDGFR- β , platelet-derived growth factor receptor beta; ROS, reactive oxygen species; S6K1, ribosomal protein S6 kinase beta-1; Scr, serum creatinine; SIRT1, sirtuin-1; Sox9, SRY-box transcription factor 9; STZ, streptozotocin; TECs, tubular epithelial cells; TGF- β , transforming growth factor- β ; TGF- β R1, transforming growth factor- β type 1 receptor; TNF- α , tumor necrosis factor- α ; Ucr, urine creatinine; VEGF, vascular endothelial growth factor; ZO-1, tight junction protein-1; 2K-1C, 2 kidneys, 1 clip model; YAP, yes-associated protein; UUO, unilateral ureteral obstruction.

over time. Exosomes have played a marked impact on the improvement of all tissues, as highlighted below.

5.1 | Type 2 diabetes mellitus

Many review articles have been written on the use of exosomes in the treatment of type 2 diabetes mellitus.^{587,588} Globally, the incidence of diabetes mellitus (DM) is rising at an alarming pace. More than 30.3 million people in the United States suffer from DM and 91.2% of them are type 2 diabetes mellitus (T2DM), which disproportionately affects African Americans and Hispanics. T2DM is the primary cause of morbidity and death for diabetic individuals and a significant risk factor for cardiovascular disease (CVD). Intensive glucose control has not been able to stop the development of macro- and microvascular-associated mortality in this population, despite major advancements in T2DM therapy. This emphasizes the need to better understand the underlying molecular pathways that contribute to CVD in the context of type 2 diabetes. The development of vascular problems brought on by diabetes, such as diabetic nephropathy (DN) and CVD, is significantly influenced by endothelial dysfunction (ED). Furthermore, as the costs of managing and treating type 2 diabetes and its multisystem consequences continues to climb, the troubling rise in the disease's prevalence in the United States and other countries has put a heavy burden on healthcare systems, policymakers, families, and caregivers.

The intricate molecular foundations of type 2 diabetes and its associated consequences are made up of an interconnected system of lifestyle, genetic, and epigenetic variables that function within a physical, social, and cultural context.⁵⁸⁹ In particular, it is generally believed that genetic predisposition, excessive energy-dense food intake, and inactivity cause T2DM by interfering with the feedback loops that regulate insulin action and secretion.⁵⁹⁰ Persistent hyperglycemia may arise from this disturbance, which may also affect β -cell activity, insulin signaling, and glucose metabolism. Insulin resistance and β -cell dysfunction both arise early in the etiology of type 2 diabetes and, over time, propel the development of IGT from normal glucose tolerance (NGT) to T2DM.⁵⁸⁹ Furthermore, the development of T2DM is influenced by several variables, such as population aging, intrauterine environment, alterations in the microbiota, and pollution.

5.1.1 | Exosomal miRNAs as markers of endothelial dysfunction

Exosomes may be detected in several body fluids, including blood, saliva, urine, tears, and lactation. This characteristic makes exosomes appealing as noninvasive biomarkers for the illness. Exosomes serve as distinctive markers of the cells they originate from and the physiological state of that cell.⁵⁹¹ Hence, the diverse concentrations

of exosomes and their abundant payload, including as proteins, mRNAs, and miRNAs, may serve as valuable indicators for predicting and diagnosing diseases. Furthermore, it may provide a valuable understanding of the intricate biological mechanisms involved in disorders like type 2 diabetes, as well as the underlying endothelial dysfunction that contributes to the vascular problems associated with this metabolic disorder. Tables 19 and 20 demonstrate examples of commonly utilized exosomal miRNAs and microRNAs used as biomarkers. In fact, nowadays, various exosomes derived from various body fluid sources have now been utilized for the detection of DM and its subsequent disease (Table 21).

5.1.2 | Exosomes, glucose, and lipid metabolism

Changes in the metabolism and absorption of glucose is the primary issue in diabetes, ultimately resulting in hyperglycemia.⁶⁴⁸ The ghrelin system has a role in improving glucose metabolism in the treatment of T1DM, and hyperbaric oxygen may influence the ghrelin system, hence impacting glucose metabolism.⁶⁴⁹ Clinical investigations on diabetic individuals have shown that GLP-1 may have a function in lowering blood glucose levels.⁶⁵⁰ Exosomes have been found to drastically reduce gluconeogenesis and promote the metabolism and absorption of fats and carbohydrates.⁶⁵¹ Notably in DM, autophagy suppression reduces the therapeutic effect of exosomes produced from MSCs. Further research showed that exosomes triggered AMP-activated protein kinase (AMPK) to increase autophagy by upregulating Beclin-1 and LC3-II, which in turn promoted lipid metabolism and glucose absorption.⁶⁵¹

5.1.3 | Exosomes and β -cell function

The pancreas contains a large number of Langerhans islets, whereby each islet is made up of around 1000 β cells and weighs 0.9 g in total.⁶⁵² The β cells are affected in diabetes mellitus and obesity, among other diseases. Hyperinsulinemia, or elevated insulin levels in the plasma of obese patients, is a compensatory mechanism that keeps insulin resistance at bay. Thus, the mass of β cells increases, resulting in hyperinsulinemia.⁶⁵³ An obese rat model demonstrated that obesity might increase β cell mass up to three fold.⁶⁵⁴ T1DM is characterized by the loss of β cells or their inability to secrete insulin and the mass of β cells in T1DM patients has been shown to decline to nearly 100% over an extended period of time.⁶⁵⁵⁻⁶⁵⁷

MSC-Exos have been used in an attempt to treat T1DM. After the exosomes were injected intravenously into animal models, insulin and glucose levels were assessed 6 weeks later. According to the findings,⁶⁵⁸ exosomes stimulated β -cell regeneration in the pancreas and increased their bulk which is crucial for increasing insulin release in type 1 diabetes. Apart from increasing insulin levels, MSC-Exos markedly reduced serum glucose levels.

TABLE 19 Extracellular miRNAs associated with type 2 diabetes mellitus and endothelial dysfunction.

miRNA	Normoglycemic	T2DM	Targets/pathways	Source	Complication	References
miR-126	Up	Down	SPRED-1 PIK3R2/p85- β -PLK4	Plasma	Impaired angiogenesis	592-594
miR-26a	Up	Down	TRPC6	Plasma	Impaired angiogenesis	595
miR-133b	Down	Up	MAPK/ERK signaling	Serum	Diabetic nephropathy	596,597
miR-342	Down	Up	SRY-box 6 (SOX-6)	Serum	Diabetic nephropathy	596,598
miR-30a	Down	Up	TGF- β 1 Becn1	Serum	Diabetic nephropathy	596,599
miR-326		Up	ADIPOR-2 (adiponectin)	Plasma	T2DM	600
Let-7a	Down		Glucose metabolism	Plasma	T2DM	600
Let-7f	Down		Glucose metabolism	Plasma	T2DM	600
miR-20b-5p	Down	Up	AKT-interacting protein	Serum	T2DM	601
miR-21-5p	Down	Up	WWP1 (WW domain- containing protein 1) Endothelial progenitor cell proliferation	Plasma	T2DM	602,603
miR-375-3p	Down	Up	β -cell function	Serum	T2DM	604
miR-362-3p	Down		ADAMTS1	Plasma	Atherosclerosis (CAD)	605
miR-15a	Down	Up	UCP-2	RBC	T2DM	606,607
miR-15b	Down	Up	TNF-alpha SOCS3	RBC	Pre-T2DM	606
miR-499	Down	Up	PTEN	RBC	Pre-T2DM	606
miR-7	Down	Up	mTOR signaling	Serum	Vascular complications	608
miR-25-3p		Up	CDH1 and PTEN	Plasma	Diabetic retinopathy	609
miR-320b	Down	Up	Angiogenesis	Plasma	Diabetic retinopathy	609
miR-495-3p	Down	Down		Plasma	Diabetic retinopathy	609
miR-34a	Down	Up	Sirt1	Mouse aortic endothelial cells	Endothelial dysfunction	610
miR-210	Down	Up	Cell proliferation	Serum	Diabetic retinopathy	611
miR-210	Up	Down	PTP1B	RBC	T2DM	612
miR-139-5p	Down	Up	c-Jun	Peripheral blood	T2DM	613

Note: Reprinted with permission from Fluitt et al.⁵⁸⁸

5.1.4 | Exosomes and insulin resistance

The management of patients with diabetes mellitus is becoming increasingly more difficult due to their insulin resistance, which causes a change in the insulin concentration–effect curve and a tendency for greater concentration levels to have a comparable effect.⁶⁵⁸ Commonly seen in T1DM and sometimes in obese individuals, insulin resistance is caused by cells that are not sensitive to insulin.

Insulin increases the expression and activity of GLUTs, including GLUT4, to improve the absorption of glucose by cells.⁶⁵⁹ Nonetheless, there are a few causes of insulin resistance. For example, fat buildup in obese individuals causes a reduction in adipokine release, an increase in lipolysis, and the production of pro-inflammatory cytokines. These effects result in elevated amounts of free glycerol and fatty acids linked to insulin resistance.⁶⁶⁰

Exosomes have been shown in experiments to have therapeutic effects by increasing insulin sensitivity, a crucial aspect of treating

diabetes mellitus. A high-fat diet, which includes fatty acids like palmitic acid (PA), may cause insulin resistance. High PA intake causes the hepatocytes to develop the insulin resistance characteristics. In addition to inhibiting IRS-2 phosphorylation and preventing glucose absorption as a consequence of insulin stimulation, the PA may cause apoptosis in INS-1 cells. The neutral ceramidase (NCDase)-enriched exosomes inhibit insulin resistance brought on by excessive PA consumption and halts the production of reactive oxygen species, perhaps helping to reduce INS-1 cell death.⁶⁶¹ In a different experiment, exosomes isolated from umbilical cord MSCs were used to treat T1DM. The authors demonstrated the ability of exosomes to increase insulin-induced glucose absorption, lasting for at least 48h. Additionally, exosomes lower leptin levels and dramatically increased SIRT1 and IRS-1 mRNA expression, demonstrating their ability to increase adipocyte insulin sensitivity.⁶⁶²

One of the most well-known cargos of exosomes are miRNAs, and the effect of exosomes on target cells will vary depending on

TABLE 20 Exosomal microRNAs associated with type 2 diabetes mellitus and endothelial dysfunction.

miRNA	Normaglycemic (concentration/excretion)	T2DM (concentration/excretion)	Targets/pathways	Source of exosomes	Method of isolation	Complication	References
miR-451	Low	High	YWHAZ CAB39	Urine	Ultra-centrifugation	Diabetic nephropathy	614–616
Let-7c-5p	Low	High		Urine	Differential centrifugation	Diabetic nephropathy	617
miR-192	Low	High	Egr1	Urine	Ultra-centrifugation	Diabetic nephropathy	618,619
Let-7e-5p	Low	High	FASLG (migration and tube formation of endothelial progenitor cells)	Urine	miRCURY™ exosome isolation kits (Qiagen)	Diabetic nephropathy	620–622
miR-15b	Low	High	TNF-alpha	Urine	Ultra-centrifugation	Diabetic nephropathy	623
miR-34a	Low	High	Growth arrest specific 1 (GAS1)				
miR-636	Low	High	Adipogenesis				
miR-30b-5p	Low	High	Epithelial-to-mesenchymal transition	Urine	Ultra-centrifugation	Diabetic nephropathy	620
miR-20-5p	Low	High	Wnt9b/β-catenin signaling pathway	Peripheral blood	Ultra-centrifugation	Wound healing angiogenesis	624,625
miR-126	Low	High	Angiogenesis vascular integrity	Mouse brain endothelial cells	ExoQuick-TC (exosome precipitation solution kit, system biosciences)	Stroke	454
miR-320c	Low	High	ADAMTS5 CDK6 TSP-4 BMP6	Urine	ExoQuick-TC (exosome precipitation solution kit, system biosciences)	Diabetic nephropathy	626
Let-7i-3p	Low	High	Wnt/β-catenin signaling cascade, activin	Urine	miRCURY™ exosome isolation kits (Qiagen)	Diabetic nephropathy	627
miR-24-3p	Low	High	receptor signaling, and cell differentiation and proliferation			Diabetic nephropathy	
miR-27b-3p	Low	High				Diabetic nephropathy	
miR-15b-5p	High	Low				Diabetic nephropathy	
miR-25-3p	Low	High	Endothelial cell proliferation	Plasma	ExoQuick-TC (exosome precipitation solution kit, system biosciences)	Diabetic retinopathy	609
miR-320b	Low	High	Angiogenesis				
miR-495-3p	Low	High					

Note: Reprinted with permission from Fluit et al.⁵⁸⁸

TABLE 21 Exosomes derived from body fluid can act as novel biomarkers for early diagnosis of DM and diabetic complications.

Disease	Target content in exosome	Sample	Method	Scientific mechanism	References
T2DM	Counts of cell-derived exosomes ↑	Serum	Flow cytometry meta-analysis	1. Total annexin V-positive blood cell microparticles—procoagulant activity could be involved in vascular complications. 2. Endothelial microparticles stimulated by elevated glucose change their molecular composition and increase their biological activity, which may lead to progressive endothelial damage and subsequent cardiovascular complications in diabetes	629–631
Diabetes nephropathy	Counts of cell-derived exosomes ↑	Urinary	Flow cytometry	Mir-26a-5p from adipose-derived mesenchymal stem cell-derived EVs protect against DN	632
	Dipeptidyl peptidase-IV ↑	Urinary	ELISA	The urinary level of microvesicle-bound microvesicle-dipeptidyl peptidase-IV is associated with the severity of diabetic kidney disease	633
	Wilms tumor-1 ↑	Urinary	Western blotting	Among podocyte-derived signal transduction factors in urinary exosomes, WT1 mRNA levels reflected damage of diabetic glomeruli in the patients	634
	AMBP, MLL3 ↑ VDAC1 ↓	Urinary	LC-MS/MS	Comparing DN urine exosomes and healthy controls, it was discovered in a panel of three proteins (AMBP, MLL3, and VDAC1) that they were differentially found in urinary exosomes from DN patients	635
	Mir-130, mir-145, mir-155, mir-424 ↑	Urinary	TaqMan qPCR	High glucose will stimulate mesangial cells and increase the content of mir-145 in mesangial cells and their derived exosomes	636
	Mitochondrial DNA ↓	Urinary	Intrarenal Gene Expression Analysis	Urine exosomes from patients with diabetes and CKD had less mitochondrial DNA, and kidney tissues from patients with diabetic kidney disease had lower gene expression of PGC1a	637
	Elf3 ↑	Urinary	Western blotting	AGE treatment induced the secretion of Elf3-containing exosomes from cultured podocytes, which was dependent on the activation of the TGF-β-Smad3 signaling pathway	638
	Mir-16 ↓	Urinary	RT-qPCR	Mir-16 identified as the most stable endogenous reference gene in data set, making it a suitable endogenous reference gene for miRNA studies of urinary exosomes derived from CKD patients	639
	Gelatinase ↓ ceruloplasmin ↑	Urinary	ELISA	Gelatinase (decreased activity) and ceruloplasmin (increased levels), in the urinary exosomes of diabetic kidney patients were in agreement with the alterations of these two proteins in the kidney tissue	640
	Counts of exosomes ↑	Blood	Flow cytometry	Exosomes from diabetic rats no longer activated the ERK1/2 and HSP27 cardioprotective pathway and were no longer protective in a primary rat cardiomyocyte model of hypoxia and reoxygenation injury. Exosomes from diabetic plasma have lost the ability to protect cardiomyocytes, but protection can be restored with exosomes from nondiabetic plasma	641
Diabetic cardiomyocytes	Hsp20 ↓	Serum	LC-MS/MS	Elevation of Hsp20 in cardiomyocytes can offer protection in diabetic hearts through the release of instrumental exosomes	642
	Mir-320 ↑	Serum	TaqMan qPCR	Cardiomyocytes exert an anti-angiogenic function in type 2 diabetic rats through exosomal transfer of mir-320 into endothelial cells	643
	Mir-126 ↓	Serum	TaqMan qPCR	Mir-126 targets insulin receptor substrate (IRS)-1 expression via PI3K/Akt signaling pathways suggests that it is involved in IR modulation	644
	Mir-7 ↑	Serum	RT-qPCR	Mir-7 was demonstrated to be involved in b-cell dysfunction and insulin secretion	608
Diabetic Charcot neuroarthropathy (CN)	Counts of exosomes ↑	Plasma	Flow cytometry	The concentration of EVs is related to elevation of markers of inflammation (CRP and foot temperature difference) in acute diabetic CN	645
	Counts of endothelial cell exosomes ↑	Serum, plasma	Western blotting, RT-qPCR	Exosomal Ang2 secretion is regulated by the PI3K/Akt/eNOS and syndecan-4/syntenin pathways	646,647

Note: Reprinted with permission from Sun et al.⁶²⁸

Abbreviations: AGE, advanced glycation end-product; CKD, chronic kidney disease; CRP, C-reactive protein; DM, diabetes mellitus; DN, diabetic nephropathy; EV, extracellular vesicle; LC-MS/MS, liquid chromatography–tandem MS; T2DM, type 2 diabetes mellitus.




FIGURE 18 Exosomes and insulin resistance/sensitivity in DM. The exosomes demonstrate dual function in DM and can mediate both insulin resistance and sensitivity. For instance, exosomes derived from M1 macrophages can induce insulin resistance and are rich in PI3K and Ptch; on the other hand, exosomes derived from mesenchymal stem cells and hepatocytes induce insulin sensitivity via affecting molecular pathways such as SIRT1 and Akt as well as transferring miRNAs. Reprinted with permission from Ashrafizadeh et al.⁶⁶⁵

the miRNAs' activity. It has been proposed that miRNA-3075 acts as an antidiabetic agent by improving insulin sensitivity which has been shown in hepatocyte-derived exosomes.^{663,664} The pro-inflammatory role of macrophages and the progression of insulin resistance are connected (Figure 18, Table 22).

5.1.5 | Exosomes and diabetic wound healing

In diabetic individuals, the healing process of wounds is slowed down, making chronic non-healing wounds much more common.⁶⁷³ Pro-inflammatory cytokines and proteases are abundant in diabetic wounds, decreasing growth factors in the area.^{674–676} Medications for diabetics are administered to about 70% of individuals with the disease. Inflammation, re-epithelialization, collagen deposition, angiogenesis, and proliferation are just a few of the many cellular and molecular processes involved in the intricate process of wound healing.^{677,678}

A recent systematic review published by Bailey et al. 2022 titled: "Extracellular Vesicles Derived from MSCs to Treat Diabetes-Related Injuries: A Comprehensive Review and Meta-Analysis of Preclinical Animal Research"⁶⁷⁹ discovered numerous benefits of exosomes in the treatment of diabetic wounds. Wound closure was the main result that was assessed, but secondary outcomes demonstrated many

advantages including (1) increase in blood vessel density and number, (2) re-epithelialization, (3) collagen deposition, (4) scar width, and (5) inflammation (Figure 19).⁶⁷⁹

5.1.6 | Diabetic-induced neuropathy

Due to focal and widespread nervous system impairments, neuropathy is seen in around 50% of DM patients. Diabetic individuals often have neuropathy, which may cause discomfort and a lower quality of life.⁶⁸⁰ In the USA, diabetic neuropathy is the cause of \$10 billion in expenses each year.⁶⁸¹ Consequently, actions aimed at managing and avoiding diabetic neuropathy should be taken. Notably, anti-diabetic drugs have shown promise in reducing brain damage caused by DM. Anti-diabetic medications have the ability to greatly reduce neuroinflammation and oxidative damage, hence preventing the onset of neurological disorders like Alzheimer's and Parkinson's diseases. Moreover, antidiabetic medications like GLP-1R and DPP-4i have the potential to enhance the process of neurogenesis and improve cognitive performance.⁶⁸²

Exosomes have been shown to possess an ability to lower apoptosis in neurons.⁶⁸³ Administering paeoniflorin enhances the therapeutic effects of Schwann cell-derived exosomes and supports the stability and integrity of the endoplasmic reticulum. Moreover, the

TABLE 22 Exosomes in regulating insulin resistance, β -cell function, glucose, and lipid metabolism.

Source of exosome	Function	Signaling network	Remarks	References
Hepatocyte	Insulin sensitivity	miRNA-3075/FA2H	The exosomes can deliver miRNA-3075 for decreasing FA2H expression and increasing insulin sensitivity	666
Human umbilical cord mesenchymal stem cells	Insulin sensitivity	SIRT1 IRS-1	Decreasing leptin levels Upregulating expression level of SIRT1 and IRS-1 in mediating insulin sensitivity	662
Mesenchymal stem cells	Insulin sensitivity	IRS-1 Akt GLUT4	Promoting glucose metabolism Increasing phosphorylation of Akt and IRS-1 Elevating membrane translocation of GLUT4	667
INS-1 cells	Insulin sensitivity	IRS-2	NCDase-enriched exosomes promote insulin sensitivity via enhancing IRS-2 phosphorylation Increasing insulin uptake in cells Preventing palmitic acid-induced insulin resistance	661
-	Insulin resistance	TLR4/NF- κ B	Silencing SIRT1 induces exosome secretion via autophagy inhibition Exosomes induce insulin resistance via triggering TLR4/NF- κ B axis	668
Adipocyte	Insulin resistance	Ptch PI3K	Exosomes possess high levels of Sonic Hedgehog Inducing M1 polarization of macrophages via Ptch and PI3K upregulation	669
Mesenchymal stem cell	β cell protection	miRNA-21/Grp94/apoptosis p38/caspase-3/PARP/apoptosis	The exosomes alleviate ER stress and decrease apoptosis in β cells Exosomes delivering miRNA-21 decrease Grp94 expression to inhibit apoptosis in β cells Exosomes downregulate p38 expression to prevent apoptosis	670
M1 macrophages	β cell dysfunction	miRNA-212-5p/SIRT2/Akt/ GSK-3 β / β -catenin	The exosomal miRNA-212 reduces SIRT2 expression to inhibit Akt/GSK-3 β axis Providing nuclear translocation of β cells Preventing insulin secretion by β cells	671
Mesenchymal stem cell	Glucose and lipid metabolism	AMPK/autophagy	Ameliorating T1DM via upregulating AMPK expression, promoting Bedlin-1 and LC3 levels, resulting in autophagy Autophagy induction enhances lipid and glucose metabolism in cells	672

Note: Reprinted with permission from Ashrafzadeh et al.⁶⁶⁵




FIGURE 19 Forest plot demonstrating increased wound closure rates of diabetic wounds receiving MSC-EVs. Note the positive impact of exosomes on wound closure in diabetic in vivo models. Reprinted with permission from Bailey et al.⁶⁷⁹

intervention of paeoniflorin enhances the ability of exosomes to inhibit programmed neuron cell death by reducing the expression of GRP78 and IRE1 α . These interactions are crucial for the prevention of neuropathy in individuals with diabetes mellitus.⁶⁸³ There is hope that exosomes produced by MSCs might alleviate diabetic peripheral neuropathy. It has been demonstrated that exosomes with a high concentration of miRNA-146a have the ability to increase the speed at which nerves transmit signals while also potentially reducing the threshold for feeling heat and pressure. To lessen diabetic neuropathy, these exosomes delivered via IV suppress endothelial cell activation and reduce monocyte inflammation. To lessen diabetic neuropathy, the TLR-4/NF- κ B axis may be suppressed by miRNA-146a-enriched exosomes.⁶⁸⁴

5.1.7 | Diabetic-induced nephropathy

Another DM consequence that leads to end-stage renal disease is nephropathy.⁶⁸⁵ Nephropathy occurs in 25% of patients and develops for a number of reasons. By producing Amadori and advanced glycation end-products, hyperglycemia may lead to diabetic nephropathy. Moreover, the electron transport cycle is triggered by hyperglycemia, which dramatically raises ROS levels and contributes to diabetic nephropathy.⁶⁸⁶

Exosomes may have a therapeutic effect on diabetic kidney disease. MSC-Exos have elevated concentrations of miRNA-125b, which inhibits TRAF6 and triggers Akt signaling. This leads to the activation of autophagy and a reduction in apoptosis, both of which are crucial for mitigating the symptoms of diabetic nephropathy.⁶⁸⁷ Autophagy is a crucial biochemical mechanism involved in cell death. In summary, autophagy facilitates the breakdown of aging organelles and macromolecules that are hazardous to primary cell homeostasis. Autophagy has context-dependent roles in both pro-survival and pro-death processes.^{682,688–692} Notably, pro-survival autophagy may stop a cell from going through apoptosis.⁶⁹³ The exosomes produced from MSCs also inhibit mTOR signaling, a negative regulator of autophagy, in order to significantly enhance the levels of Beclin-1

and LC3. These proteins play a crucial role in promoting autophagy and preventing apoptosis helping alleviate diabetic nephropathy.⁵⁶⁹

5.1.8 | Diabetic-induced endothelial dysfunction

Around the globe, gestational diabetes is prevalent in around 5%–20% of pregnancies. Endothelial dysfunction is one of the diseases that arise in gestational diabetes mellitus.⁶⁹⁴ Fetal-placental endothelial dysfunction is caused by increased production of nitric oxide (NO) and improved transport of L-arginine.^{695,696} While experiments are still in their infancy, exosomes lower the levels of a number of substances during disease progression, including IL-8, MDA, SOD, and—more importantly—ICAM-1, which helps avoid endothelial failure.⁶⁹⁷

5.1.9 | Diabetic-induced erectile dysfunction

One known risk factor for erectile dysfunction is diabetes mellitus which affects 19%–90% of individuals and may manifest up to 10 years before the onset of diabetes mellitus, lowering a man's quality of life and self-esteem.^{698–700} In addition to diabetes mellitus, additional risk factors for the development of erectile dysfunction include smoking, cardiovascular illnesses, dyslipidemia, hypertension, and obesity.⁷⁰¹ Vasculogenic variables that mediate arterial inflow or venous outflow problems are the most prevalent causes of erectile dysfunction.⁷⁰²

ADSC-Exos have been injected intravenously to increase smooth muscle level relative to collagen and increase intracavernous pressure, as shown by CD31 overexpression.⁷⁰³ In addition, exosomes reduce the degree of caspase-3 production while simultaneously enhancing the expression of Bcl-2 to inhibit apoptosis. Exosomes obtained from ADSCs have been shown to be effective therapeutic agents for relieving erectile dysfunction in individuals with diabetes mellitus.⁷⁰³ Exosomes generated from adipose stem cells have the ability to promote angiogenesis and speed endothelial cell

proliferation in the treatment of erectile dysfunction. Moreover, these exosomes decreased fibrosis in the corpus cavernosum.

Further research showed that large quantities of pro-angiogenic factors miRNA-126, -130a, and -132, as well as antifibrotic factors miRNA-let7b and -let7c, were present in exosomes produced from adipose stem cells.³¹⁷ According to the trials, the best way to treat erectile dysfunction brought on by diabetes is to avoid fibrosis. Smooth muscle cell exosomes increase the amount of smooth muscle and reduce the buildup of collagen to have an antifibrotic effect. In order to improve erectile dysfunction in diabetics, exosomes upregulate the expression levels of eNOS and nNOS to trigger the NO/cGMP axis.⁷⁰⁴

5.1.10 | Diabetic-induced cardiovascular diseases

Diabetes is also greatly linked to a number of cardiovascular disorders including atherosclerosis, which was previously discussed.^{705,706} By inhibiting TGF- β signaling, the exosomes derived from MSCs may lower the expression level of Smad2, therefore improving cardiac fibrosis in diabetic patients.⁷⁰⁷ These studies suggest that exosomes may prevent cardiac damage and help avoid DM-mediated heart damage.

5.1.11 | Diabetic-induced eye disorders

The protective effect of exosomes on the retina seems to depend on the route in which they are administered.⁷⁰⁸ Exosomes administered subconjunctivally provide superior outcomes than those administered intravenously and arrange the retina's cellular constituents in a well-organized fashion. The optimal method is thought to be intraocular delivery, which enhances retinal layers that resemble the natural retina. Hyperglycemia has been shown to decrease miRNA-222 expression. To slow down retinal damage in diabetic macular degeneration, MSC-Exos increase the expression of miRNA-222.⁷⁰⁸

5.1.12 | Summary of diabetic-induced complications

Diabetes affects millions of individuals worldwide, and the associated complications caused by diabetes are both numerous and difficult to treat. These studies show that a range of diabetic problems arise such as ocular disorders, cardiovascular illnesses, delayed wound healing, erectile dysfunction, endothelial dysfunction, and neuropathy which may be ameliorated by exosomes. Table 23 summarizes the use of MSC-Exos for the treatment of many diabetic-related complications.⁷⁰⁹ Figure 20 highlights all the benefits related to the use of exosomes for the treatment of diabetes.

5.2 | Hematology disorders

Hematology disorders are often degenerative by nature and can lead to serious complications and life-threatening conditions. MSC-Exos

have the ability to transport intricate loads and maintain balance within altered homeostasis, hence regulating diseases or malignancies. Recently, the use of MSC-Exos has had significant impacts on many hematological conditions such as graft-versus-host disease (GVHD),⁷⁴⁰ multiple myeloma,⁷⁴¹ acute myeloid leukemia (AML),⁷⁴² chronic myeloid leukemia (CML),⁷⁴³ chronic lymphocytic leukemia (CLL),⁷⁴⁴ lymphomas,⁷⁴⁵ and myelodysplastic syndrome (MDS).⁷⁴⁶ An excellent article by Shen and Chen 2021⁷⁴⁷ focused exclusively on the role of exosomes for hematology disorders and highlighted many studies in Table 24 and a summary of all related benefits found in Figure 21.

5.3 | Musculoskeletal degeneration

There are major obstacles to treating musculoskeletal degeneration injuries such as fracture nonunion, muscle fibrosis, reinnervation, compartment syndrome, and infection and inflammation. In the realm of musculoskeletal regeneration, there has been a recent focus on exosomes. Various cell types release these vesicles, which are crucial for cell communication since they transport functional signaling molecules including proteins and RNAs. A multitude of these cargo molecules may be used for reparative functions in skeletal illnesses such as osteoporosis, osteogenesis imperfecta, sarcopenia, and fracture healing. A research conducted by Youssef El Baradie and colleagues in 2021⁷⁶² titled "Therapeutic application of extracellular vesicles for musculoskeletal repair and regeneration" comprehensively investigated the many uses of exosomes in both repairing and regenerating musculoskeletal tissues.

5.4 | Osteoradionecrosis and radiation therapy

In a study titled: "An indispensable tool: Exosomes are involved in the treatment of radiation-induced damage," Li and colleagues investigated the numerous benefits of exosomes following radiation therapy.⁷⁶³ Approximately 70% of individuals with tumors receive radiation at various time intervals.⁷⁶⁴ Radiotherapy may improve the rate of tumor control and the quality of life for patients. However, normal tissues often experience radiation-induced damage resulting from radiotherapy. Recent research has shown that exosomes have the potential to serve as biomarkers for many illnesses and have a therapeutic role in radiation-induced damage. They further possess the ability to control the inflammatory response, amplify the regenerative impact on injured tissue, and facilitate the restoration of damaged tissues and cells, hence prolonging their lifespan.

Various degrees of radiation damage reduction may be achieved by using MSC-Exos, which include inherent benefits such as minimal immunogenicity, facile culture separation, potent immunosuppressive properties, robust regenerative capabilities, and versatile multi-differentiation capacities.⁷⁶⁵⁻⁷⁷⁰ Recently, there has been significant research on MSCs as they pertain to radiation damage. A growing body of data suggests that the therapeutic benefits of MSC therapy

TABLE 23 MSC-Exos used in the treatment of DM complication.

DM complication	MSC types	Effect/involved non-coding RNA	References
Diabetes wound	Bone marrow MSC	Vascularization	710
	Adipose-derived MSC	Vascularization	711
	Adipose-derived MSC	Vascularization/miR-21-5p	712
	Bone marrow MSC	Vascularization	713
	Bone marrow MSC	Regulate inflammation/lncRNA H19	714
	Adipose-derived MSC	Vascularization	715
	Synovium MSC	Vascularization/miR-126-3p	716
	Urine MSC	Vascularization	717
	Adipose-derived MSC	Vascularization and regulate inflammation	718
	Induced pluripotent stem cell	Vascularization	719
	Menstrual blood-derived MSC	Vascularization and regulate inflammation	720
	Adipose-derived MSC	Vascularization	721
	Adipose-derived MSC	Vascularization	722
	Adipose-derived MSC	Vascularization/mmu_circ_0000250	723
	Bone marrow MSC	Vascularization and regulate inflammation	724
	Bone marrow MSC	Vascularization/miR-221-3p	725
	Umbilical cord MSC	Vascularization	726
Diabetic nephropathy	Adipose-derived MSC	Podocyte repair/miRNA-215-5p	727
	Urine MSC	Podocyte repair	728
	Bone marrow MSC	Anti-fibrosis and promote renal function recovery	572
	Bone marrow MSC	Anti-fibrosis and promote renal function recovery	569
	Adipose-derived MSC	Podocyte repair/miR-486	729
	Urine MSC	Podocyte repair/miR-16-5	730
	Umbilical cord MSC	Reduce kidney inflammation and improve kidney function	731
	Umbilical cord MSC	Reduce kidney inflammation and improve kidney function	732
Diabetic retinopathy	Adipose-derived MSC	Retinal repair/miR-222	708
	Umbilical cord MSC	Retinal repair and regulation of inflammation/miR-126	141
Erectile dysfunction	Adipose-derived MSC	Vascularization and anti-apoptosis	703
	Adipose-derived MSC	Promote angiogenesis and anti-fibrosis/miR-126, miR-130a, miR-132, miR-let7b, miR-let7c	317
	Adipose-derived MSC	Vascularization and anti-inflammatory	733
	Bone marrow MSC	Vascularization and anti-inflammatory/miR-21-5p	734
Cognitive dysfunction	Bone marrow MSC	Nerve repair	735
	Bone marrow MSC	Nerve repair and anti-inflammatory/miR-146a	736
	Bone marrow MSC	Nerve repair	280
Diabetic stroke	Bone marrow MSC	Nerve repair/miR-9	737
	Bone marrow MSC	Nerve repair/miR-145	738
Submandibular gland dysfunction	Bone marrow MSC	Salivary gland function repair	739
Diabetic cardiomyopathy	MSC	Reduce myocardial injury and fibrosis	707

Note: Reprinted with permission from Xiong et al.⁷⁰⁹




FIGURE 20 Exosomes and diabetic complications. The various kinds of diabetic complications, including eye disorders, cardiovascular diseases, nephropathy, neuropathy, delayed wound healing, and endothelial dysfunction, can be ameliorated by exosomes. Neuropathy and nephropathy are the most common diabetic complications. Apoptosis, autophagy, angiogenesis, and fibrogenesis are among the most common molecular mechanisms affected by exosomes in alleviating diabetic complications. Reprinted with permission from Ashrafizadeh et al.⁶⁶⁵

are mostly attributed to the vesicles released via their paracrine pathways, particularly the actions of exosomes.⁷⁷¹ In the article proposed by Li et al.,⁷⁶³ exosomes were shown to play a role following radiation on (1) lungs, (2) skin, (3) intestinal, (4) testicular, (5) bone, and (6) hemopoietic system (Figure 22, Table 25). A recent study titled: "The therapeutic potential of exosomes derived from mesenchymal stem cells in the treatment of osteoradionecrosis." further highlighted the potential for exosomes to improve necrotic tissues caused by osteoradionecrosis.⁷⁷² While this field is still in its infancy and protocols remain to be developed, there is great therapeutic potential for utilizing exosomes to treat radiation damage.

5.5 | Respiratory diseases

Recent interest has focused on the ability for MSC-EVs to treat chronic respiratory diseases. MSC-EVs have the potential to be a revolutionary cell-free treatment for pulmonary fibrosis, COPD, asthma, and pulmonary arterial hypertension, among other chronic respiratory disorders. The two most common ways to administer MSC-EVs are intravenous and intratracheal. Target cells may receive

microRNAs and proteins from MSC-EVs, which further amplifies their therapeutic effects. It is interesting to note that exosomes have been used in several clinical studies for the diagnosis and/or treatment of respiratory disorders and that number is only predicted to rise sharply in the years to come (Figures 23 and 26). Figure 24 summarizes research to date on the topic of exosomes and their associations regarding their regenerative potential.

5.5.1 | Chronic obstructive pulmonary disease

Chronic obstructive pulmonary disease (COPD) is a prevalent chronic lung illness marked by recurrent airway inflammation and permanent, progressive airflow restriction. It significantly impairs the patient's ability to breathe and significantly impacts their daily activities and ability to work.^{806,807} The pathophysiology of COPD is approached from a variety of angles, including protease/antiprotease, oxidative stress, epigenetics, cell aging, apoptosis, chronic inflammation, and linear green body function.⁸⁰⁸ Genetic mutations and environmental variables are involved. The main risk factor for COPD is exposure to cigarette smoke (CS), whether from

TABLE 24 MSC exosomes in hematological diseases.

Disease	MSC sources	Exosomal cargo	Disease model	Biological effect	References
Refractory GVHD	Human BMSCs	NM	Clinical case	Reduced pro-inflammatory cytokine and improved clinical GVHD symptoms	740
aGVHD	Immortalized human embryonic stem cell-derived MSCs	NM	Mouse GVHD model	Enhanced Treg production, alleviated GVHD symptoms, and increased survival by APC	748
aGVHD	Human BMSCs	miR-125a-3p	Mouse GVHD model	Prolonged the survival of mice with aGVHD and reduced the pathologic damage by suppressing the functional differentiation of T cells from a naive to an effect or phenotype	140
aGVHD	Human UC-MSCs	NM	Mouse GVHD model	Lowered the number of CD3 ⁺ CD8 ⁺ T cells; reduced levels of IL-2, TNF- α , and IFN- γ ; increased the ratio of CD3 ⁺ CD4 ⁺ /CD3 ⁺ CD8 ⁺ T cells; and rose serum levels of IL-10	749
GVHD	Human UC-MSCs	TGF- β , IFN- γ , IDO, IL-10	In vitro cell experiment	Promoted PBMCs to differentiate into Tregs via TGF- β and IFN- γ	750
cGVHD	Human BMSCs	NM	Mouse chronic GVHD	Blocked Th17 differentiation and improved the Treg phenotype	751
cGVHD	Human UC-MSCs	NM	Mouse chronic GVHD	Prevented skin fibrosis in the cGVHD mouse model by suppressing the activation of macrophages and B cell immune response	313
MM/lymphoma/leukemia	Young and elderly healthy donor BMSCs	NM	In vitro cell experiment	Antitumor effect existed in the supernatant and not in exosomes; the antiangiogenesis effect depends on the age of donors	752
MM	MM-derived BMSCs	miRNA-15a, IL-2, CCL-2, fibronectin	Mouse MM model	MM patient-derived BMSC exosomes promoted MM tumor growth while normal-derived exosomes inhibited the growth of MM cells	753
MM	Human BMSCs and mouse BMSCs	MCP-1, IP-10, SDF-1	In vitro and in vivo MM model	Favored MM cell proliferation, migration, and survival and induced drug resistance to bortezomib	754
MM	Normal donors and MM BMSCs	NM	In vitro cell experiment	Decreased cells viability, proliferation, migration, and translation initiation with exosomes from normal donor BMSCs, whereas MM MSC exosomes increased	755
MM	Old and young MM-derived BMSCs	miR-340	In vivo model of hypoxic BM in MM	Inhibited MM-induced angiogenesis with exosomes from young BMSCs, and miR-340 inhibited angiogenesis in endothelial cells	756

(Continues)

TABLE 24 (Continued)

Disease	MSC sources	Exosomal cargo	Disease model	Biological effect	References
MM	MM and normal tissue-derived MSCs	LINC00461	In vitro cell experiment	LINC0046 was highly expressed in MSC exosomes and enhanced MM cell proliferation	757
MM	Bortezomib-resistant or bortezomib-sensitive patient MSCs	IncPSMA3, PSMA3-AS1	U266-luc ⁺ xenograft models	Exosomal IncPSMA3-AS1-mediated resistance to proteasome inhibitors by regulating the stability of PSMA3	758
AML	Human BMSCs	S100A4	In vitro cell experiment	Upregulated S100A4 and driven proliferation, invasion, and chemoresistance of leukemia cells	759
AML	Human BMSCs	TGF-β1, miR-155, miR-375	Clinical sample analysis	Released TGF-β1, miR155, and miR375 to mediate extrinsic chemoresistance within the niche in AML	760
AML	HD or newly diagnosed AML patient BMSCs	miR-26a-5p, miR-101-3p, miR-23b-5p, miR-339-3p, miR-425-5p	Clinical sample analysis	Identified candidate miRNAs that provide new insights regarding leukemogenesis and new treatment strategies	742
CML	Human UC-MSCs	NM	In vitro cell experiment	Enhanced the sensitivity of K562 cells to imatinib (IM) via activation of the caspase signaling pathway	761
CML	Human BMSCs	miR-15a	CML xenograft tumor model	Inhibited CML cell proliferation, decreased their sensitivity to IM, and promoted IM resistance	743
CLL	Human BMSCs	NM	In vitro cell experiment	Rescued leukemic cells from spontaneous or drug-induced apoptosis, enhanced their migration, and induced gene expression modifications	744
Hodgkin lymphoma	MSC cell lines	ADAM10	In vitro cell experiment	Induced release of cytokines, like TNFα, sCD30, or CD30 shedding by HL cells	745
MDS	HD and MDS patient BMSCs	miR-10a, miR-15a	In vitro cell experiment	MDS BMSC-derived cargoes overexpressed miR-10a and miR-15a and enhanced cell viability and clonogenic capacity of CD34 ⁺ cells	746

Note: Reprinted with permission from Shen and Chen.⁷⁴⁷

Abbreviations: aGVHD, acute GVHD; HD, health donor; IM, imatinib; IP-10, interferon-inducible protein 10; MCP-1, monocyte chemoattractant protein 1; MDS, myelodysplastic syndrome; miRNAs, microRNAs; NM, not mentioned; SDF-1, stromal cell-derived factor 1; UC-MSC, umbilical cord MSC.




FIGURE 21 Schematic diagram of molecular mechanisms MSC exosomes in hematological diseases. (A) The action of MSC exosomes and subsequent clinical outcomes in GVHD. (B) A brief outline of exosomal cargoes and underlying mechanisms of MSC exosomes in MM. (C) Exosomal loadings and potential effects of MSC exosomes in the diseases of AML, CML, and CLL. AML, acute myeloid leukemia; cGVHD, chronic GVHD; CLL, chronic lymphocytic leukemia; CML, chronic myeloid leukemia; HLA-G, human leukocyte antigen-G; IM, imatinib; MM, multiple myeloma; PIs, proteasome inhibitors. Reprinted with permission from Shen and Chen.⁷⁴⁷

secondhand smoke or active smoking.^{809,810} In some developing nations, exposure to chemicals at work and environmental pollution are major causes of COPD.⁸¹¹ Indoor air pollution exposure is linked to chronic obstructive pulmonary disease and may even have an impact on fetuses. The World Health Organization (WHO) lists COPD as the third most common cause of mortality worldwide. According to statistical forecasts, the number of fatalities from chronic obstructive pulmonary disease will reach 4.4 million by 2040, with low-income and middle-income nations accounting for 90% of these deaths.⁸¹² Issues related to the airway epithelium, such as initial contact and prolonged exposure to CS, can cause epithelial cells to produce pro-inflammatory medium, senescence-associated secretory phenotype (MCP-1, IL-1, IL-6, and IL-8), and damage-associated molecular patterns (HIG box, heat shock

proteins, receptor for advanced glycation end products). The contents that are released into systemic and pulmonary circulation^{813,814} may accelerate the development of COPD by causing damage to the alveoli and lung tissue.⁸¹⁵ The degeneration of lung structure and function cannot be reversed by current COPD therapy approaches; they can only slow down the loss of lung function. Consequently, it is critical to comprehend the molecular process behind the onset and progression of COPD for the purpose of improving the therapeutic treatment plan.

Currently, the primary pharmaceuticals used for COPD treatment are bronchodilators, which are derived from treatments of asthma. However, in COPD patients, the addition of inhalation corticosteroids (ICSs) is associated with a reduced risk of exacerbation and mortality. Nevertheless, the majority of patients do not exhibit




FIGURE 22 Exosomes as an indispensable tool play a role in biomarker and therapy for radiation damage. Exosomes carry many proteins and nucleic acids, which become potential biomarkers for a variety of diseases. And they show broad prospect in the treatment of radiation-induced lung injury, skin injury, intestinal injury, bone injury, testicular injury, and hemopoietic system injury. Reprinted with permission from Li et al.⁷⁶³

TABLE 25 Clinical studies of exosomes affiliated with improvements post-radiation therapy.

Source	miRNAs/proteins	Signaling pathway	Function	References
Lung	miR-28-5p	PI3K/Akt	Regulates MSCs ability	773
Pulmonary epithelial cells	Caspase-3	ROCK I	Reduce lung injury	774
MSCs	KGF	Hippo-Yap	Protecting lung Prevent bone loss	775,776 777
ADSCs			Reduce lung injury Promote wound healing	778 779-783
huc-MSCs	miR-21, miR-221, etc.	β 2/SMAD2 Wnt/ β -catenin	Inhibit pulmonary fibrosis Promote wound healing	784,785 114
Ovarian carcinoma cells	miR-29b	κ B	Promote pulmonary fibrosis	786,787
MenSCs		Let-7/LOX1	Reduce lung injury	788
Human fibroblast	miR-21, miR-126, etc.		Promote angiogenesis	789
BM-SPCs			Alleviate chronic enteritis	790,791
BMSCs	miR-29b	Wnt/ β -catenin	Reduce radiation enteritis Alleviate osteoporosis Weaken radiation-induced bone loss Reverse radiation-induced bone marrow damage	792 793 794 795
Brest milk			Prevent enterocolitis	796-798
Mice macrophages	G-CSF, MIP-2	TLR4	Remain the function of testis	799
Serum		Hippo, PI3K-Akt	HSCs protection	800-802

Note: Reprinted with permission from Li et al.⁷⁶³

Abbreviations: ADSCs, adipose-derived stem cells; BMSCs, bone marrow-derived mesenchymal stem cells; BM-SPCs, bone marrow-derived stromal pro-generator cells; huc-MSCs, human umbilical cord mesenchymal stem cells; MenSCs, menstrual blood-derived endometrial stem cells; MSCs, mesenchymal stem cells.




FIGURE 23 Mechanisms of mesenchymal stem cell-derived extracellular vesicles (MSC-EVs) in treating chronic respiratory diseases. Sources of MSC-EVs: umbilical cord, bone marrow, adipose tissue and induced pluripotent stem cells (iPSCs); isolation methods: ultra-centrifugation (UCF), exosome extraction kits and anion-exchange chromatography; treatment routes: intratracheal delivery and intravenous administration; diseases: chronic obstructive pulmonary disease (COPD), asthma, pulmonary fibrosis, and pulmonary arterial hypertension (PAH). Reprinted with permission from Ma et al.⁸⁰³

sensitivity to steroids, and the administration of high doses amplifies the likelihood of developing pneumonia.⁸¹⁶ Consequently, it is imperative to provide COPD patients with a targeted therapy that addresses their important needs. Additionally, there is a significant need for medications that may prevent the worsening of the illness, as well as the development of pulmonary hypertension and other problems associated with COPD.

In CS-exposed human pulmonary microvascular endothelial cells (HPMECs), Sun et al.⁸¹⁷ observed higher miR-206 expression as well as an increase in caspase-3 and HPMEC apoptosis. MiR-206, Notch3, and VEGFA mRNA levels were negatively correlated. They found that miR-206 targets Notch3 and VEGFA to modulate COPD vascular remodeling.⁸¹⁷ Additionally, anti-miR-27-3p may halt the inflammatory process, reduce BALF inflammatory cytokines, and decrease neutrophil and macrophage infiltration in the lungs. Another study found that overexpressing miR-3202 in COPD may reduce the increase in T lymphocyte IFN- γ and TNF- α caused by CSE while increasing Fas and FasL. High miR-3202 levels decrease T-cell apoptosis and protect human bronchial epithelial cells by targeting FAIM 2.⁸¹⁸ Remarkably, 43 papers were included in a recent systematic review on the use of exosomes for COPD by Gomez et al.⁸¹⁹ The investigations were categorized based on the pathophysiological function of EVs, their origin and cargo, their contribution to COPD exacerbations, and their diagnostic value.⁸¹⁹

5.5.2 | Asthma

Asthma is one of the most common respiratory diseases for which current diagnostic and treatment strategies are still inadequate despite significant advancements.^{820–823} The symptoms associated with asthma such as wheezing, shortness of breath, coughing, chest tightness, and recurring signs of varying levels of intensity and severity, are primarily caused by reversible airway blockage resulting from easily induced bronchospasm and increased mucus production.^{824–827} Alashkar Alhamwe et al.⁸²⁸ recently reviewed the use of exosomes for the treatment and management of asthma as highlighted in Table 27.

5.5.3 | Pulmonary hypertension

Multifactorial reasons contribute to the debilitating nature of pulmonary hypertension (PH).⁸³⁵ Pulmonary vascular resistance is influenced by many etiologies and results in right heart failure. Five types of PH are distinguished based on the hemodynamic characteristics that influence pathogenesis.⁸³⁶ Based on the global prevalence, nearly 70 million individuals are impacted by PH worldwide.^{837,838} Some of the pathologic aspects include dysfunctional endothelial cells (ECs), the proliferation of fibroblasts and smooth muscle cells (SMCs), communication between pericytes, and detrimental changes in ECs and mesenchymal

TABLE 26 Clinical trials in exosomes therapy in respiratory diseases.

No	Title and sponsor	Trial ID	Location	Design	Primary outcome	Recruitment status	Phase
1.	The use of exosomes for the treatment of acute respiratory distress syndrome or novel coronavirus pneumonia caused by COVID-19 (ARDOXSO) Sponsor: AVEM HealthCare	NCT04798716	United States, California	Open label, interventional, mesenchymal stem cell exosomes for the treatment of COVID-19-positive patients with acute respiratory distress syndrome and/or novel coronavirus pneumonia N: 55	Measure and report the number of participants with treatment-related adverse events Tabulate and report the number of IMV days for patients receiving ARDOXSO™ perinatal MSC-derived exosome therapy	Not yet recruiting July 21, 2021	Phase 1 Phase 2
2.	Safety and efficiency of method of exosome inhalation in COVID 19-associated pneumonia (COVID-19EXO2) Sponsor: Olga Tyumina	NCT04602442	Russian, Samara	Randomized, interventional, the extended protocol of evaluation of safety and efficiency of method of exosome inhalation in COVID-19 associated two-sided pneumonia N: 90	Number of participants with nonserious and serious adverse events during trial Number of participants with nonserious and serious adverse during inhalation procedure	Enrolling by invitation October 26, 2020	Phase 2
3.	Evaluation of safety and efficiency of method of exosome inhalation in SARS-CoV-2 associated pneumonia. (COVID-19EXO) Sponsor: State-Financed Health Facility "Samara Regional Medical Center Dinasty"	NCT04491240	Russian, Samara	Randomized, interventional, the protocol of evaluation of safety and efficiency of method of exosome inhalation in SARS-CoV-2 associated two-sided pneumonia N: 30	Number of participants with nonserious and serious adverse events during trial Number of participants with nonserious and serious adverse during inhalation procedure	Completed November 4, 2020	Phase 1 Phase 2
4.	A safety study of IV stem cell-derived extracellular vesicles (UNEX-42) in preterm neonates at high risk for BPD Sponsor: United Therapeutics	NCT03857841	United States, Colorado, Massachusetts, Mississippi, Missouri	Randomized, interventional, a safety study of intravenous infusion of bone marrow mesenchymal stem cell-derived extracellular vesicles (UNEX-42) in preterm neonates at high risk for bronchopulmonary dysplasia N: 3	Number of subjects with treatment-emergent adverse events during the post-treatment phase	Terminated October 12, 2021	Phase 1

TABLE 26 (Continued)

No	Title and sponsor	Trial ID	Location	Design	Primary outcome	Recruitment status	Phase
5.	A clinical study of mesenchymal stem cell exosomes nebulizer for the treatment of ARDS Sponsor: Ruijin Hospital	NCT04602104	China, Shanghai	Randomized, double-blinded, controlled clinical study of allogeneic human mesenchymal stem cell exosomes (hMSC-Exos) nebulized inhalation in the treatment of acute respiratory distress syndrome N: 169	Incidence of adverse reaction Time to clinical improvement 28-day mortality	Recruiting November 2, 2021	Phase 1 Phase 2
6.	A tolerance clinical study on aerosol inhalation of mesenchymal stem cells exosomes in healthy volunteers Sponsor: Ruijin Hospital	NCT04313647	China, Shanghai	Open label, non-randomized, interventional, a tolerance clinical study on aerosol inhalation of mesenchymal stem cells exosomes in healthy volunteers N: 24	Number of participants with adverse reaction (AE) and severe adverse reaction (SAE)	Completed August 4, 2021	Phase 1
7.	Omics sequencing of exosomes in body fluids of patients with acute lung injury Sponsor: Nanfang Hospital of Southern Medical University	NCT05058768	China, Guangdong	Observational, case-control, exosomes in urine, blood, and alveolar lavage fluid from patients with acute respiratory distress syndrome (ADRS) were sequence by omics N: 180	Compare the omics differences of blood samples between the experimental and control groups Compare the omics differences of urine samples between the experimental and control groups	Recruiting September 28, 2021	Case-control
8.	Exosomes derived from placental mesenchymal stem cells as treatment for severe COVID-19: Phase 1 and 2 clinical trials Sponsor: Omid Cell and Tissue center	IRCT20200413047063N2	Tehran, Iran	Participants were randomly divided into two equal groups using a randomized double AB blocking method based on a random number table. Patients allocated randomly to two groups: (1) Intervention 1, Patients will receive Six doses of Exosomes. (2) Control, Patients will receive conventional therapy N: 50	Adverse events assessment	Recruiting July 8, 2021	Phase 1 Phase 2

(Continues)

TABLE 26 (Continued)

No	Title and sponsor	Trial ID	Location	Design	Primary outcome	Recruitment status	Phase
9.	Molecular profiling of exosomes in tumor-draining vein of early-staged lung cancer (ExOnSite- Pro) Sponsor: University Hospital, Limoges	NCT04939324	France, Limoges	Open label, single group assignment, Analyze du Profil moléculaire Des Exosomes de la Veine Pulmonaire Dans le Cancer Bronchique de Stade précoce N: 30	Evaluate size distribution, concentration and molecular profiling of pulmonary vein exosomes at inclusion	Recruiting November 11, 2021	Not applicable
10.	A pilot clinical study on inhalation of mesenchymal stem cells exosomes treating severe novel coronavirus pneumonia Sponsor: Ruijin Hospital	NCT04276987	China, Shanghai	Open label, single group assignment, a pilot clinical study on aerosol inhalation of the exosomes derived from allogenic adipose mesenchymal stem cells in the treatment of severe patients with novel coronavirus pneumonia N: 24	Adverse reaction (AE) and severe adverse reaction (SAE) Time to clinical improvement (TTIC)	Completed September 7, 2020	Phase 1
11.	Extracellular vesicle infusion treatment for COVID-19-associated ARDS (EXIT-COVID-19) Sponsor: Direct Biologics, LLC	NCT04493242	United States, Alabama, California, Pennsylvania, Texas	Randomized, double-blinded, bone marrow mesenchymal stem cell-derived extracellular vesicles infusion treatment for COVID-19 associated acute respiratory distress syndrome (ARDS): a phase II clinical trial N: 120	7 day change in partial pressure of arterial oxygen to fraction of inspired oxygen ratio	Completed December 6, 2021	Phase 2

TABLE 26 (Continued)

No	Title and sponsor	Trial ID	Location	Design	Primary outcome	Recruitment status	Phase
12.	Safety and efficacy of exosomes overexpressing CD24 in two doses for patients with moderate or severe COVID-19 Sponsor: Athens Medical Society	NCT04902183	Greece, Athens, Attica	Randomized, single, a Phase II randomized, single-blind dose study to evaluate the safety and efficacy of exosomes overexpressing CD24 in 10 ⁹ dose versus 10 ¹⁰ dose, for the prevention of clinical deterioration in patients with moderate or severe COVID-19 N: 90	Collection of serious adverse events Proportion of patients related with respiratory rate and SpO ₂ saturation	Recruiting June 15, 2021	Phase 2

Note: Reprinted with permission from Zargar et al.⁸⁰⁴

cells.⁸³⁹ Medically, these disorders result in blood clot formation, swelling, and constriction of the blood vessels. The major elements that influence pulmonary vascular remodeling are the interactions between cellular instigators such as pulmonary ECs and pulmonary artery SMCs, various immune cells, and molecular recruitment mediators such as cytokines and paracrine products.^{840–842} These pathobiological responses worsen the hemodynamic conditions of patients and lead to right heart failure which damages several organs and eventually results in death.^{843,844} An increasing amount of evidence indicates that the development of progressive vascular remodeling in PAH is closely associated with pulmonary vascular inflammation brought on by interactions between various immune cells.^{845,846}

Exosomes play a crucial role in intercellular signaling and communication throughout the physiological cellular process of vascular remodeling. Apart from the fundamental secretory channels that have been theoretically discovered so far, exosomes, and various additional EVs are novel therapeutic ways to treat pulmonary arterial hypertension.⁸⁴⁷ Understanding the traditional channels of secretory proteins that operate on these arteries, as well as the circumstances surrounding the use of nonconventional pathways, are necessary to restore damaged pulmonary vasculature. Exosomes have played a crucial role as immunomodulators in the presentation of antigens and intercellular communication that trigger immune responses. Additionally, new research suggests that exosomes may be important regulators of the inflammatory response, which is one of the processes contributing to the progression of vascular disorders. Despite concerns regarding the excessive proliferative capacity of stem cell treatments, therapies using exosomes provide a robust and promising answer for several illnesses such as PH. One benefit of exosome therapies is their ability to directly be administered at substantial quantities to damaged tissues without risks associated with their native parent MSCs. Table 28 highlights some of these findings.⁸³⁵

5.5.4 | Interstitial lung diseases

Interstitial lung diseases (ILDs) are long-lasting pulmonary illnesses that cannot be reversed and have a high impact on both the health and survival of individuals. The diagnostic methods for ILDs are complex and include several factors. Ongoing research has led to significant advancements in the development of effective treatment approaches including EVs which provides a novel approach in the field of diagnostics (biomarkers) and therapy.⁸⁵⁷ ILDs may occur idiopathically as a result of exposure to biological, chemical, or small particles, or resulting from co-morbidities such as connective tissue or autoimmune illnesses.^{858,859} Chronic, progressive, irreversible fibrosis with substantial morbidity and death is a hallmark of ILDs' natural course.^{860–863} If treating the underlying cause of the illness is crucial for managing nonidiopathic ILDs, conventional treatment methods for ILDs include lung transplantation or antifibrotic medication.^{864–868} Despite the fact that modern therapies significantly reduce morbidity and death,⁸⁶⁹ they do not possess any form of regenerative properties.




FIGURE 24 Overview of biofluid-derived extracellular vesicles in chronic respiratory disease research. Reprinted with permission from Purghe et al.⁸⁰⁵

TABLE 27 Major asthma-related cellular and systemic effects of extracellular vesicles (EVs) released by mesenchymal stem cells (MSCs).

EVs	Source cells/tissue	Recipient	Main effect(s)	Publication
MVs	Equine amniotic MSC	Horse	Reduction in TNF- α secretion and, to a lesser degree, TGF- β and IL-6 from primary alveolar macrophages	829
Exosomes	Human BM-derived MSCs	Human	Upregulation of IL-10 and TGF- β 1 secretion from PBMCs of asthmatics and promotion of proliferation and immunosuppressive capacity of Tregs	830
EVs	Human/mouse BM-derived MSCs	Mouse	Amelioration of <i>Aspergillus</i> extract-induced AAI in sensitized animals	831
miR-1470-containing Exosomes	Human MSCs	Human	Promotion of Tregs differentiation from CD4 ⁺ T cells isolated from PBMCs of acute asthmatics	142
Exosomes	Mouse adipose Tissue-derived MSCs	Mouse	Effective suppression of the maturation of BM-derived DCs as reflected by decreased IL-6 release but augmented IL-10 and TGF- β secretion	832
EVs	Human adipose Tissue-derived MSCs	Mouse	Reduced symptoms and cellular and molecular features of OVA-induced AAI as well as lung TGF- β levels in OVA-sensitized animals	833
Exosomes	Mouse adipose Tissue-derived MSCs	Mouse	Attenuating effect on airway remodeling in a model of OVA-induced AAI could be further augmented by genetic modifications of MSCs	834

Note: Reprinted with permission from Alashkar Alhamwe et al.⁸²⁸

Abbreviations: AAI, allergic airway inflammation; BM, bone marrow; DCs, dendritic cells; IL-10, interleukin-10; MVs, microvesicles; OVA, ovalbumin; PBMCs, peripheral blood mononuclear cells; TGF- β 1, transforming growth factor beta 1; Tregs, regulatory T cells.

TABLE 28 Therapeutic role of stem cells-exosome in pulmonary arterial hypertension.

Exosome source	Cargo molecules	Research object	Role	References
Human AdMSCs	PDGF VEGF FGF	Human Microvascular Endothelial cells	<ul style="list-style-type: none"> Modulating effect on pro-angiogenic and anti-angiogenic factor 	Lopatina et al. ⁸⁴⁸
Human AdMSCs	miR-191	MCT rat model Hypoxic rat model	<ul style="list-style-type: none"> Ameliorating effect on the MCT-induced PAH pathology via BMPR2 degradation 	Zhang et al. ⁸⁴⁹
Human MSCs	miR-196b	Sugen5416/hypoxic Rat model	<ul style="list-style-type: none"> Preventing and reversing effect on pulmonary artery pressure, right ventricular hypertrophy, and pulmonary vascular remodeling Modulating effect on macrophage recruitment to the lung, promote the alternative (M2) macrophage activation pathway, and increase vessel formation 	Klinger et al. ⁸⁵⁰
Human MSCs/mice MSCs	miR-21, 145, 199a	MCT mice model	<ul style="list-style-type: none"> Anti-proliferative, apoptotic, or senescent effects on a variety of hyperproliferative cells Modulating effect on pulmonary vascular remodeling 	Aliotta et al. ⁸⁵¹
Mice mesenchymal stromal cell	miR-204	Hypoxic mice model	<ul style="list-style-type: none"> Suppression effect on hyper-proliferation by STAT3 mediating signaling 	Lee et al. ⁸⁵²
Human placental MSCs	miR-210	Human placenta microvascular endothelial cells	<ul style="list-style-type: none"> Facilitating effect on placental microvascular endothelial cells migration and vascularization 	Komaki et al. ⁸⁵³
Human UC-MSCs	CD63, CD81, TS101, Alix	MCT rat model Hypoxic rat model	<ul style="list-style-type: none"> Attenuating effect on PH pulmonary vascular remodel Reduction effect on excessive proliferation PASMCs by p-GSK3β signaling in PH Protective effect on vascular remodeling and hypoxic PH Inhibitory effect on proliferative STAT3 signaling in PAECs 	Salomon et al. ⁸⁵⁴
Human umbilical cord Wharton's Jelly MSCs	CD34, CD45, CD73, CD90, CD105, HLA-DR	MCT rat model	<ul style="list-style-type: none"> Protective effect on PH vascular remodeling by regulating Wnt5a/BMPR2 signaling 	Zhang et al. ⁸⁵⁵
Human umbilical cord Wharton's Jelly MSCs	ALIX, TSG101, CD81, CD9, CD63, Flotilin-1	Hypoxic mice model	<ul style="list-style-type: none"> Restoring effect on lung architecture Decreasing effect on fibrosis and pulmonary vascular muscularization, ameliorating PH 	Willis et al. ⁸⁵⁶

Note: Reprinted with permission from Oh et al.⁸³⁵

Abbreviations: AdMSC, adipose-derived mesenchymal stem cell; FGF, fibroblast growth factor; MCT, monocrotaline; MSC, mesenchymal stem cell; PAEC, pulmonary arterial endothelial cell; PAH, pulmonary arterial hypertension; PASMC, pulmonary arterial smooth muscle cell; PDGF, platelet-derived growth factor; PH, pulmonary hypertension; STAT3, signal transducer and activator of transcription 3; UC-MSC, umbilical cord-derived mesenchymal stem cell; VEGF, vascular endothelial growth factor.

The involvement of EVs in parenchymal lung damage and interstitial fibrosis is a novel and exciting field of study.⁸⁷⁰⁻⁸⁷² Critical advantages of EVs exist in their mechanistic interventions in the therapy of ILD including reparative, antifibrotic, and immunomodulatory properties.⁸⁷³ Although therapeutic EV investigations are in their early stage of study for the management of ILD, preclinical findings are promising treatment approaches.^{871,874}

According to recent research, BMSC-EVs have been shown to inhibit lung fibrosis and fibroblast proliferation by downregulating the expression of Frizzled Class Receptor 6 (FZD6).⁸⁷⁵ While the precise

mechanism remains unknown, it was also shown that EVs modulate local pro-inflammatory activation of macrophages and monocytes. Additionally, EVs generated from MSC displayed therapeutic effectiveness in lung fibrosis produced by silica and bleomycin.⁸⁷⁶

6 | REGENERATIVE PROCEDURES

Therapeutic exosomes have been utilized successfully for many years owing to their ability to improve many conditions and diseases




FIGURE 25 Changes to circulating extracellular vesicles (EVs) with age. Together, studies suggest that there may be an age-dependent decrease in the concentration of circulating EVs and concomitant changes to EV protein (colored oval within EVs) and EV nucleic acid (curled line within EVs) with age. It remains unknown whether these changes reflect altered secretion, uptake, or rerouting of EVs with age. Autophagy is an important regulator of aging and shares substantial regulatory overlap with EV biogenesis and degradation, but potential interactions between these relationships have not been thoroughly investigated. Reprinted with permission from Lananna et al.⁸⁷⁷

due to their stability and delivery of signaling molecules in a safer manner. This section focuses more specifically on their ability to improve and regenerate damaged tissues in many fields of medicine. These include bone, cartilage, cutaneous wounds, dermatologic applications, dental, vascular, and spinal tissue regeneration.

6.1 | Antiaging

The use of exosomes has improved various cognitive disorders, including Alzheimer's disease, dementia, and multiple sclerosis, with an ability to cross the blood-brain barrier and help repair functions of various cells and their associated tissues. Owing to these abilities, many clinicians have also utilized exosome therapy as a means to prevent aging and especially cognitive function decline.⁸⁷⁷

Recent research indicates that exosomes control both the pathophysiology of age-related illnesses and systemic aging. Exosome studies have undoubtedly focused on their capacity to preserve human health span. It is interesting to note that the biological usefulness of EVs varies depending on the conditions of the cell/tissue source. Therefore, EVs released by aging or diseased cells may have deleterious effects on receiving cells, while EVs released by young or healthy cells may encourage functional improvement.⁸⁷⁷ Therefore, it is important to comprehend the duality of EVs and fully understand that the cell source may shift their positive benefits from an antiaging intervention.⁸⁷⁷

For almost 60 years, there has been credible scientific backing for the once-mythological notion that youthful blood might have antiaging advantages. Even though these early parabiosis trials were fascinating, until much more recently, it was unclear how fresh blood might have rejuvenating benefits.⁸⁷⁸ Therefore, it is becoming increasingly clear how extracellular vesicles and exosomes

support a variety of biological processes, including aging.^{371,878,879}

There is already a significant and rapidly expanding body of evidence that confirms the crucial role of EVs in regulating systemic aging and various age-related problems such as inflammation (the chronic inflammation that occurs with advancing age), cellular senescence, metabolic dysfunction, cardiovascular disease, cancer, and neurodegeneration.^{371,878,880,881} Moreover, it is possible to utilize age-related alterations in EVs as readily available indicators of aging.⁸⁸²

According to recent data, EVs undergo significant alterations as human's age (Figure 25).⁸⁷⁷ A comparable decline in the total amount of EVs circulating between young and elderly participants was observed by several research groups.^{877,883} As a preventive measure to reduce risks, many patients who suffer from disorders that impair cognitive function may receive exosomes by IV, much as in the case of Parkinson's disease, multiple sclerosis, etc. These exosomes must be made from suitable sources and specifically targeted at immune function (immunosomes) in order to enhance functional recovery with the correct signaling molecules.

6.1.1 | Blood-based antiaging therapies

Numerous research studies have demonstrated that blood-based therapy might help with deficiencies associated with aging. Heterochronic parabiosis, for example, has linked the circulatory systems of young and elderly mouse pairs, revealing a number of substances in young blood that prevent aging phenotypes and in old blood that promote aging.^{884–886} By stimulating satellite cells, exposure to a youthful circulatory system revitalizes the ability of muscles to regenerate and returns the proliferative ability of hepatic progenitors.⁸⁸⁷ Additionally, neurogenesis and related learning and

memory abilities are impacted by heterochronic parabiosis, which enhances these capabilities in elderly mice.^{884–886}

Studies have shown that, in addition to parabiosis, animal plasma obtained from exercise provides antiaging cognitive advantages,⁸⁸⁶ suggesting that the plasma EV pool undergoes significant alterations including an increase in EVs when humans exercise.⁸⁸⁸ Together, these results indicate that EVs may play a role in the advantages of blood-based treatments to preserve normal physiological processes as people age. A recent exploratory clinical experiment ([ClinicalTrials.gov](https://clinicaltrials.gov)) confirmed the favorable safety and tolerability of using young human plasma to treat individuals with Alzheimer's disease (NCT02256306).⁸⁸⁹ More thorough research is necessary to determine how much EVs may mediate the advantages and disadvantages of heterochronic parabiosis or other blood-based treatments.

6.1.2 | Effects of EVs on health span and lifespan

EVs may have a significant role in directly regulating a mammal's longevity and health. Research demonstrates that the hypothalamus is the aging control center.^{890–892} Age-related deficits in movement, motor coordination, social interaction, and memory may be lessened by supplementing with EVs generated from hypothalamic neural stem cells.⁸⁹⁰

Another recent study showed that adipose tissue secretes extracellular nicotinamide phosphoribosyltransferase (eNAMPT), a crucial enzyme for NAD⁺ biosynthesis in animals, into the bloodstream in the form of EVs.⁸⁹³ In humans and mice alike, the amount of EV-contained eNAMPT in plasma declines with age and, surprisingly in mice, is a good indicator of life expectancy. Activity levels and sleep fragmentation were two of the age-related deficiencies that were ameliorated by adipose tissue-specific overexpression of NAMPT, which also prevented the age-related drop in eNAMPT levels and increased median lifespan by 13.4%. 20-month-old rats also showed improvements in insulin secretion, glucose tolerance, memory, and ocular function.⁸⁹³

Interestingly, at the conclusion of their study, female mice receiving weekly intraperitoneal injections of EVs carrying eNAMPT which were isolated from young mouse plasma had a longer median and maximum lifespan increase by 10.5% and 16.3%, respectively.⁸⁹⁴ In addition, the administration of EVs resulted in the rejuvenation of both physical appearance and activity levels.⁸⁹⁴ The hypothalamus is one of the main destinations for EVs carrying eNAMPT released from adipose tissue. Simple NAD infusions have become increasingly popular in standard IV clinics since it is now recognized that a systemic decrease in NAD⁺ availability is a major cause of systemic aging.⁸⁹⁵ Exosomes' capacity to have positive effects on NAD levels has several benefits for lifespan.

Investigating how exosomes affect longevity and health also has enormous potential for cancer prevention. This subject has been intensively studied in various review articles.^{371,881,896} However, it should be mentioned that EVs may supply elements that favorably reinforce the relevant disease state in the context of aging,

senescence, cancer, or other pathologies in addition to having the ability to significantly rejuvenate target tissues. This paradox may best be shown by research examining how EVs produced from BMSCs affects the course of multiple myeloma.⁷⁵³ Tumor development in vivo was prevented by EVs obtained from normal BMSCs but encouraged by EVs derived from multiple myeloma BMSCs. This impact may have been mediated by changes in EV composition in MM, such as the enrichment of inflammatory molecules like IL6 and the reduction in tumor-suppressive miR-15a.⁷⁵³

6.1.3 | EVs and "Inflammaging"

EVs are well recognized for their significant roles in both immune system activation and inhibition.^{897,898} It is true that EVs released by cancerous or senescent cells cause telomere degradation and inflammation.⁸⁹⁹ Research has shown that the presence of irradiated cancer cells in cell culture medium resulted in a reduction of both telomere length and telomerase activity in the breast cancer epithelial cells that received them.⁹⁰⁰ Remarkably, the introduction of EVs produced from youthful fibroblasts resulted in a decrease in cellular aging indicators, such as senescence markers, oxidative stress, lipid peroxidation, and inflammatory factors, in several organs of aged animals.⁹⁰¹ A summary of all related benefits to Immunosome infusions is found in [Figure 26](#).

6.2 | Bone regeneration

The skeletal system primarily comprises bone and cartilage, which are composed of many cellular and molecular constituents. The interplay and synchronization of these cellular and molecular components is crucial for preserving homeostasis and effectively rejuvenating bone and cartilage.^{902,903} Health issues related to the musculoskeletal system, mostly caused by osteoporosis, tumors, and fractures, have been more prevalent in recent years.⁹⁰⁴

Many cell types found in bone engage in bidirectional or multidirectional communication with each other ([Figure 27](#)). Exosomes have a significant role in the intricate network of cell communication due to their ability to modulate the immune system. Apart from their functions in osteogenic differentiation and bone formation, bone damage often results in the disturbance of nearby blood vessels, which may hinder the successive phases of bone regeneration.⁹⁰⁵ Through the production of pro-angiogenic substances, MSCs interact with endothelial cells, promoting angiogenesis.⁹⁰⁶

Based on the results of early clinical trials, MSCs have historically been the most promising cell group (from autologous sources) to use in treating disorders affecting the bones and joints. MSCs may be extracted from a variety of tissues, including bone marrow, adipose tissue, and oral tissues including gingiva, periodontal ligament, and tooth pulp.^{907–909} MSCs, in particular, aid in tissue healing because of their capacity to migrate into damaged tissues and owing to their trophic and immunomodulatory




FIGURE 26 Extracellular vesicle theory of aging. Early in life, extracellular vesicles (EVs) serve as beneficial signaling molecules and promote tissue health. As senescent/damaged cells accumulate over the lifespan of an organism, these cells secrete EVs carrying detrimental cargo. These aging-promoting EVs circulate throughout the body, positively reinforcing aging-related tissue deterioration. At the same time, as an organism ages, the number of rejuvenating or supportive EVs decreases. Beneficial effects seen with EV treatments may then reflect a shift of this balance toward rejuvenating EVs and may be one strategy for circumventing age-related deterioration. Proteins are denoted by colored ovals and nucleic acids are denoted by curled lines. Examples of potential effector molecules found to have detrimental effects (top, aging) or beneficial effects (bottom, rejuvenating) are depicted along with supporting citation. Reprinted with permission from Lananna et al.⁸⁷⁷

effects.^{910,911} Additionally, MSCs have the capability for self-renewal and the ability to differentiate into osteoblasts, adipocytes, and chondroblasts in vitro. These characteristics allow them to be increased to therapeutic numbers for many possible applications.^{912,913} However, regulatory concerns have restricted effectiveness of MSCs, even from autologous sources.^{914,915} Over the past decade, clinical researchers have instead concentrated their efforts on the utilization of exosomes produced from MSCs for bone regeneration.^{794,916,917}

The processes of bone regeneration and resorption depend on interactions with the ECM and paracrine signaling between cells. Mechanical stimuli, as well as local and systemic hormonal modulation, impact the network of cells. According to recent research, osteocyte mechanical stimulation results in the Ca^{2+} -dependent release of EVs, including the bone-regulating proteins RANKL, OPG, and sclerostin.⁹¹⁸ This finding suggests that the osteocyte's ability to coordinate tissue-level bone adaptation and control bone metabolism in response to mechanical stimuli is derived by EV release and regulation of RANKL, OPG, and sclerostin secretion.⁹¹⁸

For the purpose of tissue regeneration, MSC-EVs have been studied on a wide variety of cell types found in bone. MSC-EVs have been shown to improve bone fracture,⁹¹⁹⁻⁹²¹ bone defect healing,^{916,922-927} facilitate repair in osteogenesis imperfecta,⁹²⁸ and have been studied in femoral head avascular necrosis brought on by steroids,⁹²⁹ osteochondral abnormalities,^{226,930,931} osteoarthritis (OA),^{210,233,238,240} and periodontal disease.^{932,933} An overview of the current knowledge between interactions among several cell types involved in bone and cartilage regeneration is presented below.

6.2.1 | The functional outcome of MSC-derived sEVs in bone regeneration

Multiple animal models have shown that MSC-EVs stimulates neo-vascularization and bone repair (Table 29). Out of these investigations, the most often used approach for EV therapy is locally injecting native EVs in a liquid mixture.^{210,233,238,240,919,930-932} One study demonstrated positive outcomes on bone regeneration even




FIGURE 27 Multiple cellular and molecular interactions during bone regeneration. Some of the representative cellular interactions and responsible molecules are illustrated in the figure. (1) MSCs—OB/osteocytes: MSCs commit to osteoblasts and terminally differentiate to osteocytes. The secretion of SDF-1 α , TGF- β and BMPs promotes the migration and differentiation of osteoblastic progenitor cells. (2) MSCs—Mo/M ϕ : MSCs regulate migration, proliferation, differentiation, and polarization of monocytes/macrophages via secretion of MCP-1, M-CSF, PGE2 and IDO. (3) MSCs/OB—Mo/OC: MSCs/osteoblasts interact with the osteoclastic lineage via secretion of M-CSF, RANKL and OPG, which regulate the proliferation, differentiation, and activation of osteoclasts. (4) Mo/M ϕ —OC: Macrophages differentially influence the activity of osteoclasts via secretion of pro- or anti-inflammatory cytokines, depending on the M ϕ phenotypes. (5) MSCs—EC: MSC-secreted VEGF and AGN promote angiogenesis via increased proliferation, migration, and tube formation of endothelial cells. AGN, angiostatin; BMPs, bone morphogenetic proteins; EC, endothelial cell; IDO, indoleamine 2,3-dioxygenase; IL-10, interleukin 10; IL-1 β , interleukin 1 β ; M1 M ϕ , pro-inflammatory M ϕ ; M2 M ϕ , anti-inflammatory M ϕ ; MCP-1, monocyte chemoattractant protein-1; M-CSF, macrophage colony-stimulating factor; Mo, monocyte; M ϕ , macrophage; OB, osteoblast; OC, osteoclast; OPG, osteoprotegerin; PGE2, prostaglandin E2; RANK, receptor activator of nuclear factor- κ B; RANKL, RANK ligand; SDF-1 α , stromal cell-derived factor 1 α ; TGF- β , transforming growth factor- β ; TNF α , tumor necrosis factor- α ; VEGF, vascular endothelial growth factor. Reprinted with permission from Wang and Thomsen.⁹³⁴

when a systematic tail vein injection was utilized.⁹²⁸ Commonly, EVs were loaded to various biomaterials including β -TCP,^{922,923} PLA,⁹²⁴ decalcified bovine bone matrix scaffolds,⁹³⁵ or encapsulated in hydrogel^{226,916,920} or collagen sponges.⁹³³

These combined findings show that MSC-EVs directly support bone regeneration by interacting with bone cells, namely those belonging to the MSC-osteoblast-osteocyte lineage. MSC-Exos from diverse origins have been shown to impact osteoblast proliferation, migration, differentiation, and mineralization.^{916,922–926,929,936,937} Additionally, current research has revealed that MSC-EVs up-regulate VEGFA and VEGFR2 expression thereby supporting both osteogenesis and angiogenesis during the process of bone regeneration.^{924,927}

One of the first vital stages of cell-surface interactions and bone-regenerating process associated with implanted prostheses is cell adhesion.^{938,939} MSC attachment to titanium surfaces was enhanced by implant surface-immobilized MSC-EVs, which also had an impact on adherent MSC behavior.⁹⁴⁰ Furthermore, MSC-EVs protected bone-forming osteoblasts from harm during challenged environments such as during implant placement.

Several cargo components and recipient cell molecules have been investigated within EVs to determine the molecular mechanisms during bone regeneration.⁹³⁶ MSC-EVs have been shown to transfer osteogenesis-related microRNAs to activate osteogenic differentiation. VEGF and RUNX2 levels increased, as did osteogenic differentiation.²¹⁰ Another study linked the pro-osteogenic

TABLE 2.9 MSC-derived sEVs in bone and cartilage regeneration.

sEVs and delivery routes	MSC source	Pretreatment of MSCs or sEVs	Experimental model	sEV administration	Functional outcome	References
Native sEVs	Human BM	-	Femoral shaft fracture in C57BL/6 wild-type and CD9 ^{-/-} mice	Local injection into the fractured part at 1 and 8 days after fracture	Promoted fracture healing in wild-type mice and rescued impaired fracture healing in CD9 ^{-/-} mice	919
	Murine BM and human BM	-	Osteogenesis imperfecta animal model in G610 mice	Tail vein injection weekly for 4 weeks	Improved bone growth as indicated by increased bone length in both femora and tibiae	928
	AT	-	Ligature-induced periodontitis model in male albino Wistar rats	Local injection into pockets as an adjunctive treatment	sEV treatment showed promising periodontal regeneration as indicated by formation of highly organized proliferating periodontal ligament tissue and well-formed dense healthy bone in the periodontal ligament space	932
	Human ESCs	-	Critical-sized osteochondral defect on trochlear grooves of the distal femurs in female SD rats	Intra-articular injection weekly for up to 12 weeks	Promoted osteochondral regeneration as indicated by complete restoration of cartilage and subchondral bone	930
	Human ESCs	-	Critical-sized osteochondral defect on trochlear grooves of the distal femurs in female SD rats	Intra-articular injection weekly for up to 12 weeks	Accelerated cartilage repair, increased cellular proliferation and M2 macrophage infiltration, reduced cellular apoptosis and inflammatory cytokine secretion, and enhanced matrix synthesis	931
	Murine BM	-	Collagenase-induced OA model in C57BL/6 mice	Intra-articular injection at Day 7	Protected cartilage and bone from degradation induced by collagenase as indicated by increased cartilage thickness and bone volume	210
	Human ESCs	-	Destabilization of the medial meniscus surgery induced OA model in C57BL/6J mice	Intra-articular injection every 3 days for 4 weeks	Promoted recovery of cartilage destruction as indicated by lower OARSI score and increased expression of Col II	233

TABLE 2.9 (Continued)

sEVs and delivery routes	MSC source	Pretreatment of MSCs or sEVs	Experimental model	sEV administration	Functional outcome	References
	Infrapatellar fat pads from OA patients	-	Induced OA model by surgical destabilization of the medial meniscus in male C57BL/6 mice	Intra-articular injection twice per week for 4 or 6 weeks	Protected articular cartilage from damage and ameliorate gait abnormality as indicated by reduced OARSI score, reversed increase in COL2 expression and downregulated ADAMTS5 and MMP13 expression	240
	Human ESCs	-	Induced TMJ OA model by injection of monosodium iodoacetate in SD rats	Intra-articular injection weekly for up to 8 weeks	Suppressed pain and modulated early gene expression changes in TMJ condylar cartilage tissue, reversed TMJ degeneration as indicated by restored matrix synthesis, alleviated subchondral bone deterioration as indicated by restored bone volume and architecture, suppressed inflammation as indicated by reduced IL-1 β + and iNOS+ cells, promoted proliferation and reduced apoptosis as indicated by increased PCNA+ cells and decreased CCP3+ apoptotic cells	238
Material carrier	Human iPSCs	-	Critical-sized calvarial bone defects in osteopenic animal model using ovariectomized SD rats	Implantation of β -TCP scaffold with lyophilized sEVs in the defect site	Dose-dependently promoted bone regeneration and neovascularization in the defect site as indicated by an increased ratio of BV/TV, BMD, area of new bone formation and mineralization, area and number of vessels, and expression of the osteogenic markers OCN and OPN and the angiogenic marker CD31	922
	Human iPSCs	-	Critical-sized calvarial bone defects in SD rats	Implantation of β -TCP scaffold with lyophilized sEVs in the defect site	Dose-dependently enhanced bone regeneration in the defect site as indicated by an increased ratio of BV/TV, BMD, area of new bone formation and mineralization, and expression of the osteogenic marker OCN	923
	Human gingiva tissue	-	Cortical calvaria bone tissue damage in male Wistar rats	Implantation of 3D printed PLA scaffold enriched with sEVs or both sEVs and MSCs to cover the damaged area	Combination of 3D-PLA with enriched sEVs and MSCs improved bone healing as indicated by higher positive staining of calcium, increased vascularization, higher rate of regeneration and integration level at the damaged site	924

(Continues)

TABLE 2.9 (Continued)

sEVs and delivery routes	MSC source	Pretreatment of MSCs or sEVs	Experimental model	sEV administration	Functional outcome	References
SD rat BM		-	Subcutaneous bone formation model in nude mice	Subcutaneous implantation of sEV functionalized DBM scaffold pre-coating with fibronectin and with or without seeding of osteogenic-induced MSCs	sEV functionalized DBM showed pro-angiogenic activity as indicated by increased CD31+ vessel formation. However, only combination of sEV functionalized DBM and MSCs showed enhanced bone regeneration as indicated by increased staining of new bone, BV and ratio of BV/TV	935
Human AT		-	Calvarial defect in male SD rats	Implantation of sEV-loaded hydrogel in the defect site	Promoted bone regeneration as indicated by micro-CT analysis and histology examination	916
Human umbilical cord		-	Stabilized fracture in femur of male Wistar rats	Injection of sEV-embedded HyStem-HP hydrogel near the fracture site	Enhanced bone healing and angiogenesis at the fracture site as indicated by increased callus formation, BMD, BV, BV/TV, vessel volume, number of CD31+ blood vessels, maximum mechanical load and bending stiffness	920
Human iPSCs		-	Surgically created periodontal intrabony defect in SD rats	Implantation of sEV-loaded collagen sponge in the defect site	Promoted periodontal regeneration with enhanced bone growth, increased functional PDL length and increased cellular infiltration and proliferation at the defect site as indicated by histological examination and micro-CT analysis	933
Human iPSCs		-	Articular full-thickness cartilage defect in New Zealand rabbits	Injection of sEV suspension or sEV-embedded in situ—formed or in vitro—preformed hydrogel glue to the defect site	In situ-formed hydrogel glue containing sEVs had best performance promoting cartilage repair and cartilage-hydrogel integration as indicated by uniform and well-organized articular cartilage structure, distribution of abundant chondrocytes in newly formed tissue, positive staining of Safranin O and Col II	226

TABLE 2.9 (Continued)

sEVs and delivery routes	MSC source	Pretreatment of MSCs or sEVs	Experimental model	sEV administration	Functional outcome	References
Modified sEVs	Rabbit BM	Transfection for expression of mutant HIF-1 α	Steroid-induced avascular necrosis of femoral head in rabbits	Local injection of single-dose sEVs into the femoral head	Promoted bone regeneration and neovascularization in the necrotic region as indicated by massive trabecular tissue generation, increased staining of newborn cartilage and increased density of CD31+ micro vessels	929
	Human umbilical cord	Hypoxic condition	Femoral fracture model in mice	Local injection of single-dose sEVs near the fracture	Promote bone fracture healing primarily via enhanced angiogenesis indicated by increased volume of callus bridging the fracture gap, vessel volume, vessel number, expression of endothelial markers CD31 and endomucin and proliferation marker Ki67	921
Material carrier	Human AT	Osteoinduction for 2 days	Critical-sized calvarial bone defects in male BALB/c mice	Implantation of an sEV-immobilized PLGA/pDA scaffold in the defect site	Increased bone formation and recruitment of SSEA-4+/CD45- MSCs at the defect site as indicated by increased new bone volume, collagen expression and expression of the osteogenic markers RUNX2 and OCN	925
	Human AT	Transfection for overexpression of miR-375	Calvarial defect in male SD rats	Implantation of sEV-loaded hydrogel in the defect site	Enhanced bone regeneration as indicated by increased ratio of BV/TV, BMD, staining of new bone and expression of the osteogenic markers OCN and BMP2	916
	Human gingiva tissue	Coating sEVs with branched PEI by using noncovalent layer-by-layer protocol	Cortical calvaria bone tissue damage in male Wistar rats	Implantation of 3D printed PLA scaffold carried with native sEVs or PEI-coated sEVs to cover the damaged area	Improved bone healing as indicated by new bone formation inside the scaffold structure and presence of blood vessels around the new bone deposition area. PEI-engineered sEVs showed better healing effects	926
	Human PDL tissue	Coating sEVs with branched PEI by using noncovalent layer-by-layer protocol	Cortical calvaria bone tissue damage in male Wistar rats	Implantation of 3D collagen membrane carried hPDLSCs and enriched with native sEVs or PEI-coated sEVs to cover the damaged area	Both native and PEI-engineered sEVs promoted vascularization as indicated by expression of the angiogenesis markers VEGFA and VEGFR2, however, PEI-sEVs improved bone regeneration and integration as indicated by the distribution of OB and OC in the naïve bone, and quantification of bone parameters including BV, BS, BV/TV, and BS/TV	927

Note: Reprinted with permission from Wang and Thomsen.⁹³⁴

Abbreviations: ADAMTS5, a disintegrin-like and metalloproteinase (reprolysin type) with thrombospondin type 1 motif 5; AT, adipose tissue; BM, bone marrow; BMD, bone mineral density; BMP2, bone morphogenetic protein 2; BS, bone surface; BV/TV, bone volume/total bone volume; CCP3, cyclic citrullinated peptide 3; Col II, type II collagen; ESC, embryonic stem cell; HIF-1 α , hypoxia-inducible factor-1 α ; IL-1 β , interleukin-1 β ; iNOS, inducible nitric oxide synthase; iPSC, induced pluripotent stem cell; MMP13, matrix metalloproteinase 13; OA, osteoarthritis; OARS1 score, osteoarthritis research society international score; OB, osteoblasts; OC, osteoclasts; OCN, osteocalcin; OPN, osteopontin; PCNA, proliferating cell nuclear antigen; PDL, periodontal ligament; RUNX2, runt-related transcription factor 2; SD rat, Sprague-Dawley rat; TMJ, temporomandibular joint; VEGFA, vascular endothelial growth factor A; VEGFR2, vascular endothelial growth factor receptor 2.

activities of MSC-EVs to their enrichment of Wnt3a and targeting of the Wnt signaling pathway, one of the most important signaling pathways regulating osteogenic development.⁹⁴¹ Previous authors⁶⁵³ demonstrated that MSC-EVs also increased MSC osteogenic differentiation by activating the PI3K/Akt signaling pathway.^{942,943} A summary of data is found in Table 29. Furthermore, Part III of this publication series further highlights the number of studies on exosomes produced from tooth tissues for bone regeneration.

6.3 | Cartilage/joint/tendon regeneration

In orthopedics, treating cartilage abnormalities such as osteoarthritis (OA) and osteochondral defects (OCD) continues to be very difficult and quite unpredictable. An estimated 654 million persons worldwide suffered from knee OA in 2020 and that number is only expected to increase.⁹⁴⁴

The two primary factors increasing the prevalence and incidence of OA are an aging population and rising obesity rates.⁹⁴⁵ Articular cartilage degrades as a result of OA, a chronic inflammatory condition that causes joint stiffness and discomfort.⁹⁴⁶ OA does not only affect the articular cartilage but also the synovium, subchondral bone, and ligaments around the joints.⁹⁴⁷ The development of OCDs occurs when OA-related cartilage lesions impact the subchondral bone.⁹⁴⁸ Based on the inflammatory component of OA, exosomes have been investigated as a means to repair bone cartilage, joints and tendons.

6.3.1 | In vivo delivery and exosome uptake

Avascular and alymphatic tissue makes up cartilage.⁹⁴⁹ Therefore, exosomes are not appropriate for intravenous infusion for the treatment of cartilage abnormalities, unlike in other tissue injuries. Essentially, since intra-articular injection improves bioavailability and decreases off-target effects (as a whole for any biomaterial), exosome delivery to cartilage is performed locally. To date, all of the included clinical trials have therefore been performed by injecting exosomes directly into the afflicted joint with or without a scaffold.⁹⁵⁰ To regenerate cartilage damage, several studies utilized a single injection, while others performed a series of injections (Table 30). Since no studies examined the therapeutic effects of single versus numerous injections, it is still unknown if multiple injections are more advantageous than single injections for cartilage regeneration. Cosenza and colleagues²¹⁰ as well as Wang et al.⁸⁰ suggested that exosomes administered intraarticularly may be visibly kept within the cartilage tissue even 1 month post-injection. This is most likely because exosomes were directly administered to the target site (cartilage is avascular and alymphatic, therefore exosomes are cleared slower). Exosomes may also remodel resident or injured cells by transferring bioactive molecules that activate regenerative mechanisms, thereby inducing cellular reprogramming.⁹⁵¹⁻⁹⁵³

Ng et al.⁹⁵⁴ conducted a comprehensive review on the use of exosomes in cartilage regeneration, resulting in a summary of 29 studies (Table 30 and illustrated in Figure 28). Direct injection without a carrier seems to be a method most preferred, although a few studies injected exosomes into the damaged cartilage tissues using a scaffold.^{226,239,955} A scaffold has been discussed as being capable of distributing the exosomes more evenly over time and has been shown to prolong the delivery of exosomes over a longer duration. Moreover, it has been suggested that the combination of exosomes with biomaterials might provide a synergistic impact, enhancing the process of cartilage regeneration. According to Liu et al.,^{955,956} in situ hydrogel glue and iPSC-MSC-Exos promoted cartilage regeneration in combination better than either group alone. The group that underwent in situ iPSC-MSC exosome hydrogel tissue patch implantation displayed consistent and well-structured articular cartilage composition, supporting histological results. In vitro research further has shown that exosomes help chondrocytes proliferate and migrate, inhibit apoptosis and the production of pro-inflammatory markers.^{210,931,955-957}

6.3.2 | EVs for tendon repair

EVs and exosomes have also been utilized for tendon repair.⁹⁶⁷ In sports and physical activities, tendon injuries are common where the recovery from injury is slow when compared to other tissues owing to a lack of blood flow. With a high risk of re-injury and the creation of scar tissue, the outcomes of conservative therapies and surgical interventions are not optimal. With 4 million new cases globally each year, tendon and ligament injuries account for 30% of all musculoskeletal consultations and represent a substantial economic and social burden.⁹⁶⁸ Tendon healing has been documented when MSCs have been applied to damaged tendons. According to recent research, MSCs facilitated tendon healing owing to their secretion of EVs/exosomes.⁹⁶⁷

To date, nine investigations utilizing MSC-EVs have demonstrated positive outcomes on tendon and ligament healing. Several benefits have been reported by clinicians using MSC-EVs from cell-based therapies when it comes to treating tendon and ligament disorders including minimal immunogenicity, good stability, no issues with cell viability after implantation, no danger of uncontrolled activity or differentiation of transplanted cells, and no risk of persistence as permanent grafts following treatment discontinuation. Because the cellular components are missing in EVs, platelet-rich plasma/fibrin (PRP/PRF) has been proposed as a delivery vehicle exhibiting benefits in treating tendon and ligament issues.⁹⁶⁹⁻⁹⁷³ Numerous review articles have highlighted the impact of EVs for treating tendons and ligaments, including the use of platelet concentrates.⁹⁶⁹⁻⁹⁷³

The transplantation of rat EVs produced from bone marrow microvascular cells was shown to enhance the repair of the rat Achilles tendon 30 days after damage in a dose-dependent manner, resulting in improvements to the tendon architecture, fiber structure,

TABLE 30 Summary of efficacy and safety findings.

References	Animal models	Method of delivery	Treatment groups	Euthanasia	Efficacy outcome			Safety outcome
					Macroscopic and functional analysis	Imaging	Histological and biochemical analysis	
Zhang et al. ²³⁰	SD rats Surgically induced OCD	Multiple intra-articular injections after surgery and thenceforth weekly	<ul style="list-style-type: none">Exo group: 100 µg/100 µL of ESC-MSC-ExosContralateral control: 100 µL PBSUn-operated control	Week 6 or 12 post-surgery	<ul style="list-style-type: none">Exo group demonstrated almost complete neotissue filling with good surface regularity and complete integration of neotissue with surrounding cartilage by 12 weeks	Not reported	<ul style="list-style-type: none">Exo group exhibited almost complete regeneration and bonding of cartilage and underlying subchondral bone after 12 weeksDistinctly higher modified O'Driscoll histological scores in the Exo groupExo group showed hyaline cartilage formation with high amount of GAG, Col II, and low amount of Col I	No detrimental responses were observed in all animals
Cosenza et al. ²¹⁰	C57BL/6 mice Collagenase-induced OA	Single intra-articular injection at day 7 after OA induction	<ul style="list-style-type: none">Cell group: 2.5 × 10⁵ BMSCs/5 µLMP group: 500 ng/5 µL of BMSC-microvesiclesExo group: 250 ng/5 µL of BMSC-ExosOA groupHealthy control	Day 42 post-collagenase induction	Not reported	<ul style="list-style-type: none">CLSM showed structural improvement in articular cartilage which comparable with healthy control group in all treatment groupsµCT showed higher bone volume, lower bone degradation, lower osteophyte formation and lower calcification of menisci and ligaments in all treatment groups compared with the OA group	<ul style="list-style-type: none">Exo group showed the greatest improvement with the lowest OA scores	Not reported

Liu et al. ²²⁶	New Zealand rabbits Surgically created OCD	Scaffold implantation or single intra-articular injection immediately after surgery	<ul style="list-style-type: none">EHG group: 20 µL in situ formed EHG tissue patch containing 1 × 10¹¹/mL iPSC-MSC-ExosHG group: 20 µL in situ formed HG tissue patchPre-EHG group: 20 µL in vitro preformed EHG containing 1 × 10¹¹/mL iPSC-MSCs-ExosInj-Exo group: 20 µL of 1 × 10¹¹/mL iPSC-MSCs-Exos suspensionOA group: Saline rinsing	Week 12 post-surgery	<ul style="list-style-type: none">EHG group showed the best repair with smooth surface with white regenerated tissue fully filled the defects and integrated with surrounding cartilage	<ul style="list-style-type: none">OCT displayed uniform and well-organized articular cartilage structure in EHG group	<ul style="list-style-type: none">EHG group has the highest ICRS score and the neotissue was almost entirely hyaline cartilage (strong safranin O and Col II staining, weak Col I staining)	Not reported
---------------------------	--	---	--	----------------------	---	---	---	--------------

(Continues)

TABLE 30 (Continued)

Efficacy outcome								
References	Animal models	Method of delivery	Treatment groups	Euthanasia	Macroscopic and functional analysis	Imaging	Histological and biochemical analysis	Safety outcome
Tao et al. ²³⁰	SD rats Surgically induced OA	Multiple intra-articular injections on the first day of week 5–8 after surgery	<ul style="list-style-type: none">Exo group: 100 µL of 10¹¹ SM-MSC-Exos particles/mLExo-miR-140-5p group: 100 µL of 10¹¹ miR-140-5p overexpressed SM-MSC-Exos particles/mLOA group: SalineHealthy control: Saline	Week 12 post-surgery	Not reported	Not reported	<ul style="list-style-type: none">Exo-miR-140-5p group showed the lowest OARSI assessment scoresSignificant increased chondrocyte count, higher Col II and aggrecan expression, and lower Col I in the Exo-miR-140-5p group compared with the Exo and OA groups	No adverse events occurred
Wang et al. ²³³	C57BL/6J mice DMM induced OA	Intra-articular injections at week 4 after surgery (once for Cell and Cell-OA group; multiple injections every 3 days for 4 weeks for Exo and Exo-OA group)	<ul style="list-style-type: none">Cell group: 5 µL of 1 × 10⁶ ESC-MSC suspensionOA group: 5 µL of PBSSham controlExo group: 5 µL ESC-MSC-ExosOA group: 5 µL of PBSSham control	Week 8 post-surgery	Not reported	Not reported	<ul style="list-style-type: none">Exo group exhibited similar regenerative effect as Cell groupExo group revealed milder OA pathology compared with the OA group which was concomitant with lower OARSI scores and stronger Col II staining and weaker ADAMT5 and aggrecan neopeptide staining	Not reported
Zhu et al. ²²⁹	C57B/L10 mice Collagenase-induced OA	Multiple intra-articular injections on day 7, 14 and 21 after collagenase administration	<ul style="list-style-type: none">iPSC-MSC-Exo group: 8 µL of 1 × 10¹⁰ particles/mLiPSC-MSC-ExosSM-MSC-Exo group: 8 µL of 1 × 10¹⁰ particles/mLSM-MSC-ExosOA group: 8 µL of PBSNormal control: 8 µL of PBS	Day 28 post-collagenase induction	<ul style="list-style-type: none">No significant differences in ICRS macroscopic analysis scores among the normal, iPSC-MSC-Exo and SM-MSC-Exo groups, but significantly higher than the OA group	Not reported	<ul style="list-style-type: none">Neotissue of iPSC-MSC-Exo group presented smooth cartilage, regular cellular organization, and normal proteoglycan content that similar to control group: SM-MSC-Exo exhibited moderate MSC-Exo scores between the iPSC-MSC-Exo and normal group, but notable lower than the SM-MSC-Exo and OA groupsMore intense Col II staining in the iPSC-MSC-Exo group compared with the SM-MSC-Exo group	Not reported

TABLE 30 (Continued)

Efficacy outcome								
References	Animal models	Method of delivery	Treatment groups	Euthanasia	Macroscopic and functional analysis	Imaging	Histological and biochemical analysis	Safety outcome
Mao et al. ¹⁴⁹	C57B/L10 mice Collagenase-induced OA	Multiple intra-articular injections on day 7, 14 and 21 after collagenase administration	<ul style="list-style-type: none">Exo group: 15 µL of 500µg/mL BMSC-ExosExo-miR-92a-3p group: 15 µL of 500 µg/mL miR-92a-3p overexpressed BMSC-ExosOA group: 15µL of PBSHealthy control: 15 µL of PBS	Day 28 after collagenase induction	Not reported	Not reported	<ul style="list-style-type: none">Exo-miR-92a-3p group demonstrated significant reduced severity of cartilage matrix loss with higher Col II and aggrecan, and lower Wnt5a and MMP13 expressions in both gene and protein levels compared to the Exo and OA groups	Not reported
Wang et al. ²⁵⁴	SD rats Surgically induced OA	Intra-articular injection	<ul style="list-style-type: none">Exo group: 100 µL of 1 × 10¹¹ particles/mL MSC-ExosTGF-β1-Exo group: 100 µL of 1 × 10¹¹ particles/mL TGF-β1-MSC-ExosTGF-β1-NC-Exo group: 100µL of 1 × 10¹¹ particles/mL TGF-β1-MSC-NC-ExosTGF-β1-miR135b inhibitor-Exo group: 100 µL of 1 × 10¹¹ particles/mL TGF-β1-MSC-miR135b inhibitor-Exos	Week 12 post-surgery	Not reported	Not reported	<ul style="list-style-type: none">Significant lower OARSI scores in the TGF-β1-Exo group than the Exo group, with high OARSI scores in the TGF-β1-miR135b inhibitor-Exo groupTGF-β1-Exo group showed elevated number of chondrocytes, while dropped in chondrocyte number was recorded in the Exo and TGF-β1-miR135b inhibitor-Exo groups	Not reported
Zhang et al. ²⁵¹	SD rats Surgically induced OCD	Multiple intra-articular injections immediately after surgery on weekly basis	<ul style="list-style-type: none">Exo group: 100µg/100 µL ESC-MSC-ExosContralateral control: 100 µL PBS	Week 2, 6, or 12	Not reported	Not reported	<ul style="list-style-type: none">Exo group demonstrated predominately hyaline cartilage regeneration (mostly Col II and very low Col I expression) and complete integration of neotissue with adjacent native cartilage with a smooth surface regularity and complete regeneration of subchondral bone at Week 12Wakitani scores of Exo group decreased from week 2 to 12 and the results were significantly lower than the contralateral controlEnhanced proliferation, attenuated apoptosis, increased M2 macrophages and reduced M1 macrophages in both cartilage andsynovium tissues, and decreased in M1-associated cytokines, that is, IL-1β and TNF-α, in synovial fluid of the Exo group	Not reported

(Continues)

(Continues)

TABLE 30 (Continued)

References	Animal models	Method of delivery	Treatment groups	Euthanasia	Efficacy outcome			Safety outcome
					Macroscopic and functional analysis	Imaging	Histological and biochemical analysis	
Chen et al. ²³⁹	New Zealand White rabbits Surgically induced OCD	3D printed scaffold implantation	<ul style="list-style-type: none">ECM/GelMA group: ECM/GelMA scaffoldGelMA group: GelMA scaffoldECM/GelMA/Exo group: 3D printed ECM/GelMA/BMSC-Exo scaffoldOA group: Untreated	Week 6 or 12 post-surgery	ECM/GelMA/Exo group exhibited smooth and intact tissues and gave the highest ICRS macroscopic analysis scores	<ul style="list-style-type: none">MRI scanning demonstrated smooth neo-cartilage and great defect filling in both ECM/GelMA and ECM/GelMA/Exo groupsµCT displayed increased ratio of bone volume to tissue volume, trabecular thickness, and ossified tissues in the subchondral bone of both ECM/GelMA and ECM/GelMA/Exo groups	<ul style="list-style-type: none">ECM/GelMA and ECM/GelMA/Exo groups showed hyaline-like cartilage regeneration in the defect sitesECM/GelMA/Exo group has higher ICRS visual histological scores at week 12Increased Col II and decreased of MMP13 expression in the synovial membrane of ECM/GelMA and ECM/GelMA/Exo groupsECM/GelMA/Exo group expressed the lowest MDA levels	No apparent pathological effects in myocardium, liver, and kidney 1 to 2 weeks after transplantation
Liu et al. ²⁵⁶	New Zealand White rabbits Surgically induced OA	Multiple intra-articular injections once a week	<ul style="list-style-type: none">Exo group: 100 µg/mL PRP-ExosPRP-As group: 100 µg/mL activated PRPOA group: Normal salineControl: Normal saline	Week 6 after surgery	Not reported	Not reported	<ul style="list-style-type: none">Exo group exhibited more regular arrangement of chondrocytes, clearer tidal line, reduced hyperplasia on articular cartilage surface and lower OARSI scores than the PRP-As and OA groupsExo group showed increased expression of Col II and RUNX 2	No adverse events occurred
Wu et al. ²⁴⁰	C57BL/6 mice DMM induced OA	Multiple intra-articular injections twice a week, starting at week 4 after surgery	<ul style="list-style-type: none">Exo group: 10 µL of 10¹⁰ particles/mL IPFP-MSC-ExosOA group: 10 µL PBSSham control: 10 µL PBS	Week 8 post-surgery	Improved gait pattern (CatWalk gait analysis) after 6 weeks of exosome treatment	<ul style="list-style-type: none">Obvious green, fluorescent dots (DIO-labeled IPFP-MSC-Exos) can be found at the defect sites	<ul style="list-style-type: none">Exo group showed integration of cartilage with smooth surface and lower OARSI scores confirming cartilage lesion was healedExo group showed higher expression of Col II as well as lower expression of ADAMTS5 and MMP13	Not reported
Zhao et al. ²⁵⁸		3 weeks of 10 µL antagonist pre-injection (once a week), starting at week 1 after surgery, followed by injection of Exos and antagonist twice a week for 4 weeks	<ul style="list-style-type: none">PBS + antagonist-NC group: 10 µL PBS + antagonist-NCExo + antagonist-NC group: 10 µL 10¹⁰ particles/mL IPFP-MSC-Exos + antagonist-NCExo + antagonist-100-5p group: 10 µL 10¹⁰ particles/mL IPFP-MSC-Exos + antagonist-100-5p	Week 8 post-surgery	Not reported	Not reported	<ul style="list-style-type: none">Exo + antagonist-100-5p group reversed the results of the Exo group	Not reported

TABLE 30 (Continued)

Efficacy outcome								
	Macroscopic and functional analysis			Imaging		Histological and biochemical analysis	Safety outcome	
Zhang et al. ²³⁸	SD rats MIA-induced TMJ-OA	Method of delivery	Treatment groups	Euthanasia	Macroscopic and functional analysis	Imaging	Histological and biochemical analysis	Safety outcome
								</

(Continues)

TABLE 30 (Continued)

References	Animal models	Method of delivery	Treatment groups	Euthanasia	Efficacy outcome			Safety outcome
					Macroscopic and functional analysis	Imaging	Histological and biochemical analysis	
Zheng et al. ⁹⁵⁹	C57BL/6 mice Surgically induced OA	Multiple intra-articular injections once per week, starting 10 days after surgery	<ul style="list-style-type: none"> Exo group: 200 µg of primary Chondrocyte-Exos OA group: 20 µL saline Sham control: Untreated 	Week 4, 6, or 8 post-surgery	Not reported	µCT demonstrated lower subchondral bone mineral density and smaller osteophyte formation at the joint margins in the Exo group	<ul style="list-style-type: none"> Both femoral and tibia cartilage of the Exo group showed nearly complete preservation No obvious synovitis appearance in all groups Exo group expressed marked repression in MMP13 staining and elevated Col II staining in joint samples Significant lower OARSI scores in the Exo group Exo group exhibited higher level of M2 macrophages infiltration in the synovium and cartilage tissues Exo-miR-136-5p group showed lesser loss of cartilage matrix followed by the Exo and OA groups Exo-miR-136-5p group revealed higher level of Col II and aggrecan gene and protein expressions, decreased in ELF3 and MMP13 gene and protein expression 	Not reported
Chen et al. ²⁴⁷	C57BL/6 mice Mechanical load induced OA	Single intra-articular injection immediately after mechanical induction	<ul style="list-style-type: none"> Exo group: 100 µL of 10¹¹ particles/mL BMSC-Exos Exo-miR-136-5p group: 100 µL of 10¹¹ particles/mL miR-136-5p overexpressed BMSC-Exos OA group: Untreated Normal control: Untreated 	1 h after injury	Not reported	Not reported	<ul style="list-style-type: none"> Exo-miR-136-5p group showed lesser loss of cartilage matrix followed by the Exo and OA groups Exo-miR-136-5p group revealed higher level of Col II and aggrecan gene and protein expressions, decreased in ELF3 and MMP13 gene and protein expression 	Not reported

TABLE 30 (Continued)

Efficacy outcome								
References	Animal models	Method of delivery	Treatment groups	Euthanasia	Macroscopic and functional analysis	Imaging	Histological and biochemical analysis	Safety outcome
He et al. ²³⁴	SD rats MIA-induced OA	Multiple intra-articular injections once a week, starting 1 week after OA induction	<ul style="list-style-type: none">• Exo group: 40 µg/100 µL BMSC-Exos• OA group: 100 µL of normal saline• Sham control: 100 µL of normal saline	Week 6 post-treatment	<ul style="list-style-type: none">• Joint injuries were alleviated in the Exo group Thermal PWL and mechanical PWT improved at Weeks 2, 4 and 6 in the Exo group	In vivo imaging showed accumulation of PKH26 labeled BMSC-Exos in the joint cavity	<ul style="list-style-type: none">• Articular cartilage of the Exo group regenerated, with small number of defects and fractures on cartilage surface• Exo group has significantly reduced OARSJ scores• Exo group demonstrated notably increased in Col 2 protein expression level in chondrocytes and ECM, and decreased of MMP13 and Col 1 protein levels in cartilage tissue• Exo group showed reduced expression of CGRP and iNOS in DRG• Lower inflammatory cytokines (IL-1β, IL-6 and TNF-α) and higher anti-inflammatory cytokine (IL-10) in serum of Exo group	Not reported
Jin et al. ²⁶⁰	SD rats Surgically induced OA	Intra-articular injection, starting 2 weeks after OA induction	<ul style="list-style-type: none">• Exo group: BMSC-Exos• OA group: Normal saline	Week 7 post-treatment	Not reported	Not reported	<ul style="list-style-type: none">• Lower observation and Mankin scores in the cartilage of the Exo group• Notably decreased in inflammatory factors, AKP content and oxidative stress injury indicators, but increased in SOD in synovial fluid of the Exo group• Cartilage tissue of Exo group showed significant reduced MMP13, OCN and COMP gene and protein expressions	Not reported

(Continues)

TABLE 30 (Continued)

References	Animal models	Method of delivery	Treatment groups	Euthanasia	Efficacy outcome			
					Macroscopic and functional analysis	Imaging	Histological and biochemical analysis	
Jin et al. ²³⁵	Wistar rats Surgically induced OA	1-week intra-articular injection of treatments after surgery	<ul style="list-style-type: none">Exo-miR-26a-5p group: 250 ng/5 μL of miR-26a-5p overexpressed BMSC-ExosExo-miR-NC group: 250 ng/5 μL of miR-NC BMSC-ExosOA group: UntreatedSham control: UntreatedHealthy control: Untreated	Week 8 post-surgery	Not reported	Not reported	<ul style="list-style-type: none">Exo-miR-26a-5p group has less pathological changes with the concomitant of reduced synovial tissue proliferation and suppressed inflammationExo-miR-26a-5p group has lower MMP3 and MMP13 expression and higher apoptotic index in synovial cellsNotably increased of miR-26a-5p expression and declined of PTGS2 expression in synovial tissue of the Exo-miR-26a-5p groupExo-miR-26a-5p group showed reduced serum IL-1β levels	Not reported
Liang et al. ²⁶¹	SD rats DMM induced OA	Multiple intra-articular injections post-injury, once per week	<ul style="list-style-type: none">Exo group: 100 μg of Exos particles in 100 μL PBSExo-miR140 group: 100 μg of Exo-miR140 particles in 100 μL PBSCAP-Exo-miR140 group: 100 μg of CAP-Exo-miR140 particles in 100 μL PBSOA group: UntreatedSham control: Untreated	Week 8 post-surgery	Not reported	Fluorescence microscopy showed that CAP-Exo-miR-140 mainly stayed in the articular cavity, while Exo-miR-140 distributed to other body parts and enriched in kidney	<ul style="list-style-type: none">CAP-Exo-miR140 group displayed smooth and flat cartilage surface, small joint space, proper cell alignment, normal subchondral bone and dense proteoglycan which was almost identical to the sham control and comparable OARSJ scores with sham controlCAP-Exo-miR140 group suppressed the MMP13 and Adamts5 protein levels in cartilage tissueUpregulation of miR-140 and downregulation of MMP13 gene expression in cartilage tissue in the CAP-Exo-miR140 group	Nontoxic to major organs (heart, liver, kidney, lung, and spleen)

TABLE 30 (Continued)

References	Animal models	Method of delivery	Treatment groups	Euthanasia	Efficacy outcome			Safety outcome
					Macroscopic and functional analysis	Imaging	Histological and biochemical analysis	
Liu et al. ⁹⁵⁵	SD rats Surgically induced OCD	Intra-articular implantation	Exo group: Col-Tgel hydrogel with BMSC-Exos	Weeks 4, 6, and 8 post-surgery	Better cartilage regeneration in the KGN-Exo group as indicated by smoother articular surface and better integration of newly formed cartilage with adjacent host cartilage	Not reported	KGN-Exo group revealed better cartilage reconstruction with hyaline cartilage predominantly, corresponding to notable high ICRS visual histological scores	Not reported
			KGN-Exo group: Col-Tgel hydrogel with KGN-BMSC-Exos		Exo group exhibited higher ICRS macroscopic scores which were comparable with the normal group at 8 weeks			
Qiu et al. ²⁴⁸	Mice Surgically induced OA	Not mentioned	Exo-Cur group: Curcumin pretreated BMSC-Exos	Not reported	Not reported	Not reported	Exo-Cur group showed higher gene expression of miR-124 and miR-143 as well as lower protein expression of ROCK1, NF-κB and TLR9 that were similar to the sham control group	Not reported
			Exo group: BMSC-Exos					
Wong et al. ⁹⁶²	New Zealand white rabbits Surgically induced OCD	Multiple intra-articular injections at day 0, day 7 and day 14 post-surgery	Exo + HA group: 1 mL of 3% (w/v) hyaluronic acid with 200 μg ESC-MSC-Exos	Week 6 or 12 post-treatment	Significant improvement in ICRS scores of Exo + HA group at week 6 and 12, associated with marked improvements of neotissue integration at the border zone	Not reported	Exo + HA group showed improvements in cartilage regeneration over time as confirmed by complete defect coverage by neotissue characterized by the presence of hyaline cartilage, normal cellularity and chondrocytic-like cells, high GAG and Col II deposition, and lower Col I deposition at week 12	No adverse events
			HA group: 1 mL of 3% (w/v) hyaluronic acid		Exo + HA group displayed improved mean Young's moduli and stiffness of the repaired cartilage that approximated the normal tissue			

TABLE 30 (Continued)

References	Animal models	Method of delivery	Treatment groups	Euthanasia	Efficacy outcome		
					Macroscopic and functional analysis	Imaging	Histological and biochemical analysis
Yan and Wu ⁹⁶³	New Zealand rabbits Surgically induced cartilage defect	Multiple intra-articular injections on a weekly interval	2D-Exo group: 500 µL of 1 × 10 ¹⁰ particles/mL 2D cultured UC-MSC-Exos 3D-Exo group: 500 µL of 1 × 10 ¹⁰ particles/mL 3D cultured UC-MSC-Exos OA group: 500 µL PBS	Week 4 post-treatment	Significant improvement in ICRS macroscopic scores in the 3D-Exo group compared to the 2D-Exo and OA groups 3D-Exo group demonstrated more neotissue formation with smoother surface and better integration with the native hyaline cartilage at the surrounding	Not reported	3D-Exo group displayed partly hyaline cartilage and the defects showed greater surface regularity and better thickness of cartilage than the 2D-Exo group 3D-Exo group exhibited lower Wakitani score than the 2D-Exo and OA groups
Zavatti et al. ⁹⁶⁴	CD rats MIA-induced OA	Intra-articular injection 3 weeks after OA induction (once for Cell and OA groups; twice for Exo group with 10 days interval)	Cell group: 50 µL of 5 × 10 ⁵ AFSCs Exo group: 50 µL of 100 µg AFSC-Exos OA group: 50 µL of glucose/PBS Contralateral control	Week 3 Post-treatment	Cell and Exo groups have higher pain tolerance, and the results were comparable to the normal control group	Not reported	Exo group showed better cartilage regeneration as indicated by complete neotissue formation with good surface regularity compared to the cell-treated defects which exhibited few fissures on the cartilage surface Uniformed GAG distribution was demonstrated in the Exo group Both Exo and cell groups exhibited improved OARSI scores Cell group displayed more intense Col II staining, whereas Exo group showed more regular distribution of Col II

No adverse inflammatory responses were observed

No adverse inflammatory responses were observed

No adverse inflammatory responses were observed

No adverse inflammatory responses were observed

TABLE 30 (Continued)

References	Animal models	Efficacy outcome					Safety outcome	
		Method of delivery	Treatment groups	Euthanasia	Macroscopic and functional analysis	Imaging		Histological and biochemical analysis
Zhang et al. ²⁰⁹	SD rats Surgically induced OA	Intra-articular injection every 3 days for 4 weeks, starting at Week 4 Post-operation	<ul style="list-style-type: none">• Exo group: 10 μL 1010 particles/mL BMSC-Exos• OA group: 10 μL PBS• Sham control	Week 4 post-treatment	Not reported	<ul style="list-style-type: none">• μCT showed less cartilage degradation, near-normal chondrocyte morphology and distribution, and less osteophyte formation around the joint treated with Exo• Dil-labeled Exo were observed in the knee joint and taken up by the synovial cells	<ul style="list-style-type: none">• Lower OARSI scores in the Exo group compared with the OA group• Exo group demonstrated upregulated chondrogenic gene expressions and downregulated hypertrophic gene expressions• Exo group has decreased synovial hyperplasia and cell filtration• Exo group has less M1-positive cells and more M2-positive cells• Exo group exhibited reduced pro-inflammatory cytokines (IL-1β, TNF-A) and increased anti-inflammatory cytokine (IL-10) in synovial fluidExo group showed significant reduction in synovitis scores	Not reported

Zhou et al. ²³⁶	C57BL/6J mice Collagenase VII induced OA	Multiple intra-articular injections at Days 7, 14, and 21 after collagenase induction	<ul style="list-style-type: none">• Exo group: BMSC-Exos• pExo group: polydactyl BMSC-Exos• OA group: Saline• Normal control	Day 28 post-collagenase VII injection	Not reported	Not reported	<ul style="list-style-type: none">• pExo group alleviated cartilage damage evidenced by the significant lower OARSI scores than the OA and Exo groups	Not reported
----------------------------	--	---	---	---------------------------------------	--------------	--------------	---	--------------

(Continues)

TABLE 30 (Continued)

Efficacy outcome								
References	Animal models	Method of delivery	Treatment groups	Euthanasia	Macroscopic and functional analysis	Imaging	Histological and biochemical analysis	Safety outcome
Wang et al. ⁹⁶⁵	C57BL/6J mice Surgically induced OA	Single intra-articular injection at Week 4 post-operation	<ul style="list-style-type: none">Sham-Exo group: 200µg sham-ExosOA-Exo group: 200µg OA-ExosATF4-OA-Exo group: 200µg ATF4-OA-ExosOA groupSham control	Week 8 post-surgery	Not reported	µCT showed reduced osteophyte formation in Sham-Exo and ATF4-OA-Exo groups, while OA-Exo group displayed enlarged osteophytes	<ul style="list-style-type: none">Sham-Exo and ATF4-OA-Exo groups alleviated pathological injury of articular tissues observed in the OA group, whereby the ATF4-OA-Exo group exerted greater therapeutic effect and lesser proteoglycan lossATF4-OA-Exo group exhibited lower Mankin scores than the Sham-Exo group, while OA-Exo group aggravated the cartilage damage and exhibited the highest Mankin scores among all groupsATF4-OA-Exo group gave stronger effect in upregulating Col II levels and downregulating MMP13 levelsATF4-OA-Exo group was more potent in decreasing inflammatory cytokine levels in the cartilage ATF4-OA-Exo group partially restored the impeded autophagy of the OA cartilage	Not reported
Wang et al. ²³²	BALB/C mice 4°C water stimulated OA	Multiple intra-articular injection once a day, starting from day 20 after OA induction	<ul style="list-style-type: none">Exo group: 30µL of 10¹¹ particles/mL SM-MSC-ExosExo-155-5p group: 30µL of 10¹¹ particles/mL miR-155-5p overexpressed SM-MSC-ExosOA group: normal salineNormal control	2 weeks post-treatment	Not reported	Not reported	<ul style="list-style-type: none">Exo-155-5p group revealed lower OARSI scores and higher chondrocyte number in femoral condyleExo and Exo-155-5p groups reversed the increased caspase-3 and decreased Col II expressions in OA femoral condyle sections, but Exo-155-5p was more effective	Not reported

TABLE 30 (Continued)

References	Animal models	Method of delivery	Treatment groups	Euthanasia	Efficacy outcome			Safety outcome
					Macroscopic and functional analysis	Imaging	Histological and biochemical analysis	
Yan et al. ⁹⁵⁴	SD rats Surgically induced artilage defects	Multiple intra-articular injections on a weekly interval	<ul style="list-style-type: none">• S-Exo group: 100 µL of 1 mg/mL rotary cell culture system cultured UC-MSC-Exos• si-Exo group: 100 µL of 1 mg/mL rotary cell culture system cultured siRNA H19 transfected UC-MSC-Exos• OA group: 100 µL PBS	Week 4 or 8	<ul style="list-style-type: none">• S-Exo group revealed reduced pain with higher LWT at Week 3• Defects covered with neotissue at Week 4 and were filled with uniform tissue and obscured boundaries at Week 8 in the S-Exo group• S-Exo group showed the highest ICRS scores	<ul style="list-style-type: none">• Defects of S-Exo group displayed similar intensity to the adjacent cartilage through MRI at Week 8• S-Exo group exhibited the lowest T2 values at Weeks 4 and 8	<ul style="list-style-type: none">• S-Exo group has the best surface regularity, highest glycosaminoglycan, most orderly tissues and best subchondral bone repair among the groups• S-Exo group has significant lower Wakitani scores compared to the other groups• S-Exo group displayed the highest matrix synthesis which consists mainly of Col II	Not reported

Note: Reprinted with permission from Ng et al.⁹⁵⁴

Abbreviations: ADAMTS, ADAM metalloproteinase with thrombospondin motifs; AFSC, amniotic fluid stem cell; AKP, alkaline phosphatase; BAX, Bcl-2 associated X protein; BMSC, bone marrow-derived mesenchymal stem/stromal cell; CAP, chondrocyte-affinity peptide; CCP3, cleaved caspase-3; CGRP, calcitonin gene-related peptide; CLSM, confocal laser scanning microscopy; col I, type I collagen; col II, type II collagen; COMP, cartilage oligomeric matrix protein; DMM, destabilization of medial meniscus; DRG, dorsal root ganglion; ECM, extracellular matrix; EHG, exosome encapsulating hydrogel; ELF3, E74-like factor 3; ESC-MSC, embryonic stem cell-derived mesenchymal stem/stromal cell; Exo, exosome; GAG, glycosaminoglycan; GelMA, gelatin methacrylate; HWT, head withdrawal threshold; ICRS, International Cartilage Repair Society; IL, interleukin; iNOS, inducible nitric oxide synthase; IPFP-MSC, infrapatellar fat pad-derived mesenchymal stem/stromal cell; iPSC-MSC, induced pluripotent stem cell-derived mesenchymal stem/stromal cell; KGN, kartogenin; LWT, leg withdrawal threshold; MDA, malondialdehyde; MIA, monosodium iodoacetate; MMP, matrix metalloproteinase; MRI, magnetic resonance imaging; MSC, mesenchymal stem/stromal cell; NC, negative control; NGF, nerve growth factor; OA, osteoarthritis; OARSI, Osteoarthritis Research Society International; OCD, osteochondral defect; OCN, osteocalcin; OCT, optical coherence tomography; P75NTR, p75 neurotrophin receptor; PBS, phosphate-buffered saline; PCNA, proliferating cell nuclear antigen; PRP, platelet-rich plasma; PWL, paw withdrawal latency; PWT, paw withdrawal threshold; RUNX2, runt-related transcription factor 2; SD rat, Sprague-Dawley rat; SMA, smooth muscle actin; SM-MSC, synovial membrane-derived mesenchymal stem/stromal cell; SOD, superoxide dismutase; TGF-β1, transforming growth factor beta 1; TIMP, tissue inhibitor of metalloproteinase; TMJ, temporomandibular joint; TNF-α, tumor necrosis factor alpha; TrkA, tropomyosin receptor kinase A; UC-MSC, umbilical cord-derived mesenchymal stem/stromal cell; µCT, micro computed tomography.




FIGURE 28 Overview of the studies. The exosomes tested in the included studies were derived from human, murine, or rabbit amniotic fluid stem cells (AFSCs), embryonic stem cell-derived mesenchymal stem/stromal cells (ESC-MSCs), induced pluripotent stem cell-derived MSCs (iPSC-MSCs), bone marrow-derived MSCs (BMSCs), polydactyl BMSCs, synovial membrane-derived MSCs (SM-MSCs), infrapatellar fat pad-derived MSCs (IPFP-MSCs), umbilical cord-derived MSCs (UC-MSCs), chondrocytes, dendritic cells, platelet-rich plasma (PRP), and serum. The exosomes were administered to the osteoarthritic joint through intra-articular injection or scaffold implantation. The exosomal bioactive compounds played an important role in cartilage and subchondral bone repair and regeneration. Overall, exosome therapy restored joint function, reduced joint pain, and improved the joint macroscopic, histological, and biochemical features. Reprinted with permission from Ng et al.⁹⁵⁴




FIGURE 29 Mechanisms of MSC-EVs on tendon repair. Reprinted with permission from Liu (2021).⁹⁶⁷

and collagen type 1 expression.⁹⁷⁴ The BMSC-EV group exhibited decreased expression of collagen type III and increased vascularity compared to both the PBS control group and the BMSC group, suggesting that the BMSC-EV group had superior healing quality.⁹⁷⁴ Similarly, tendon recovery was faster in rats when fibrin glue was added to BMSC-EVs transplanted to the patellar tendon window incision than in either the fibrin glue alone or untreated groups. Evidence of this included more Col1a1 gene expression and better collagen fiber alignment 4 weeks after the damage.⁹⁷⁴ Yu et al.⁹⁷⁵ showed that the biomechanical and histological regeneration of the rat patellar tendon was greatly enhanced by the injection of BMSC-EVs in fibrin glue, indicating that BMSC-EVs may improve the biomechanical characteristics of damaged tendons. Another study found that local distribution of BMSC-EVs in the hydrogel improved tendon-bone healing in mice, with greater fibrocartilage and better biomechanical properties than the hydrogel-only and control groups.⁹⁷⁴

Intratendinous injections of exosomes from tendon-derived stem cells (TDSCs) was also shown to accelerate tendon regeneration both biomechanically and histologically in a collagenase-induced rat Achilles tendinopathy model.⁹⁷⁶ It has been shown that part of the mechanism by which repair occurs is via macrophage polarization

and immune cell regulation.⁹⁷⁶ Previous research has shown that MSC-Exos have the ability to shift the macrophage response to tendon damage from an inflammatory M1 phenotype to a regenerative M2 phenotype via paracrine processes.^{977–979} Figure 29 and Table 31 summarize and highlight the benefits of exosomes for tendon repair.

6.4 | Cutaneous wounds

A number of studies and systematic reviews have now pointed to the benefit of exosomes on cutaneous wound healing.^{985–996} A study by Zeng and Liu pointed to a number of studies to date that have favored better healing with stem cell-derived exosomes.⁹⁹⁵ In that study, exosomes were investigated on the many phases of wound healing. It was found that exosomes improved the hemostasis phase by favoring faster blood clot formation,⁹⁹⁷ aided the inflammatory phase by decreasing inflammation and improving M2 macrophage polarization,^{128,998} stimulated the proliferative phase by targeting fibroblast activity, improving collagen type III formation,⁹⁹⁹ new vessel formation,⁷⁸⁴ and lastly enhanced the remodeling phase by minimizing scarring.^{720,779} (Table 32). Interestingly, a number of studies have further pointed to the benefit of exosomes on diabetic

TABLE 3 1 Studies investigating the therapeutic potential of MSC-EVs on tendon repair:

Animal model	EV (source/dosage/frequency)	Time point	Results	References
Achilles tendon transection and repair model in nude mouse	EEM (Human BMSC-derived EVs were used to educate human macrophages to M2 phenotype)/1 × 10 ⁶ /immediately after repair/once	Day 7 and Day 14 post-injury	EEM treatment substantially improved the biomechanical properties of the healing tendon but showed no improvement in collagen fiber organization. The EV or BMSC treatment showed biological responses had but have no effects on the biomechanical properties of tendon nor collagen fiber organization. The EEM treatment reduced the expression of type I collagen and type I/type III collagen ratio while the expression of type III Collagen remained unchanged compared to the injured control. The injection of EVs similarly reduced the expression of type I collagen in the healing tendon. However, it had no treatment effects on the expression of type III collagen and type I/type III collagen ratio	Chamberlain et al. ⁹⁸⁰
A case study of a horse suffering from suspensory ligament injury	Allogeneic MVs derived from 5-azacytidine (AZA) and resveratrol (RES)-treated ASCs isolated from horse with metabolic syndrome/dosage not reported/twice (7 days after injury and 9 months after first injection)	10 months and 12 months after first injection	The application of MVs improved lesion filling, angiogenesis, and elasticity of injured tissue	Kornicka-Garbowska et al. ⁹⁸¹
Rat patellar tendon window injury model	Rat BMSC-EVs/25 µg in 10 µL fibrin glue/once	2 and 4 weeks post-injury	The local administration of BMSC-EVs promoted tendon healing with improvement in collagen fiber Alignment, expression of tendon matrix genes and tenogenic differentiation markers compared to the fibrin glue-only group and untreated group Inflammation and accumulation of apoptotic cells were suppressed while the numbers of tendon progenitor cells increased at the healing site	Shi et al. ⁹⁷⁴
Rat Achilles tendon collagenase-induced tendinopathy model	Rat Achilles tendon TDSC-derived Ex/dosage not reported/twice a week at 1-week post-injury	5 weeks post-injury	The injection of exosome derived from TDSCs promoted tendon repair both histologically and biomechanically compared to the injury group and the treatment effects were comparable to the TDSCs treatment	Wang et al. ⁹⁷⁶

TABLE 31 (Continued)

Animal model	EV (source/dosage/frequency)	Time point	Results	References
Rat Achilles tendon injury model	Rat BMSC-EVs/ 2.8×10^{12} or 8.4×10^{12} /once	30 days	BMSC-EVs accelerated tendon repair in a dose-dependent manner. Higher doses of BMSC-EVs produced better restoration of tendon architecture, tendon-fiber alignment, and lower vascularity. Higher concentration of EVs induced higher expression of collagen type I and lower expression of collagen type III compared to the PBS control group and BMSC group	Gissi et al. ⁹⁶⁹
Mouse Achilles tendon two-third partial transection and repair model	Mouse naïve and IFN- γ -primed ASC-EVs in collagen sheet/ $5-6 \times 10^9$ from 5×10^5 ASCs/once	1, 3, and 7 days post-injury	Compared with the untreated control group, primed ASC-EVs, not their unprimed counterparts, further reduced the rate of post-repair tendon gap formation and rupture and promoted collagen formation at the injury site. Primed ASC-EVs, but not unprimed EVs, attenuated the early tendon inflammatory response after injury via modulation of the macrophage inflammatory response	Shen et al. ⁹⁸²
Mouse Achilles tendon-bone reconstruction model	Mouse BMSC-derived Ex in hydrogel/dosage not reported/once	7 days, 14 days and 1 month post-surgery	At 1 month after surgery, there was more fibrocartilage in the exosome group than in the other groups. The biomechanical properties of the tendon-bone junction significantly improved in the exosome group	Shi et al. ⁹⁸³
Rabbit rotator cuff repair	ADSC-derived Ex/dosage not reported/once	18 weeks post-surgery	The injection of Ex reduced fatty infiltration, increased the histological score with more fibrocartilage and improved biomechanical properties of the tendon-bone junction compared to the saline group	Wang et al. ⁹⁸⁴
Rat patellar tendon window injury model	Rat BMSC-derived Ex in fibrin glue/ $20 \mu\text{g}$ in $10 \mu\text{L}$ /once	3 days, 1, 2, 4 weeks post-injury	The transplantation of Ex improved the histological scores, promoted the proliferation of resident tendon stem cells, enhanced the expression of tenomodulin and type I collagen as well as biomechanical properties of neotendon	Yu et al. ⁹⁷⁵

Note: Reprinted with permission from Po and Liu.⁹⁶⁷

TABLE 3.2 Effects of mesenchymal stem cell exosomes on cutaneous wound healing.

Phase	Exosome source	Nomenclature	Related exosomal cargo	Secreted factors or expressed genes affected	Outcome	References
Hemostasis phase	Human mesenchymal stem cells (MSCs) from the umbilical cord	EVs	-	Phosphatidylserine (+)	Umbilical MSCs and extracellular vesicles derived from them have a reasonably high procoagulant potential	997
Inflammatory phase	Human jaw bone marrow-derived MSCs and bone marrow MSCs	Exosomes	miR-223	TNF- α \downarrow IL-10 \uparrow	Accelerated wound healing in mice Induced M2 macrophage polarization (CD206 ⁺ macrophage \uparrow)	998
	Human umbilical cord (UC)-MSCs	Exosomes	let-7b	TLR4, p-p65, iNOS \downarrow p-STAT3, p-AKT, ARG1 \uparrow	Alleviated inflammation and enhanced diabetic cutaneous wound healing in rats Induced M2 macrophage polarization Inhibited TLR4 signaling pathway	128
	Human UC-MSCs	Exosomes	miR-181c	TNF- α , IL-1 β , TLR4, p65, p-p65 \downarrow IL-10 \uparrow	Reduced burn-induced inflammation in rats Reduced neutrophil and macrophage infiltration (MPO ⁺ cell, CD68 ⁺ cell \downarrow) Inhibited TLR4 signaling pathway	135
	Human menstrual blood-derived MSCs (MenSCs)	Exosomes	-	iNOS \downarrow ARG1, VEGF \uparrow	Resolved inflammation and ameliorate cutaneous non healing wounds in diabetic mice Induced M2 macrophage polarization	720
Proliferative Phase	Human bone marrow MSC-derived exosomes	Exosomes	TGF- β /Smad	TGF- β 1, Smad2, Smad3, Smad4 \downarrow TGF- β 3, Smad7 \uparrow	Effectively promoted the cutaneous wound healing by inhibiting the TGF- β /Smad signal pathway	1000
	Human adipose MSCs (ASCs)	Exosomes	-	N-cadherin, cyclin 1, PCNA, collagen I/III, elastin \uparrow	Facilitated cutaneous wound healing via optimizing the characteristics of fibroblasts	779
	Human ASCs	Exosomes	-	Collagen I/II, TGF- β 1/3, MMP1/3 α -SMA \downarrow	Promoted ECM reconstruction in cutaneous wound repair by regulating the ratios of collagen type III: type I, TGF- β 3:TGF- β 1, and MMP3:TIMP1, and by regulating fibroblast differentiation to mitigate scar formation	999
	Human fetal dermal MSCs	Exosomes	Jagged 1	Collagen I/III, elastin, fibronectin mRNA \uparrow	Promoted wound healing by activating the ADF cell motility and secretion ability via the Notch signaling pathway	1001
	Human UC-MSCs	Exosomes	Wnt4	CK19, PCNA, collagen I \uparrow	Stimulated the AKT pathway to protect immortalized keratinocytes from heat-induced apoptosis stimulated the AKT pathway to protect immortalized keratinocytes from heat-induced apoptosis	1002

TABLE 32 (Continued)

Phase	Exosome source	Nomenclature	Related exosomal cargo	Secreted factors or expressed genes affected	Outcome	References
	Human UC-MSCs	Exosomes	Akt, ERK, STAT3	HGF, IGF1, NGF, SDF1 ↑	Promoted the proliferation and migration of fibroblasts in normal and chronic wounds. This effect was positively correlated with the dose of exosomes	713
	Induced pluripotent stem cell-derived MSCs	Exosomes	-	Collagen ↑	Increased the secretion of collagen by HaCaT cells to accelerate skin cell proliferation	1003
	Adipose mesenchymal stem cells (ADSCs)	Exosomes	AKT/HIF-1α	-	Promoted the proliferation and migration of HaCaT cells by regulating the activation of the AKT/HIF-1α signaling pathway, thus promoting wound healing	1004
	Human UC-MSCs	Exosomes	-	PARP-1, PAR ↑	Suppressed HaCaT cell apoptosis induced by H ₂ O ₂ by restraining the nuclear translocation of apoptosis-inducing factor (AIF) and promoting poly (ADP-ribose) (PAR) and poly ADP ribose polymerase 1 (PARP-1) expression	1005
	Human adipose-derived MSCs (adMSC-Exo)	Exosomes	miR-125a	Angiogenic inhibitor delta-like 4 (DLL4) ↓	Transferred miR-125a to endothelial cells and promoted angiogenesis by repressing DLL4	1006
	Mouse BM-MSCs	Exosomes	miR-17 miR-23a miR-125b	TNF-α, IL-1β, iNOS, TLR4, IRAK1, p65 ↓ ARG1, IL-10, TGF-β ↑	Decreased the threshold for thermal and mechanical stimuli in mice Increased nerve conduction velocity, the number of intraepidermal nerve fibers, myelin thickness, and axonal diameters	1007
	Rat BMSCs	Exosomes	-	MDA, HIF1α, NOX2, Caspase 3, BAX, PARP1, MPO, ICAM1, IL-1β, NF-κB ↓ SOD, CAT, GPX, HO-1, BCL2, IL-10, bFGF, HGF, SOX9, VEGF ↑	Decreased histopathological score of kidney injury in rats Reduced the levels of blood urea nitrogen (BUN) and Creatinine Reduced the level of oxidative stress Increased antioxidant status Reduced apoptosis and inflammation Improved regeneration and enhanced angiogenesis	1008
	Human endometrial MSCs	Exosomes	-	Tie2, VEGF, Ang1, Ang2 ↑	Increased the expression of angiogenesis markers, including Tie2, VEGF, Ang1, and Ang2, and increased the proliferation, migration, and angiogenesis of HUVECs	1009

(Continues)

TABLE 3 2 (Continued)

Phase	Exosome source	Nomenclature	Related exosomal cargo	Secreted factors or expressed genes affected	Outcome	References
	Human umbilical cord mesenchymal stem cells (hUC-MSCs)	Exosomes	-	Ang2 ↑	huc-MSC-Ex-derived Ang-2 plays a significant role in tube formation of HUVECs and promotion of angiogenesis	1010
	Human UC blood-MSCs	Exosomes	-	Ang, Ang1, HFG, VEGF ↑	Human umbilical cord blood (UCB)-MSC-derived exosomes pretreated with thrombin could accelerate skin wound healing in rats with full-thickness wounds. Exosomes from human UCB-MSCs increased angiogenesis factors, such as VEGF, HGF, and Ang1, and decreased TNF-α and IL-6	1011
	Human UC-MSCs	Exosomes	Wnt4	β-catenin, N-cadherin, PCNA, Cyclin D3 ↑	Enhanced angiogenesis in rats through the Wnt4/β-catenin pathway. When the expression of Wnt4 was knocked out by shRNA, the pro-angiogenic effect of hUC-MSC-derived exosomes was eliminated	114
	Human UC-MSCs	Exosomes	-	α-SMA, collagen I ↓	Increased the formation and maturation of new blood vessels at the wound site, although the mechanism is still unclear	784
	Human UC-MSCs	Exosomes	GSK3β-Wnt/β-catenin	-	Alleviated hepatic IRI by transporting miR-1246 via regulated GSK3β-mediated Wnt/β-catenin pathway	1012
Remodeling Phase	Human gingival MSCs	Exosomes	-	Collagen ↑	Reduced the formation of scars by inhibiting the accumulation of mouse myofibroblasts	1013
	Adipose mesenchymal stem cells (ASCs)	Exosomes	-	N-cadherin, cyclin-1, PCNA collagen I, III ↑	Facilitates cutaneous wound healing via optimizing the characteristics of fibroblasts	779
	ASCs	Exosomes	ERK/MAPK	Matrix metalloproteinases-3 (MMP3) ↑	A promoted ECM reconstruction in cutaneous wound repair by regulating the ratios of collagen type III: type I, TGF-β3:TGF-β1, and MMP3:TIMP1, and by regulating fibroblast differentiation to mitigate scar formation	999
	MenSCs	Exosomes	-	iNOS ↓ ARG1, VEGF ↑	Resolved inflammation and ameliorated cutaneous non-healing wounds in diabetic mice Induced M2 macrophage polarization	720

Note: Reprinted with permission from Zeng and Liu.⁹⁹⁵

Abbreviations: AIF, apoptosis-inducing factor; Ang, angiotensin; ASC, human adipose mesenchymal stem cell; BMMSC, bone marrow MSC; BMSC, bone marrow-derived stem cells; CTGF, connective tissue growth factor; DLL4, delta-like 4; ECM, extracellular matrix; EGF, epidermal growth factor; EVs, extracellular vesicles; FD, human fetal dermis; FGFs, fibroblast growth factors; HaCaTs, human keratinocytes; hBM, human bone marrow; HDFs, human dermal fibroblasts; HGF, hepatocyte growth factor; hUC, human umbilical cord; HUVECs, human umbilical vein endothelial cells; IFNs, interferons; IGF-1, insulin growth factor 1; IL, interleukin; ILVs, intraluminal vesicles; iNOS, inducible nitric oxide synthase; iPSC, induced pluripotent stem cell; JMMSC, jaw bone marrow MSC; KGFs, keratinocyte growth factors; LPS, lipopolysaccharide; MMP, matrix metalloproteinases; MSCs, mesenchymal stem cells; MVBs, multivesicular bodies; NAP-2, neutrophil activating peptide-2; PDGF, platelet-derived growth factor; PF-4, platelet factor 4; ROS, reactive oxygen species; SDF-1α, stromal cell-derived factor-1; TGF-β, transforming growth factor; TIMP1, tissue inhibitor of metalloproteinase 1; TNF-α, tumor necrosis factor alpha; VEGF, vascular endothelial growth factor.

wounds, burn healing, and post-irradiation (Table 33).⁹⁹⁴ A study by An et al.⁹⁸⁷ demonstrated that exosomes improved healing via the following four mechanisms:

- Control of inflammation and immune system
- Encouraging wound angiogenesis
- Accelerating the growth and regeneration of skin cells
- Controlling the process of collagen remodeling to prevent excessive scar formation

In summary, previous studies have revealed that exosomes had a positive impact on fat grafting, diabetic ulcer wound healing, scarless wound healing, and could effectively be utilized in conjunction with carriers such as hydrogels or fibrin (Figure 30).⁹⁸⁷ Figure 31 summarizes how exosomes assist during cutaneous wound healing.

6.5 | Dermatology and skin regeneration

Much like the effects of exosomes on cutaneous wound healing, there are many benefits to utilizing MSC-Exos for dermatological conditions. Their ability to be utilized for skin healing has been popularized not only in medicine but also in the field of facial esthetics since it encompasses a multi-billion-dollar industry where specific exosomes for dermatological conditions (Dermasomes) have been utilized for skin health and antiaging benefits.

6.5.1 | Atopic dermatitis

Often referred to as atopic eczema, atopic dermatitis (AD) is a common inflammatory skin condition marked by persistent, uncontrollably high inflammatory reactions.¹⁰²⁴ About 8% of adults and 20% of children have been diagnosed with the condition.¹⁰²⁵ Unfortunately, it remains unknown what exact molecular pathway leads to the pathogenesis of the illness. Major pruritus and erythematous lesions are the disease's primary symptoms having a major negative impact on a patient's quality of life.¹⁰²⁴ Pharmacological intervention, such as corticosteroids and calcineurin inhibitors, is the mainstay of therapy for AD. A number of negative side effects as well as the development of drug resistance have been reported. As a result, several investigators have stated that the development of novel, more potent, and less hazardous treatments is critically needed in this field. MSCs have been useful in AD healing, much like in the repair of chronic wounds.¹⁰²⁴ MSC-Exos provide a stronger treatment option for AD as they are more stable and have less immunogenicity than the cells from which they are isolated, yet they perform the same biological tasks.¹⁰²⁶ This circumvents the majority of issues related to live MSC-based treatment options.

In recent years, greater Th2 cytokine levels have been primarily linked to increased vulnerability to AD (inside-outside theory).^{186,1026,1027} Because of this, the majority of research has focused on lowering Th2-mediated immune responses. However,

current research has shown a clear link between aberrant gene expression-driven epidermal barrier defects and immune response dysregulation, such as increased Th2 cytokine levels. Cho et al. showed that human ADSC-Exos may provide a potential cell-free therapeutic approach for the treatment of AD.¹⁸³ In an in vivo animal model, researchers found that ADSC-Exos reduced the production of various inflammatory cytokines, including IL-4, IL-23, IL-31, and TNF-alpha. In an oxazolone-induced dermatitis model, Shin et al. found that subcutaneous ASC-exosome injection substantially decreased TSLP, IL-5, IL-13, TNF-, IFN-, and IL-17.¹⁸⁶

6.5.2 | Psoriasis

The most prevalent chronic inflammatory skin condition, psoriasis, affects around 125 million individuals globally.¹⁰²⁸ Impaired keratinocyte differentiation and proliferation are linked to the condition. Furthermore, it has been observed that the immune system cells massively invade troubled areas.¹⁰²⁹ It manifests as red squamous plaques that are restricted to the head, elbows, knees, and sacroiliac area.¹⁰²⁸ Unfortunately, the exact cause of psoriasis is still unknown. However, past studies indicate that the condition is influenced by T cells and dendritic cells. The immunology of the disease has been well explained by Lowes et al.¹⁰³⁰ EVs have a role in the pathophysiology of psoriasis, just as they do in AD.¹⁰²⁹ Research is still being done on the use of EVs as therapeutic agents to control psoriasis. On the other hand, Zhang and colleagues reported encouraging results using exosomes produced from human umbilical cord mesenchymal stem cells (huc-MSCs-Exo).¹⁰³¹ Injection of huc-MSCs-Exo subcutaneously decreased the levels of IL-17, IL-23, CCL20, and STAT3/p-STAT3. Additionally, exosomes inhibited the maturation activation of dendritic cells and prevented IL-17's positive impact on keratinocytes.¹⁰³¹

6.5.3 | Medical esthetics

People are now more than ever more conscious of their physical appearance. Consequently, self-esteem is often affected by apparent skin aging processes, scars, or hair loss. This has spurred the growth of aesthetic medicine, which is continuously looking for novel, minimally invasive, and highly successful cosmetic interventions. Increased research on exosome use for skin rejuvenation, scar removal, and hair loss has been spurred by promising results with exosomes in regenerative medicine (discussed later in this section). In a study titled: "Stem cell-derived exosomes: A groundbreaking development in cosmetic dermatology," Shen et al. discovered that exosomes produced from stem cells play a critical role in several aspects of skin cosmetology, including wound healing, skin aging, and scar formation.¹⁰³² Today many commercial entities exist selling dermal exosomes (aka Dermasomes).

TABLE 33 Studies that evaluated an in vivo role for MSC-EVs in wound healing.

Study	EV source	Model	Findings
Fang et al. ⁷⁸⁴	Human UC-MSC	Mouse skin wound <ul style="list-style-type: none"> Local injection In vitro dermal fibroblasts	EVs reduced scar formation and myofibroblast accumulation EVs suppressed TGF- β induced myofibroblast formation. EVs were enriched in miR-21, miR-23a, miR-125b, and miR-145. miRNA delivery reduced TGF- β /SMAD2 signaling in fibroblasts
Hu et al. ⁷⁷⁹	Human AD-MSC	Mouse skin wound <ul style="list-style-type: none"> Local injection Mouse skin wound <ul style="list-style-type: none"> Intravenous injection In vitro fibroblasts	EVs improved rate of wound healing, increased Col1 and Col3 mRNA on Day 3 and Day 5 post wounding, and decreased Col1 and Col3 mRNA on Days 7 and 14 EVs migrated to wound site (Days 5–14) and spleen and promoted wound healing EVs promoted fibroblast proliferation and migration, increased mRNA for N-cadherin, COL1, COL3, and elastin
Zhang et al. ¹⁰¹⁴	Human AD-MSC	Mouse skin wound <ul style="list-style-type: none"> Local injection In vitro fibroblasts	EVs improved rate of wound healing, decreased scar size, and neoangiogenesis EVs promoted fibroblast proliferation and migration, and increased mRNA for COL1, COL3, MMP1, FGF2, and TGF- β 1. Fibroblasts had increased p-AKT. Application of PI3K/AKT inhibitor Ly294002 abrogated the EV-induced effects on fibroblasts
He et al. ⁹⁹⁸	Human BMSC	Mouse skin wound <ul style="list-style-type: none"> Intravenous injection In vitro human monocytes/macrophages	EVs promoted wound healing and polarization of macrophages to M2 phenotype EVs promoted M2 macrophage polarization in part through transfer of miR-223
Ren et al. ¹⁰¹⁵	Human AD-MSC	Mouse skin wound <ul style="list-style-type: none"> Local injection In vitro fibroblasts, keratinocytes (HaCaT), and endothelial cells (HUVEC)	EVs accelerated wound healing, re-epithelialization, collagen deposition, and neovascularization EVs promoted proliferation and migration, and stimulated AKT and ERK signaling
Cheng et al. ¹⁰¹⁶	Human UC-MSC	Mouse skin wound <ul style="list-style-type: none"> Local injection In vitro dermal fibroblasts and keratinocytes (HaCaT)	EVs accelerated re-epithelialization and promoted collagen fiber maturation EVs promoted proliferation and migration. The effect was blocked by miR-27b inhibitor. Proposed miR-27b acts by suppressing ITCH, thereby activating JUNB/IRE1 α
Jiang et al. ¹⁰¹⁷	Human BMSC	Mouse skin wound <ul style="list-style-type: none"> Local injection 	EVs from MSCs with TSG-6 overexpression (TSG-6-EVs) and knock-down (TSG-6-KD-EVs). EVs reduced scar formation, reduced production of TGF- β 1, Collagen I and III, and α -SMA protein, and suppressed SMAD2/3 signaling. TSG-6-EVs enhanced the effect of EVs, the effect was lost in TSG-6-KD-EVs, and when TSG-6 neutralizing antibodies were present
Liu et al. ¹⁰¹⁸	Mouse BMSC	Mouse skin wound <ul style="list-style-type: none"> Topical in pluronic F127 hydrogel In vitro mouse macrophages	Topical EVs accelerated wound healing, limited inflammatory infiltrate, and decreased scar size EVs polarized macrophages toward M2 phenotype. Conditioned media from EV treated macrophages promoted fibroblast proliferation and migration

TABLE 33 (Continued)

Study	EV source	Model	Findings
Qiu et al. ¹⁰¹⁹	Mouse BMSC	Mouse skin wound <ul style="list-style-type: none"> Local injection In vitro endothelial cells (HUVECs)	EVs from MSCs treated with EVs from neonatal serum and adult serum. MSC-EVs accelerated wound healing and promoted neoangiogenesis. Neonatal serum stimulated MSC-EVs showed more robust effect MSC-EVs promoted HUVEC proliferation, migration, and tube formation, and increased p-AKT and p-eNOS. Neonatal serum stimulated MSC-EVs showed more robust effect
Zhang et al. ¹⁰⁰⁴	Human AD-MSC	Mouse skin wound <ul style="list-style-type: none"> Local injection In vitro HaCaT keratinocytes	EVs promoted mouse wound healing, proposed to occur in AKT/HIF-1 α dependent fashion EVs promoted HaCaT keratinocyte proliferation
Zhao et al. ¹⁰⁰⁵	Human UC-MSC	Mouse skin wound <ul style="list-style-type: none"> Local injection In vitro keratinocytes (HaCaT)	EVs enhanced re-epithelialization and neoangiogenesis EVs stimulated keratinocyte proliferation, migration, and suppressed ROS induced apoptosis. Proposed effect was through suppression of AIF nuclear translocation and PARP-1 activation
Li et al. ¹⁰²⁰	Human AD-MSC	In vitro human hypertrophic scar fibroblasts	EVs decreased collagen deposition, transdifferentiation of fibroblasts-to-myofibroblasts, and formation of hypertrophic scar. EVs were noted to express miR-192-5p, which can suppress IL-17RA/SMAD axis
<i>Diabetic wounds</i>			
Wang et al. ⁷²¹	Mouse AD-MSC	Mouse diabetic wound <ul style="list-style-type: none"> Topical in complex hydrogel (Pluronic F127, oxidative hyaluronic acid, and Poly-L-lysine) 	EVs improved wound healing and neovascularization. The effect was improved when EVs were loaded in complex hydrogel
Li et al. ⁷¹⁴	Mouse BMSC	Mouse diabetic wound <ul style="list-style-type: none"> Local injection In vitro human fibroblasts from diabetic foot ulcers and health control	EVs from MSCs overexpressing lncRNA H19 (H19-EVs). Only H19-EVs promoted wound healing, decreased inflammatory infiltrate, and increased granulation tissue formation H19-EVs reduced miR-152-3p expression in fibroblasts from diabetics and increased PTEN expression
Shi et al. ⁷²³	Mouse AD-MSC	Mouse diabetic wound <ul style="list-style-type: none"> Local injection In vitro endothelial cells (HUVECs)	EVs accelerated wound healing, increased angiogenesis, suppressed apoptosis, and increased autophagy markers SIRT1 and LC3. The effects were further enhanced with EVs from mmu_circ_0000250 overexpressing MSCs EVs promoted HUVEC survival under high glucose conditions and increased autophagy. This was enhanced by loading with mmu_circ_0000250, which was shown to increase SIRT1 mediated autophagy
Yang et al. ⁷²⁶	Human UC-MSC	Mouse diabetic wound <ul style="list-style-type: none"> Topical in Pluronic F127 hydrogel 	EVs accelerated wound healing and angiogenesis, increased expression of VEGF and TGF- β 1

(Continues)

TABLE 33 (Continued)

Study	EV source	Model	Findings
Pomatto et al. ¹⁰²¹	Human BMSC AD-MSC	Mouse diabetic wound • Topical in carboxymethylcellulose In vitro fibroblasts, keratinocytes, and endothelial cells	AD-MSC-EVs, but not BMSC-EVs, promoted the rate of wound healing. Comparative in vivo analysis of scar and angiogenesis was not performed BMSC-EVs promoted proliferation of keratinocytes and endothelial cells, and promoted viability of fibroblasts, keratinocytes, and endothelial cells. AD-MSC-EVs promoted only the proliferation of endothelial cells. Protein and miRNA analysis indicated BMSC-EVs are enriched for proliferative factors, whereas AD-MSC-EVs are enriched in pro-angiogenic factors
Ti et al. ¹²⁸	Human UC-MSC	Rat diabetic wound • Local injection In vitro human monocytes (THP-1)	EVs from LPS preconditioned MSCs (LPS Pre-EVs) decreased inflammatory cell infiltration and polarized macrophages toward M2 LPS Pre-EVs induced M2 polarization. EVs transferred Let-7b, reducing TLR-4 expression and NF-κB activation
Li et al. ⁷¹⁸	Human AD-MSC	Rat diabetic wound In vitro human epithelial progenitor cells (EPC)	EVs from MSCs overexpressing NRF2 (NRF2-EVs). Endothelial progenitor cells (EPC) + NRF2-EVs promoted wound healing better than EPC + AD-MSC-EVs, and both were better than EPC alone or control EVs decreased EPC senescence under high glucose conditions. NRF2-EVs inhibited inflammatory cytokines and ROS
Ding et al. ⁷¹⁰	Human BMSC	Rat diabetic wound • Local injection In vitro endothelial cells (HUVECs)	EVs from deferoxamine stimulated MSCs (DFO-EVs). EVs promoted wound healing and neoangiogenesis, and DFO-EVs were more effective DFO-EVs were more potent stimulators of HUVEC proliferation and tube formation than EVs. DFO-EVs proposed to transfer miR-126 to HUVECs, which suppresses PTEN, and thereby activates AKT signaling
Liu et al. ⁷²⁴	Human BMSC	Rat diabetic wound • Local injection In vitro mouse macrophages (RAW264.7)	EVs from MSCs treated with melatonin (MT-EVs). EVs promoted wound closure, Collagen I and III expression, and M2 macrophage polarization; MT-EVs enhanced the effect of EVs MT-EVs were more potent than EVs at polarizing macrophages to M2 phenotype
Yu et al. ⁷²⁵	Human BMSC	Rat diabetic wound • Local injection In vitro endothelial cells (HUVECs)	EVs from MSCs treated with atorvastatin (ATV-EVs). EVs promoted wound healing and angiogenesis. ATV-EVs were more effective EVs promoted proliferation, migration, and tube formation, increased VEGF secretion, and activated AKT/eNOS signaling. ATV-EVs produce a larger magnitude effect compared to standard EVs. ATV-EVs proposed to work by upregulating miR-221-3p in endothelial cells
Burn wounds			
Shafei et al. ⁷¹¹	Human AD-MSC	Mouse burn wound • Topical in alginate hydrogel	EVs accelerated wound closure, increased epithelial thickness, collagen deposition, and neovascularization

TABLE 33 (Continued)

Study	EV source	Model	Findings
Zhang et al. ¹⁰²²	Human iPSC-MSC	Rat burn wound • Local injection	EVs accelerated re-epithelialization, reduced scar width, promoted collagen maturation, and stimulated neoangiogenesis. Effects depended on EV transfer of Wnt4
		In vitro fibroblasts and endothelial cells (HUVECs)	EVs stimulated proliferation and migration, stimulated Collagen I and III, and elastin secretion, and promoted tube formation
Li et al. ¹³⁵	Human UC-MSC	Rat burn wound • Intravenous injection	EVs reduce inflammation following burn wounds. EVs transfer miR-181c and reduce TLR4 signaling
		In vitro mouse macrophages (RAW264.7)	EVs suppress LPS-induced macrophage inflammation

Note: Reprinted with permission from Bray et al.⁹⁹²




FIGURE 30 Representative photographs shown of full-thickness excision wound area of the rat treated with PBS (control) or hBMSCs-Exos. Reprinted with permission from Jiang et al.¹⁰⁰⁰

6.5.4 | Potential capacity of stem cell-derived exosomes in skin aging

Aging skin is caused by both external and internal factors. Age dependency and natural aging are referred to as intrinsic aging, which is influenced by genes, while external aging, sometimes referred to

as photoaging, is caused by environmental variables.¹⁰³³ UVB light is the most powerful environmental element that causes skin photoaging. Cell damage is caused by ultraviolet B induced DNA mutations and oxidative stress.^{1034,1035}

The ECM and dermal fibroblast activity are important factors of youthful skin. Studies show that reductions in collagen type I and III




FIGURE 31 Role of mesenchymal stem cell-derived extracellular vesicles (MSC-EVs) in wound healing. (A) MSC-EVs in hemostasis. MSC-EVs contain pro- and anticoagulant factors, which balance and regulate blood coagulation. (B) MSC-EVs in inflammation. MSC-EVs support anti-inflammatory processes, reducing reactive oxygen species (ROS) synthesis, alleviating apoptosis, and inducing macrophage phenotype change from pro-inflammatory (M1) to anti-inflammatory (M2). (C) MSC-EVs in proliferation. MSC-EVs stimulate fibroblast migration and proliferation to the wound site, resulting in raised levels of extracellular matrix (ECM) components synthesis. Also, MSC-EVs can promote vascularization. (D) MSC-EVs in remodeling. Bone marrow MSC-EVs (BMSC-EVs) increase collagen I production, smooth muscle actin (SMA) and fibroblast differentiation to myofibroblast; however, they decrease collagen III synthesis. Besides, BMSC-EVs boost new cutaneous appendage formation. Adipose mesenchymal stem cell extracellular vesicles (AdMSC-EVs) act opposite and lead to scar reduction. Reprinted with permission from Narauskaite et al.¹⁰²³

synthesis and fragmentation of collagen fibrils define the molecular structure of aged skin.¹⁰³⁶ One of the most significant ECM proteins that provides tissues a younger, more full appearance is collagen. The key to skin elasticity is elastin.¹⁰³⁷ But as fibroblasts age, their capacity to proliferate and synthesize collagen is diminished.¹⁰³⁸ Furthermore, the elevated production of matrix metalloenzymes expedites the degradation of collagen, ultimately resulting in the formation of wrinkles and other visible signs of skin aging.¹⁰³⁴ Numerous researchers have shown that MSC-Exos play a significant role in mitigating skin aging and deterioration.¹⁰³⁹

Exosomes have been shown to have a greater impact on fibroblast senescence than other constituents.⁹⁰¹ Exosomes produced from human-induced pluripotent stem cells (iPSCs), for instance, have the ability to prevent HDFs from aging. iPSCs-Exos have the ability to control the amount of MMP-1/3 and enhance the production of type I

collagen in senescent HDFs.¹⁰³⁴ MMP-1, functioning as a collagenase, participates in the production of a secretase enzyme that breaks down interstitial collagens I, II, and III.¹⁰⁴⁰ Multiple studies have shown a correlation between the increase in fibroblast growth and the activation of collagen production, which is influenced by the presence of TGF- β and PDGF.^{1034,1041} Previous studies have shown that human dermal fibroblasts take up MSC-Exos and stimulate cell migration and synthesis of skin collagen I and elastin.¹⁰⁴² Zhang and colleagues showed that exosomes produced from ADSCs enhanced fibroblasts' levels of type I and type III collagen and accelerated wound healing via the PI3K/Akt signaling pathway.¹⁰¹⁴ These investigations demonstrate that stem cell-derived exosomes have significant potential use in postponing the aging of the skin by stimulating the production of collagen. Figure 32 highlights some benefits of exosomes on skin healing.




FIGURE 32 Main composition of exosomes and their functions on skin injuries. Despite the different origin and mode of biogenesis, exosomes display a similar appearance, and common composition. Exosomes are featured by tetraspanins, receptors or cell-type specific proteins, lipids, adherin proteins, intracellular trafficking proteins. Deep analysis of extracellular vesicle composition reveals that they convey various cargoes, including nucleic acids, proteins (biogenesis elements, chaperones, and signaling pathway molecules) and lipids, which all vary widely between cells and conditions. The vast information will directly affect the fate and function of exosomes in skin injuries. Exosomes participate in multiple cutaneous diseases including laceration, burn, aging, diabetic wound healing, scar formation, skin autoimmune disease, and skin cancers, through different biological pathways. Reprinted with permission from Shi et al.¹⁰⁴³

6.5.5 | Stem cell-derived exosomes in repair, regeneration, and angiogenesis

Inflammation, hyperplasia, and remodeling are all part of the intricate and dynamic process of skin wound healing. Type M2 polarization of macrophages regulates inflammation. Exosomes released by human mesenchymal stem cells are capable of lowering neutrophil counts and preventing the recruitment of macrophages during the inflammatory phase. Therefore, exosomes from human MSCs may lessen the inflammatory response and are crucial for skin repair and regeneration during the proliferative stage because they encourage fibroblast migration, proliferation, and endothelial cell creation.¹¹⁴ Bin and colleagues found that huc-MSC exosomes activate the Wnt4/β-catenin signaling pathway to promote endothelial cell migration, proliferation, and formation.¹¹⁴ Huc-MSC-Exos also promoted β-catenin nuclear translocation and activity via the Wnt4 pathway, hence increasing fibroblast migration and proliferation.¹⁰⁰² Jieyuan discovered that the angiogenic activities of endothelial cells might be facilitated by exosomes released from endothelial progenitor cells via the Erk1/2 signaling pathway, ultimately stimulating cutaneous wound healing and regeneration.¹⁰⁴⁴

Exosomes generated from human umbilical cord blood (UCB-Exos) have been shown to inhibit phosphatase and tensin homolog (PTEN) and sprout homolog 1 (Spry1), hence promoting wound healing and reducing scarring. These effects could be connected to UCB-Exos high expression of miR-21-3p.¹⁰⁴⁵ Exosomes produced

from menstrual blood-derived mesenchymal stem cells were shown by Dalirfardouei to promote neoangiogenesis via VEGF. Accordingly, the acceleration of epithelial regeneration in mice was directly linked to the overexpression of the NF-κB p65 subunit and the activation of the NF-κB signaling pathway.⁷²⁰ These findings imply that exosomal treatment created from stem cells could be a potential therapeutic strategy and that administering stem cell-derived exosomes may improve endothelial cells' angiogenic potential, fibroblast migration, and proliferation.

6.5.6 | Potential capacity of stem cell-derived exosomes in reduced scar formation

Scar development takes place mostly during the wound remodeling period. Early gestation fetuses are observably capable of scarless wound healing as compared to adult wounds.⁹⁹⁹ The ultimate objective of cosmetic dermatology is to achieve scarless recovery. Scarless tissue differs from scar tissue in many ways. It is characterized by the presence of fine reticular collagen, a lower degree of crosslinking, less inflammation, and a reduced number of myofibroblasts. The ratios of TGF-β3 to TGF-β1, type III to type I collagen, and MMPs to matrix metalloproteinase tissue inhibitors (TIMPs) were all greater in scar-free tissue.⁹⁹⁹

Activated fibroblasts replace the injured epithelial or endothelial cells during the process of tissue regeneration and repair after injury.




FIGURE 33 Exosomes derived from multiple sources could regulate wound healing by affecting effector cells. The exosomes from various sources, including MSCs, keratinocytes, endothelial cells, immune cells, and body fluid, could regulate the wound healing process through different mechanisms. Especially, the UBE2O mRNA in saliva-exos, the miR-135a in amnion-derived MSC-exos, the MALAT1 and mmu_circ_0000250 in adipose-derived MSC-exos, and the H19 lncRNA in bone marrow-derived MSC-exos played positive roles in enhancing the functions of main effector skin cells to accelerate wound healing. The miR-20b-5p in exosomes of patients plasma and miR-15a-3p in exosomes from patients blood inversely impaired the functionality of the endothelial cells, exerting healing delay effects. Mesenchymal Stem Cells, MSCs. Reprinted with permission from Xiong et al.¹⁰⁴⁹

Fibroblasts have the capacity to overproduce fibronectin, hyaluronic acid, type I and III collagens, and other ECM components, which may result in scarring and matrix overgrowth.¹⁰⁴⁶ Exosomes have been shown in several studies to have a significant role in preventing the development of scars (Figure 33).^{720,1047}

During wound healing, HDFs develop into myofibroblasts. Following wound healing, a significant fraction of myofibroblasts undergo apoptosis. Myofibroblasts, however, are still active and multiplying in hypertrophic scar patients.¹⁰⁴⁸ Research has shown that human ADSC-Exos have the ability to control the ratios of TGF- β 3 and collagen type III to type I: TGF- β 1, MMP3, and TIMP-1 by inhibiting fibroblast development into myofibroblasts.⁹⁹⁹

Chen et al.¹⁰⁴⁸ demonstrated that by preventing the function of miR-200C3p in fibroblasts, lncRNAs LNCs5088 from M2 macrophage exosomes may contribute to the development of scars. This provides evidence that exosomes may facilitate intercellular communication in the management of hyperplasia in scarring.¹⁰⁴⁸ Therefore, stem cell-derived exosomes have shown to be a successful noninvasive scar reduction therapy. Nevertheless, further studies are required to fully understand how stem cell-derived exosomes contribute to the reduction of scars.^{999,1017}

6.6 | Dental regeneration

A number of systematic reviews have now investigated the use of exosomes for the treatment of various dental tissues.¹⁰⁵⁰⁻¹⁰⁶⁰ In

fact, part 3 of this 3-part series is dedicated entirely to the use of exosomes in dentistry. In that systematic review, a total of 944 articles were identified using exosomes in the dental field for either their regenerative/therapeutic potential or for diagnostic purposes derived from salivary exosomes. In total, 113 research articles were selected for their regenerative potential (102 in vitro, 60 in vivo, and 49 studies were both). Therapeutic exosomes were most commonly derived from dental pulps, periodontal ligament cells, gingival fibroblasts, stem cells from exfoliated deciduous teeth, and the apical papilla have all been shown to facilitate the regeneration of a number of tissues including bone, cementum, the periodontal ligament, nerves, orthodontic tooth movement, and temporomandibular joint disorders, among others. Results demonstrate that the use of exosomes led to positive outcomes in 100% of studies. In the bone field, exosomes were found to perform equally as well or better than rhBMP2. Periodontitis animal models were treated with simple gingival injections, and benefits were even observed when the exosomes were administered intravenously. Exosomes are much more stable than growth factors commonly utilized in the dental space and, as such, were shown to be far more resistant to degradation by periodontal pathogens found routinely in a periodontitis environment. Comparative studies in the field of periodontal regeneration found better outcomes for exosomes even when compared to their native parent stem cells. In total, 47 diagnostic studies have revealed a role for salivary/crevicular fluid exosomes in the diagnosis of birth defects, cardiovascular disease, diabetes, gingival recession detection, gingivitis, irritable bowel syndrome, neurodegenerative disease, oral

lichen planus, oral squamous cell carcinoma, oro-pharyngeal cancer detection, orthodontic root resorption, pancreatic cancer, periodontitis, peri-implantitis, Sjögren syndrome, and various systemic diseases. Hence, dental exosomes (Periosomes) were described as possessing “remarkable” potential, serving as a valuable tool with significant advantages for dental clinicians.

6.7 | Erectile dysfunction

A study by Khodamoradi et al. 2022 titled: “Exosomes as Potential Biomarkers for Erectile Dysfunction, Varicocele, and Testicular Injury”, examined the possibility of using exosomes as biomarkers for a number of ailments, including erectile dysfunction.¹⁰⁶¹ It was determined that exosomes are a viable therapeutic method since they may contain cargo such as specific medications and therapeutic compounds. As discussed earlier (Section 5.1.8), erectile dysfunction is a blood flow issue which is the reasoning behind its association with diabetes specifically as well as other noncommunicable diseases such as smoking, cardiovascular diseases, and obesity.⁷⁰¹ Therefore, the use of exosomes in the treatment of arterial inflow or venous outflow problems will undoubtedly have a positive effect concerning erectile dysfunction. Additionally, exosome therapy for several male sexual and reproductive diseases has been tested in a number of clinical trials.¹⁰⁶¹ Gaining insight into the significant function of exosomes in the reproductive system may provide new opportunities for developing novel treatment approaches and diagnostic markers.¹⁰⁶² This may be a promising method for managing conditions including testicular damage, varicocele, and erectile dysfunction.¹⁰⁶¹

6.8 | Hair regrowth

Alopecia is a prominent reason for seeking dermatological consultations. However, the management of this condition continues to be challenging. Androgenic alopecia is the predominant cause of non-scarring hair loss, which affects around 80% of Caucasian men and 50% of women by the age of 70. On the other hand, central centrifugal cicatricial alopecia is the leading cause of hair loss among African Americans.¹⁰⁶³ Patients with alopecia have severe psychological effects.¹⁰⁶³

Because alopecia is so common, a lot of research has been done with the aim of developing safe and efficient therapies. Topical minoxidil and oral/topical finasteride are two FDA-approved therapies that may partially stop hair loss but not fully stimulate hair growth. Additionally, side effects and difficulty sticking to regular treatment regimens may further restrict the usage of these drugs.¹⁰⁶³ Although low-level laser treatment has FDA approval as well, its effectiveness is unknown.¹⁰⁶⁴ Patients may be able to have surgical therapy via hair transplantation, but the expense may prevent them from doing so.¹⁰⁶⁴ Therefore, current research is focused on regenerative therapies that not

only stop hair loss but also encourage growth. In many clinical trials, platelet-rich plasma injection therapy (PRP), a minimally invasive treatment, has been shown to improve hair density and volume.¹⁰⁶⁵

In a study titled: “Exosome Therapy in Hair Regeneration: A Survey of the Literature on the evidence, difficulties, and prospects for the future”, Kost et al. thoroughly examined the exosomes' capacity for regenerative hair growth.¹⁰⁶³ Exosome therapies have been shown in preclinical trials to have distinct advantages in regenerative medicine and hair loss therapy. Clinical investigations support the safety of using exosomes in medicine, although there is a shortage of information regarding the effectiveness and safety of exosome treatment for alopecia.¹⁰⁶³

The majority of preclinical data supporting the use of exosomes in alopecia comes from exosomes produced by dermal papilla cells (DPCs), a collection of specialized mesenchymal cells found at the base of hair follicles (HFs).¹⁰⁶⁶ They have been shown to be essential for controlling the creation, development, and cycling of HF.¹⁰⁶⁷ Studies in mice have demonstrated that injecting DPC exosomes accelerates the start of HF anagen while slowing down HF catagen. Immunohistochemistry studies revealed elevated amounts of β -catenin and Sonic Hedgehog (Shh) in the treated skin, which control the hair cycle, as well as H&E-stained tissue sections revealing anagen VI phase in the treated animals. Treatment of outer root sheath cells with DPC exosomes further confirmed the overexpression of β -catenin and Shh10 in vitro. There is evidence that DPC-exosome therapy enhances the migration and proliferation of cells found in the outer root sheath, namely those cells found in the hair follicle bulge.¹⁰⁶⁷ Similar to what Kwack et al. found when they subcutaneously gave DPC exosomes to mice, exosomes have been shown to promote protracted anagen of hair follicles.¹⁰⁶⁸ Research in mice with alopecia areata suggests that DPC exosomes may reduce inflammation in hair follicles and stop hair loss.¹⁰⁶⁹

In addition, DPCs are critical for HF stem cell differentiation. Epithelial cells near the follicular base die during catagen and telogen, but the DP stays alive and moves up to the HF bulge. There, it releases signals that initiate the anagen phase by stimulating the differentiation of HF stem cells and regenerating HFs.¹⁰⁷⁰ Yan et al. showed that the co-culture of HF stem cells with DPCs stimulated HF stem cell development. Additionally, transmission electron microscopy observation revealed that DPC exosomes were adhered to the surface of HFSCs during co-culture.¹⁰⁷¹ Ultimately, they found that 111 miRNAs were expressed differently in DPC exosomes than in DPCs, suggesting that DPC exosomes are important for HF regeneration.¹⁰⁷¹ It is worth mentioning that DPC exosomes have demonstrated the ability to influence ADSCs, causing their transformation into DPC-like cells that produce genes related to hair growth, such as β -Catenin and hair-inducing miRNAs.¹⁰⁷²

There are a number of other exosome sources known to exist for hair growth treatment. In vitro, the maintenance of hair human dermal papilla cells was shown using exosomes derived from hair outer root sheath cells.¹⁰⁷³ Furthermore, compared to a control

group, exosomes from the outer root sheath increased markers of hair growth induction by a ratio of 2.1, 1.7, and 1.3, respectively.¹⁰⁷³

Lastly, exosomes produced by immune cells have also shown potential in the therapy of alopecia. In a mouse model of alopecia areata, exosomes from myeloid-derived suppressor cells, which inhibit T cell growth, increased FoxP3 and arginase 1 levels to prevent T-cell hyperreactivity.¹⁰⁷⁴ Additionally, in a mouse model and in vitro studies using human hair follicles, macrophage exosomes have shown noticeably higher amounts of hair-inductive indicators of DPCs.¹⁰⁷⁵ It has also been shown that exosomes produced by the bacterium *Leuconostoc holzapfelii*, which was isolated from human

scalp tissue, regulate hair development via the Wnt/ β -catenin signal transduction pathway.¹⁰⁷⁶

Zhou et al.¹⁰⁶⁷ found that in mice models, DP cell exosomes derived from healthy human scalp specimens sped up the beginning of anagen and postponed catagen, leading to higher levels of β -catenin and Shh. Accordingly, human dermal papillae exposed to activated human dermal fibroblasts (hDFs) release the non-Wnt ligand Norrin via stimulated extracellular vesicles (st-EVs).¹⁰⁷⁷ Frizzled-4 (Fzd4), the particular receptor for Norrin given by hDF, is thought to promote the β -catenin pathway's subsequent activation, which leads to the observed increased hair follicle development ex vivo (Figure 34).¹⁰⁷⁷




FIGURE 34 St-EVs improve human hair growth. (A) Schematic representation of DP cell treatment by EVs isolated from DF stimulated by 2GFs. (B) Individual HFs were dissected from adult human scalp tissue and placed in culture in the presence of PBS (nontreated [NT]), cyclosporin A (10^{-7} M), ctrl-EVs, or st-EVs (5×10^9 particles per milliliter) and two different concentrations of bFGF and PDGF-AA named 2GF-dose1 and 2GF-dose2 at D0 and D4. Pictures of follicles were taken at three points (D0, D4, and D7) and the length of each follicle was measured over time. (C) Quantification of the length of hair fibers overtime relative to NT. Data represent experiments performed on different donors at Days 4 and 7 following HF extraction (D4/D7): NT, ctrl-EVs, st-EVs, and cyclosporin A were obtained from four donors at D4 and three donors at D7; 2GF-dose1 and 2GF-dose2 were obtained from three donors at D4 and two donors at D7 and are expressed as mean \pm SD. * $p < 0.05$ and *** $p < 0.001$ compared with NT HFs using one-way analysis of variance followed by Dunnett's multiple comparison test for D4 and Kruskal-Wallis test followed by Dunn's multiple comparison test for D7. 2GF, two growth factors; bFGF, basic fibroblast growth factor; D0/4/7, Day 0/4/7; DF, dermal fibroblast; DP, dermal papilla; EVs, extracellular vesicles; HF, hair follicle; PDGF-AA, platelet-derived-growth factor A; st-EVs, EVs from stimulated DF. Reprinted with permission from Riche et al.¹⁰⁷⁷

When combined, research indicates that exosomes from different tissue sources may be very advantageous for hair regeneration and rejuvenation inside the HF dermal papilla, outer root sheath, and HF bulge cells.

6.9 | Spinal cord injury and intervertebral disc repair

A potentially fatal and catastrophic lesion to the spinal cord, spinal cord injury (SCI) results in either temporary or permanent alterations to the cord and partial or whole loss of motor, sensory, and autonomic functioning.^{1078,1079} SCI commonly results in paralysis, sometimes known as quadriplegia or paraplegia, with sensory impairment occurring below the site of the damage.¹⁰⁸⁰ In most cases, it causes patients and healthcare systems to bear very heavy psychological and financial costs.^{1081,1082} It also has adverse effects on most fundamental body processes, including breathing, the operation of the bowel and bladder, hormone production, and sexual function. This is because the peripheral nerve system and brain are no longer connected.¹⁰⁸³ The prevalence and incidence of SCI are estimated to be 236–4187 per million persons globally, with up to 770000 new cases annually. Males under the age of 30 are more likely to have SCI than females.^{1084–1086}

The first mechanical insult, which may be produced by physical forces including contusion, compression, transection, or stretching of the spinal column, often leading to disruption of the spinal cord and results in primary damage.^{1087–1091} An instantaneous mechanical damage to the spinal cord is referred to as a primary injury since it is an irreversible procedure.^{1092,1093} A series of secondary injuries follow the initial damage, worsening the spinal cord's health.^{1094,1095} Secondary injury occurs shortly after the initial mechanical injury

and is characterized by local vascular damage, ionic changes, thrombosis, edema, ischemia, progressive hemorrhage, oxidative stress caused by the release of free radicals, lipid peroxidation, excitotoxicity, and cell death facilitated by apoptosis and cell necrosis.¹⁰⁷⁹ Moreover, the inflammatory response and excessive growth of glial cells, which occur after the subsequent suppressive environment and scar tissue, impede the regrowth of nerve fibers and restrict the effectiveness of treatment.^{1096,1097}

Recovery from SCI remains significantly restricted, even though recent therapeutic advancements in SCI care have shown some improvement in patients' quality of life.^{1098,1099} For the purpose of treating SCI, three main groups of pathologic targets may be distinguished. First, at the site of the original trauma, surgical decompression, and the removal of mechanical spinal cord compressing material.^{1079,1100–1104} Second, anti-inflammatory therapies for the area around the damaged spinal cord.^{1105–1107}

Third, the ultimate objective for treating spinal cord injuries is axonal regrowth at the location of the damage.¹⁰⁸³ Following the initial SCI, the damaged lesion is heavily invaded by macrophages, which help to create a cavity of injury (COI) surrounding the site of injury, stopping neuronal regrowth.^{1108–1111} Scarring interferes with axonal regrowth,¹¹¹² and the COI lesions that were filled with fluid prevent axons from traversing the liquid-filled COI without the aid of bridge-like structures.¹¹¹³ Furthermore, granulomatous infiltration surrounding the injured spinal cord, known as arachnoiditis, aids in the development of a mature scar devoid of astrocytes or other glial cells.¹⁰⁹⁷ Patients with SCI continue to have a dismal prognosis, a high death rate, and a noticeably shorter life expectancy.¹¹¹⁴ Figure 35 highlights the phases of injury.¹¹¹⁵

Astrocytes are crucial to the process of SCI because they may either impede or facilitate the central nervous system's (CNS) recovery.^{1116–1120} A1 and A2 reactive astrocyte phenotypes are




FIGURE 35 Schematic diagram for damage stages and responses in spinal cord injury. Reprinted with permission from Kim et al. 2021.¹⁰⁸⁰

pre-existing and are brought on by ischemia and neuroinflammation. A1 astrocytes may cause neuronal and oligodendrocyte death because of their neurotoxic effects on myelin, synapses, and neurons. They are often generated just after SCI that is triggered by IL-1, TNF-alpha, and C1q.¹¹²¹ On the other hand, by upregulating the production of certain neurotrophic factors, A2 astrocytes provide a protective effect.¹¹¹⁶ As a result, one possible therapeutic option for SCI might be selective suppression of A1 astrocytes. Moreover, it has recently been shown that red blood cells (RBCs) around SCI lesions are eliminated by reactive astrocytes via phagocytosis. This process, known as astrocytic erythrophagocytosis, is thought to aid in the quick evacuation of dispersed RBCs from the area of injury to stop macrophage aggregation and the ensuing harmful inflammation. MicroRNAs (miRNAs) have been implicated in tissue damage and regeneration processes in recent times, and several miRNAs have garnered interest as possible targets for the treatment of SCI.

6.9.1 | Regenerative potential of stem cell-derived extracellular vesicles in spinal cord injury

Spinal cord injury is a severe ailment that is progressive and very difficult to treat, requiring urgent medical treatment owing to its intricate pathophysiology and impact on social status and financial load. A paper published recently by Herbert et al.¹¹²² found that the mechanism of action of exosomes on spinal cord injury involves at least eight processes including (1) tissue sparing and neuroprotection, (2) alleviation of oxidative stress, (3) induction and progression of angiogenesis, (4) pericyte role restoring integrity of the blood-spinal cord barrier, (5) combating endoplasmic reticulum stress, (6) modulation of inflammatory response, (7) activation of autophagy, and (8) attenuation of apoptosis. Interestingly, animal studies have demonstrated significant benefit when utilized typically within 1 hour of injury but also following subsequent IV infusions up to 21 days (Tables 34 and 35). Noteworthy, protocols could be developed within an early time frame for athletes playing contact sports such as rugby or American football, where the likelihood of injury is more significant. Protocols have been proposed and are in development to improve patient outcomes following trauma. Figure 36 highlights the benefits of exosomes in SCI.

6.9.2 | Use of exosomes in intervertebral disc degeneration

In today's world, low back pain (LBP) is a common health issue that has serious socioeconomic ramifications.^{1157,1158} According to epidemiological data, 80% of individuals will, at some point in their lives, have lower back discomfort.¹¹⁵⁹ LBP is thought to be one of the main factors limiting the amount of labor that individuals under 45 may be capable of.^{1160,1161} In the United States, the direct cost of treating LBP is estimated to be over 30 billion US dollars, while the

indirect socioeconomic losses are estimated to be around 100 billion US dollars.¹¹⁶²

Previous studies¹¹⁶²⁻¹¹⁶⁵ found that the most common cause of LBP currently is IVD. Despite the fact that the exact cause of IVD is still unknown, age, metabolic diseases, mechanical stress, trauma, illness, diet, and genetic susceptibility are some of the contributing variables.^{211,1160}

Currently, bed rest, the use of nonsteroidal anti-inflammatory medicines (NSAIDs) and other analgesics, lumbar discectomy, and interbody fusion are the major clinical treatments for LBP.¹¹⁶² Nevertheless, these methods just concentrate on providing temporary relief for the symptoms rather than directly addressing the underlying cause, resulting in the inability to halt or stop the advancement of IVDD.^{212,1166} Hence, innovative therapeutic approaches are required for the treatment of IVDD. Recently, therapies based on MSCs and exosomes derived from MSCs have shown promising benefits and may have practical applications.^{1165,1167-1170}

Noor et al.¹¹⁷¹ found that exosomes produced from MSCs not only have an essential function in tissue repair and regeneration but also possess a longer-lasting, more powerful, and easier-to-maintain role when compared to parent cells. While the use of MSC-Exos in IVD therapy is still in its infancy, there is rising interest in this approach. An increase in oxidative stress, decreased ECM content, inflammation, and cell death are the pathophysiological characteristics of IVD. Targeting these four aspects of the illness, MSC-Exos may help to improve all symptoms related to IVD. The results of previous research on IVD regeneration and repair utilizing exosomes produced from MSCs are compiled in Table 36. In 2022, research by Lu et al.¹¹⁷² compiled 16 research studies and discovered that MSC-Exos were obtained from various cell sources for the treatment of IVD. These sources included human-induced pluripotent stem cells (1 study), unspecified tissues (2 studies), BMSCs (10 studies), ADSCs (1 study), umbilical cord MSCs (1 study), and placenta-MSCs (1 study). In a preclinical model of IVD, MSC-Exos exhibited comparable therapeutic benefits despite variable tissue sources.¹¹⁷² These effects are primarily attributed to the restoration of extracellular matrix integrity, promotion of cell proliferation, reduction of cell death, regulation of inflammatory response, and attenuation of oxidative stress.¹¹⁷²

6.10 | Vascular regeneration

Several pathologies affecting the vasculature, particularly the microvasculature, lead to a lack of physiological homeostasis control of patency and adequate perfusion to meet tissue metabolic needs. Most diseases involving failing organs and tissues include microvascular dysfunction as an essential underlying component. Vascular decreased angiogenic potential density, endothelial dysfunction, ER stress, mitochondrial dysfunction, oxidative stress, increased senescence, and apoptosis are pathogenic variables that contribute to this dysfunction.¹¹⁸⁸⁻¹¹⁹⁰

TABLE 34 Mechanistic overview of EV therapy and molecular outcomes in SCI.

Route of administration	Window period	Therapeutic intervention	Dose	TF/cytokines/signaling pathway	Outcome	References
Tail vein	1 h	BMMSC exosome	100 µg	↓ C3, C4b, C6, C5, Mbl, Cfp, C1q and Cfh ↓ p-p65 (NF-κB) and p-IκBα	↓ Complement system activation and inflammation	1123
Tail vein	1 h and day 7	hUCB MSC-derived Nano-Vesicle	25 µg	↓ TNFα, IL1b, Nos2 ↑ Arg1, IL10, and VEGF	M1 → M2 neuroprotection, anti-inflammation, angiogenesis	1124
Tail vein	Every 3 days till 27 days	BMMSC exosome	200 µg	↓ Bax, Cleave caspase-3 ↑ Wnt/b-catenin ↑ Bcl-2, caspase-3 ↑ LEF-1 and TCF-1	↓ Neuronal apoptosis ↑ Motor recovery	1125
Tail vein	Immediate	NSC-EV-14-3-3t	200 µg	↓ Cleave caspase-3, TNF-α, IL-1β and IL-6 ↑ Bcl-2	↓ Apoptosis, inflammation ↑ Autophagy	1126
Tail vein	Immediate	UCB-CD34 ⁺ CM	-	↓ GFAP, IL-1b, TNF-a, and MPO ↑ IL-10	↓ Oxidative stress, inflammation, astrogliosis	1127
Lumbar cistern	24 h	Subventricular zone-derived EV	1 µg	↓ TNF-α IL-1β, IL-18, and IL-6	Motor function recovery and ↓ inflammation	1128
Tail vein	30 min	BM MSC exosome	40 µg	↓ C3 and GFAP TNF-α, IL-1α, and IL-1β, p-IKBα and nuclear p65	↓ A1 astrocyte, inflammation, apoptosis	1129
Tail vein	Immediate	GIT1-overexpressing BMMSC exosome	200 µg	↑ PI3K/AKT ↑ CS56 ↓ TNF-a, IL-1b and IL-6	↓ Inflammation, Apoptosis and glial scarring	1130
Tail vein	Immediate	Epidural fat (EF)-mesenchymal stem cells (MSCs)	-	↓ NLRP3, ASC, and active-caspase-1, IL-1β, IL-18, and TNF-α	↓ Inflammation, expression of activated microglia	1131
Intra thecal	1 day	"Neuro cells" (HSC, MSC)	100 µL	↓ GFAP	↓ Astrogliosis	1132
Tail vein	1 day	BMMSC EV	200 µL EV (200 µg/mL)	↓ NF-κB	↓ Pericyte migration ↑ Integrity of the BSCB	1133
Tail vein	Immediate	NSC-derived exosomes	100 µg	-	↓ Apoptosis ↑ Nerve regeneration	1134
Tail vein	-	MSC miR-21 and miR-19b exosome	-	↓ PTEN	↓ Apoptosis	1135
Tail vein	1 h	BMMSC exosome	100 µg	↓ FasL gene	↓ Apoptosis	1136
			200 µg	-	↑ Autophagy	1137
			100 µg	↓ IL-1a, TNF-a, and IL-17B and IL-36b	↓ Inflammation	1138

(Continues)

TABLE 34 (Continued)

Route of administration	Window period	Therapeutic intervention	Dose	TF/cytokines/signaling pathway	Outcome	References
Tail vein	24 h	BMMSC mir133b exosome	100 µg	↑ GAP43, pERK1/2, CREB and STAT3	↑ Axonal outgrowth	1139
Intrathecal	Week 1,2,3	hWJMSC CM	-	↑ GAP43 ↓ Irf5, Mrc1 ↓ Il1b, TNFα	↓ Glial scar ↑ Axonal outgrowth	1140
Scaffold implantation	Immediate	hMSC exosomes	-	↓ GFAP ↓ iNOS	↓ Astrogliosis ↓ Inflammation	1141
Tail vein	Immediate	Bone MSC miR-216a-5p exosome	200 µg	↑ Arg1, CD206, YM1/2, PI3K/AKT (↓ TLR4/NF-κB)	↓ Inflammation Functional recovery	1142
Tail vein	1 h	BM MSC CM	30 µg (15 × 10 ⁹ EV)	↓ IL-1β	Improved spatial learning impairments ↓ Inflammation	1143
Tail vein	1 h	BMSC-miRNA-29b exosome	100 µg	↑ GAP-43, NF200	Neuronal regeneration	1144
In vitro	-	BMSC CM	100 µM	↓ Notch1, Hes1 Bax, Cleave caspase-3 ↑ Bcl-2, caspase-3	↓ Oxidative stress and apoptosis	1145
Tail vein	30 min	hUC-MSC exosome	200 µg	↓ iNOS, TNF-α, IL-6, IFN-γ, G-CSF, MCP-1, and MIP-1α, ↑ IL-10, IL-4	Macrophage polarization M1 → M2 ↓ Inflammation	1146
Tail vein	30 min	MSC miR-126 exosome	100 µg (1 × 10 ¹⁰)	↓ SPRED1 and PIK3R2 ↓ Bax, Cleave caspase-3 ↑ Bcl-2 ↑ VEGF	↑ Angiogenesis ↑ Neurogenesis ↓ Apoptosis	1147

Note: Exosomes exert their therapeutic potentials in SCI through the regulation of various molecular mechanisms. The administration, therapeutic window period, route, and dosage are vital players determining the extent of functional recovery. There is dynamic regulation of molecules, crosstalk between pathways, and homeostasis involved in the outcomes to contain the secondary injury that is the key for functional recovery in SCI (↑ – Upregulation, ↓ – downregulation). Reprinted with permission from Herber et al.¹¹²²

Abbreviations: ANGPT1, angiotensin; Arg1, arginase 1; ASC, apoptosis-associated Speck-like protein; BDNF, brain-derived neurotrophic factor; bFGF, fibroblast growth factor; GDNF, glial cell line-derived neurotrophic factor; GFAP, glial fibrillary acidic protein; GM-CSF, granulocyte macrophage colony-stimulating factor; HGF, hepatocyte growth factor; HLA-G, human leukocyte antigen-G; HSC, hematopoietic stem cell; hUCBMSC, human umbilical cord blood-derived mesenchymal stem cell; hUC-MSC, human umbilical cord mesenchymal stem cell; Iba-1, ionized calcium binding adaptor molecule 1; IGF-1, insulin-like growth factor; IL10, interleukin 10; IL-4, interleukin 4; IL-6, interleukin 6; iNOS, inducible nitric oxide synthase; iPSC, induced pluripotent stem cell; lncRNA, long non-coding RNA; MBP, myelin basic protein; MCP-1, monocyte chemoattractant protein-1; miRNA, microRNA; MSC, mesenchymal stromal cells; NF-κB, nuclear factor kappa B; NGF, nerve growth factor; PGE-2, prostaglandin E2; PI3K, phosphatidylinositol-3-kinase biosynthesis class F protein; PIK3R2, phosphoinositide-3-kinase regulatory subunit 2; PKA, protein kinase cAMP-dependent; PPAR-γ, peroxisome proliferator-activated receptor gamma; SOD, superoxide dismutase; SPRED, sprouty-related EVH1 domain-containing protein 1; TGF-β, transforming growth factor beta; TLR4, Toll-like receptor 4; VASH, vasohibin 1; VEGF, vascular endothelial growth factor.

TABLE 35 List of publications on EVs' therapeutic effect in SCI.

EVs sources	Routes	Models	↑ Score	Proposed mechanisms
Rat BMSC	iv	Contusion	4.5 at Week 4	Anti-apoptosis, anti-inflammation, pro-angiogenesis ¹¹⁴⁹
Rat BMSC	iv	Contusion	4.5 at Week 4	neuroprotection, reduce A1 astrocytes, anti-inflammation, anti-apoptosis ¹¹²⁹
Rat BMSC	iv	Contusion	3 at Week 4	Anti-microglia and A1 neurotoxic reactive astrocytes, anti-inflammation, anti-apoptosis, reduce scar, pro-angiogenesis ¹¹⁵⁰
Rat BMSC	iv	Hemisection	3 at Week 4	Inhibit complement activation ¹¹²³
Rat BMSC	iv	Contusion	2 at Week 4	Anti-apoptosis ¹¹²⁵
Rat BMSC	iv	Hemisection	6 at Week 4	Anti-apoptosis ¹¹³⁶
Rat BMSC	iv	Contusion	6 at Week 4	Inhibit pericyte migration, decrease BSCB permeability ¹¹³³
Rat BMSC (miR-133b-enriched)	iv	Compression	3 at Week 2	Decrease RhoA expression, axon growth ¹¹³⁹
Rat BMSC (miR-29b-enriched)	iv	Contusion	11 at Week 8	Neuroprotection ¹¹⁴⁴
Rat BMSC (miR-126-enriched)	iv	Contusion	7 at Week 4	Anti-apoptosis, pro-neurogenesis, pro-angiogenesis ¹¹⁴⁷
Rat BMSC (hypoxic)	iv	Contusion	4 at Week 4	Macrophage polarization ¹¹⁴²
Human BMSC	iv	Contusion	3 at Week 2	Decreases reactive microglia and astrocytes ¹¹⁵¹
Human UC-MSC	iv	Contusion	2 at Week 8	Macrophage polarization, anti-inflammation ¹¹⁴⁶
Human adipose-MSC	iv	Contusion	5 at Week 4	Attenuate NLRP3 inflammasome activation ¹¹³¹
Rat-NSC	it	Compression	6 at Week 4	Anti-inflammation ¹¹²⁸
Rat-NSC	it	Contusion	7 at Week 4	Inhibit NLRP3 inflammasome complex formation ¹¹⁵²
Rat-NSC (14-3-3t-enriched)	iv	Contusion	4 at Week 4	Enhance autophagy, anti-apoptosis, anti-inflammation ¹¹⁵³
Rat-NSC (IFG-1-stimulated)	iv	Contusion	5 at Week 4	Anti-apoptosis, anti-inflammation ¹¹³⁴
Mouse NSC	iv	Contusion	4 at Week 4	Activate autophagy, anti-apoptosis, anti-inflammation, anti-microglia ¹¹²⁶
Mouse pericytes	iv	Contusion	3 at Week 2	Improve microcirculation, protect BSCB, anti-apoptosis ¹¹⁵⁴
Human BMSC (PTEN siRNA)	in	Full transection	7.5 at Week 8	Anti-inflammation, anti-scarring, pro-angiogenesis, axon growth ¹¹⁵⁵

Note: Reprinted with permission from Guo et al.¹¹⁴⁸

Abbreviations: ↑ Score, increased mean BBB or BMS locomotor score from EVs treatment groups, compared to untreated controls; BM, bone marrow; BSCB, blood-spinal cord barrier; in, intranasal; it, intrathecal; iv, intravenous; MSC, mesenchymal stem/stromal cells; NSC, neural stem cells; UC, umbilical cord.

Many therapeutic settings now use pharmacologic treatments that focus on a limited aspect to alleviate symptoms of pathology rather than adopting a complete strategy to tackle the underlying cause. In response to this issue, much attention has been directed to cellular treatments and cell-free therapies, such as exosomes, which can address the complex causes of vascular and microvascular dysfunction.¹¹⁸⁸

Furthermore, Kawasaki disease, antineutrophil cytoplasmic autoantibody (ANCA)-associated vasculitis, giant cell arteritis, Behçet's disease, and other multisystem autoimmune diseases are included in the diverse group of multisystem autoimmune disorders known as systemic vasculitis.^{1191,1192} Thrombosis, vascular stenosis/occlusion, aneurysm development, or bleeding are frequent complications of

vasculitis, which is defined by inflammation in specific-sized arteries.¹¹⁹³ Patients continue to face issues like early death, recurrence, co-morbidities, and a lower quality of life. Despite advances in earlier diagnosis and novel immunotherapies in recent decades,¹¹⁹⁴ the therapeutic potential of exosomes has attracted increased attention, and studies in this area are ongoing.¹¹⁹⁵

7 | INFECTIOUS DISEASES

It is interesting to note that exosomes have been effectively used to treat a variety of infectious disorders. Exosomes were first used to treat infectious disorders like hepatitis in an effort to replace tissues




FIGURE 36 Therapeutic effects of exosomes derived from different MSCs in the treatment of SCI. MSCs can be obtained from bone marrow, the umbilical cord, the amniotic membrane, and adipose tissue. Exosomes derived from MSCs have anti-inflammatory and anti-apoptotic effects, as well as inhibit A1 astrocytes, promote axonal regeneration and macrophage polarization, and protect the BSCB from spinal cord injury. Reprinted with permission from Liu et al.¹¹⁵⁶

and organs that the viruses had damaged.¹¹⁹⁶ Globally, morbidity and mortality have been significantly attributed to infectious illnesses; pneumonia and respiratory infections are among the leading causes of death globally.¹¹⁹⁷ Finding novel therapeutic approaches to fight infections and repair organ and tissue damage caused by infections is essential given the rising frequency of infectious disease outbreaks and the lack of efficient therapies.¹¹⁹⁸

Coronavirus disease 2019 or COVID-19 was one of the leading clinical uses of exosomes where numerous clinical trials were started

to minimize the 'cytokine storm' leading to many injuries/deaths. The global impact of the COVID-19 pandemic may be attributed to the emergence of a new virus known as severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) belonging to the coronavirus family. This infectious agent has affected a significant number of individuals on a global scale.^{1199,1200} Acute respiratory distress syndrome (ARDS) brought on by COVID-19 dramatically raises death rates in patients who are older or have chronic illnesses such as diabetes, heart, lung, or kidney problems. No effective treatment

TABLE 3.6 Preclinical studies investigating the therapeutic potential of stem cell-derived exosomes on IVDD.

Study	Cell type	Species of cell source	Size of EV	Isolation	Model	Modeling method	In vitro appraisement	In vivo appraisement	Additional manipulation
Qi et al. ¹¹⁷³	UC MSC	Human	-	Ultra-centrifugation	In vitro	-	<ul style="list-style-type: none">MSC exosome has the potential to alleviate HG induced extracellular matrix degradation via the p38 MAPK pathwayHG also significantly inhibited collagen II and aggrecan expression in NPMSCs, and leads to an increase in the rate of NPMSCs apoptosismiRNAs array results and bioinformatics analysis predicted that miR-221-3p, let-7a-5p, and miR-21-5p derived from hUC-MSCs-EXO may be closely related to extracellular matrix synthesis of NPMSCs	-	High glucose (HG) induced degradation of IVDD
Li et al. ¹¹⁷⁴	BM MSCs	Human	Around 100 nm	Ultra-centrifugation	In vitro	-	<ul style="list-style-type: none">In the pathological acid environment, MSC-derived exosome promotes the expression of chondrocyte extracellular matrix, collagen II, and aggrecan, and reduces matrix degradation by downregulating matrix-degrading enzymesMSC-derived exosomes were able to prevent and mitigate NPC apoptosis through repressing caspase-3 expression and attenuating caspase-3 cleavage induced by acidic pH	-	Acidic pH-induced NPCs apoptosis
Li et al. ¹¹⁷⁵	BM MSCs	Human	Around 100 nm	Ultra-centrifugation	In vitro	-	<ul style="list-style-type: none">BMSC-derived exosomes inhibit IL-1β-induced inflammation and apoptosis of AF cells by suppressing PI3K/AKT/mTOR signaling pathway-mediated autophagyBMSC exosomes supported AF cell viability after IL1-β treatment	-	IL1- β treatment (10 ng mL ⁻¹)
Yuan et al. ¹¹⁷⁶	PLMSCs	Human	30–150 nm	Ultra-centrifugation	In vitro In vivo	-	<ul style="list-style-type: none">MSC exosomes rich in antagomir-4450 ameliorates NPC damage by promoting proliferation and migrationEV-derived Antagomir-4450 decreased MMP13, IL6, IL1-β, CASP3 expression, and increased COL2 and ACAN expression 3. miR-4450 was significantly upregulated while ZNF121 was downregulated in IDD and miR-4450 exacerbated NPC damage by targeting ZNF121	<ul style="list-style-type: none">The EXO-antagomir-4450 attenuated IVDD damage by repressing miR-4450 and upregulating ZNF121 expressionThe EXO-antagomir-4450 could partially decline the pro-inflammatory factors and MMPs in IDD mice	TNF- α (10 ng mL ⁻¹) treatment

(Continues)

TABLE 36 (Continued)

Study	Cell type	Species of cell source	Size of EV	Isolation	Model	Modeling method	In vitro appraisement	In vivo appraisement	Additional manipulation
Zhu et al. ¹¹⁷⁷	BMMSCs	Rat	Mainly 109.3nm	Ultra-centrifugation	In vitro	-	<ul style="list-style-type: none"> MSC exosomes could inhibit TNF-α-induced increase of apoptotic process, activation of apoptotic proteins, imbalance of anabolism/catabolism levels, and accumulation of collagen I in NPC through the delivery of miR-532-5p RASSF5 is a direct target gene of miR-532-5p to mediate the cell apoptosis regulation 	-	TNF- α (20ng mL ⁻¹) treatment
Wen et al. ¹¹⁷⁸	BMMSCs	Rat	About 80nm	Ultra-centrifugation	In vitro In vivo	Injection of absolute ethanol into sub-endplate in rat tail to induce IVD degeneration	<ul style="list-style-type: none"> The expression of col II and Aggrecan, SA-β gal positive cells and apoptosis rate of NPCs were decreased after MSC exosomes intervention Reducing miR-199a carried by MSC exosomes led to an impaired protective effect of exosomes on NPCs miR-199a from MSC exosomes promotes repair by targeting GREM1 and downregulating the TGF-β pathway 	<ul style="list-style-type: none"> After MSCs-exosomes treatment, MMP-2, MMP-6, TIMP1 and TUNEL-positive cells were decreased, and miR-199a was increased in IDD mice BMMSCs-EVs promote proliferation of NPCs and inhibit apoptosis 	-
Zhu et al. ¹¹⁷⁹	BMMSCs	Mouse	80nm	Ultra-centrifugation	In vitro	-	<ul style="list-style-type: none"> MSCs exosomes alleviated NPCs apoptosis by reducing IL-1β-induced inflammatory cytokines secretion and MAPK signaling activation MSCs exosomes inhibited NPCs apoptosis and MAPK signaling by delivering miR-142-3p that targets mixed lineage kinase 3 (MLK3) 	-	IL1- β treatment (10ng mL ⁻¹)
Cheng et al. ¹¹⁸⁰	BMMSCs	Human	Average 87nm	Ultra-centrifugation	In vitro In vivo	The model of disc degeneration was established by needle puncture (needle-stab model)	<ul style="list-style-type: none"> MSC exosomes were taken up by NPCs and suppressed NPC apoptosis induced by TNF-α miR-21 in MSC exosomes alleviated TNF-α induced NPC apoptosis Delivery of miR-21 in MSC exosomes inhibited NPC apoptosis by targeting PTEN through PI3K-Akt pathway 	<ul style="list-style-type: none"> Intradiscal injection of MSC exosomes alleviated the NPC apoptosis and IVD degeneration in a rat model 	TNF- α (5 ng mL ⁻¹) treatment
Zhang et al. ¹¹⁸¹	MSCs	Human	Around 100nm	Ultra-centrifugation	In vitro In vivo	The IVDD model was established by annulus fibrosus (AF) needle puncture	<ul style="list-style-type: none"> MSCs-derived exosomes play an anti-pyoptosis role by suppressing the NLRP3 pathway MSC-derives exosomes treatment inhibit LPS-induced pyroptosis in NPCs miR-410 derived from MSC exosomes inhibit LPS-induced pyroptosis in NPCs 	<ul style="list-style-type: none"> MSCs-exosomes and miR-410 treatment alleviated the severity degree of IVDD 	LPS (5mmol L ⁻¹) treatment

TABLE 36 (Continued)

Study	Cell type	Species of cell source	Size of EV	Isolation	Model	Modeling method	In vitro appraisement	In vivo appraisement	Additional manipulation
Hingert et al. ¹¹⁸²	BMMSCs	Human	Average 144 ± 2.22 nm	Serial centrifugation	In vitro	/	<ul style="list-style-type: none">• MSC exosomes treatment increased cell viability and proliferation of degenerated disc cells• MSC exosomes treatment induced early production of crucial ECM components such as proteoglycan, aggrecan, and collagen type II• EV treatment suppressed apoptosis and the secretion of MMP-1 in disc cells	/	No
Sun et al. ¹¹⁸³	IMSCs	Rat	80–200 nm	Ultra-centrifugation	In vitro In vivo	The IVDD model was established by annulus fibrosus (AF) needle puncture	<ul style="list-style-type: none">• miR-140-5p enriched in IMSC exosomes played a pivotal role in the IMSC-sEV-mediated therapeutic effect by upregulating of anabolism markers of the ECM (collagen II and aggrecan), and downregulating of catabolism markers of the ECM (MMP-3 and ADAMTS-4)• IMSC-sEVs could rejuvenate senescent NPCs and restore the age-related function by activating the Sirt6 pathway in vitro	<ul style="list-style-type: none">• NPC senescence and IVDD were significantly improved after intradiscally injecting IMSC exosomes	No
Lu et al. ¹¹⁸⁴	BMMSC	Human	30–100 nm	Ultra-centrifugation	In vitro	-	<ul style="list-style-type: none">• EV treatment increased proliferation activity of NPC• EV treatment generate a healthier extracellular matrix by upregulating expression levels of anabolic/matrix protective genes (aggrecan, collagen II, sox-9 and TIMP-1) and downregulating matrix degrading genes (MMP1 and MMP3)	-	No

Xia et al. ¹¹⁸⁵	BMMSCs	Mouse	50–130 nm	Ultra-centrifugation	In vitro In vivo	An IVD degeneration rabbit model was induced with a fine needle puncture	<ul style="list-style-type: none">• BMSC-derived exosomes attenuate apoptosis in NP cells treated with H₂O₂• Exosomes dampen H₂O₂-induced inflammatory marker expression and matrix degradation in NP cells• Exosome attenuate TXNIP-NLRP3 inflammasome activation and caspase-1 cleavage induced by H₂O₂• Exosomes replenish mitochondrial-related proteins and attenuate mitochondrial dysfunction	<ul style="list-style-type: none">• Exosomes attenuate the progression of IVDD• Exosomes delay matrix degradation during the progression of IVDD	H ₂ O ₂ 500 × 10 M ⁻⁶
----------------------------	--------	-------	-----------	----------------------	---------------------	--	---	---	--

(Continues)

TABLE 36 (Continued)

Study	Cell type	Species of cell source	Size of EV	Isolation	Model	Modeling method	In vitro appraisement	In vivo appraisement	Additional manipulation
Bari et al. ¹¹⁸⁶	ASCs	Human	171.8 ± 18.3 nm	Ultrafiltration	In vitro	-	<ul style="list-style-type: none">At concentrations between 5 and 50 mg/mL, freeze-dried secretome showed in vitro counteract the oxidative stress damage induced by H₂O₂ on nucleus pulposus cellsFreeze-dried secretome became cytotoxic to NPCs at a concentration of over 50 mg mL⁻¹	-	H ₂ O ₂ (1 × 10 ⁻³)
Xie et al. ¹¹⁸⁷	MSCs	Rat	30–200 nm	Ultra-centrifugation	In vitro In vivo	The IVDD SD rat model was established by needle puncture	<ul style="list-style-type: none">Exosomes reduced apoptosis and calcification of EPCs induced by TBHPThe downregulated level of miR-31-5p in exosomes impaired exosomal protective effects on EPCmiR-31-5p negatively regulated ATF6-related ER stress and inhibited apoptosis and calcification in EPCs	<ul style="list-style-type: none">Sub-endorplate injection of MSC exosomes can ameliorate IVDDDownregulation of miR-31-5p from Exosomes inhibited exosomal protective effects on EPC	TBHP (20 × 10 ⁻⁶ to 60 × 10 ⁻⁶ m)
Liao et al. ¹¹⁶³	BMMSCs	Human	Average 94.3 nm	ExoQuick reagent	In vitro In vivo	Injection of AGEs into in SD rat tail to induce IVD degeneration	<ul style="list-style-type: none">MSC-exos reduced AGEs-induced ER stress and ameliorated NP cells apoptosisMSC-exos inhibited the activation of UPR under AGEs-induced ER stress; and decreased CHOP expressionMSC-exos attenuate ER stress-induced apoptosis by activating AKT and ERK signaling in human NPCs	<ul style="list-style-type: none">MSC-exos modulated ER stress-related apoptosis and retarded IDD progression in a rat tail model	AGEs (200 µg mL ⁻¹)

Note: Reprinted with permission from Lu et al.¹¹⁷²

Abbreviations: AFC, annulus fibrosus cell; AGEs, Advanced glycation end products; ASC, adipose-derived mesenchymal stromal cell; BMSC, bone marrow-derived mesenchymal stem cell; CEP, cartilage endplate chondrocyte; CHOP, C/E homologous protein; EPCs, endplate chondrocytes; ER, endoplasmic reticulum; MSC, mesenchymal stem cell; NPC, nucleus pulposus cell; PLMSC, placental mesenchymal stem cell; TBHP, tert-butyl hydroperoxide; UC-MSC, umbilical cord-derived mesenchymal stem cell; UPR, unfolded protein response.

medicine has been identified for COVID-19, with the exception of three drugs: camostat mesylate, favipiravir, and remdesivir, which acts as an agent against the Ebola virus.¹²⁰¹ It has been shown that SARS-CoV-2 infection-induced a cytokine storm that may cause ARDS and eventually multiple crucial organ failures. About 80% of COVID-19-affected individuals have minor symptoms that are restricted to the upper respiratory tract, yet in 20% of cases, serious illness results.¹²⁰² There have been dozens of clinical studies focused on the use of exosome therapy for the treatment of COVID-19 in the past 3 years alone, as well as a growing number for the management and mitigation of "long-COVID."¹²⁰³⁻¹²²²

Despite the fact that there are currently a number of vaccines available to immunize people, scientists worldwide are constantly researching potential treatment approaches to treat infected populations due to the daily discovery of new information regarding the structure, pathogenicity, transmission mechanism, and immunological characteristics of the SARS-CoV-2 virus.¹²⁰³⁻¹²²²

7.1 | Treatment of COVID-19 using exosomes

As mentioned previously, there is no specific/effective vaccine/therapeutic option for combating the negative effects/side effects of COVID-19 such as the cytokine storm. Previous studies have showed the potential of exosomes for treating SARS coronavirus infection.¹²⁰³⁻¹²²² In a study by Kuate et al.,¹²²³ it was demonstrated that exosomes carrying SARS coronavirus spike S protein was successful in inducing neutralizing antibody titers. These results indicated that exosomes can be utilized for treating SARS coronavirus infection.¹²²³

A hot topic in the COVID-19 research world is using MSCs in treating the disease. Particularly, exosomes derived from these cells have attracted even more attention because they are much safer. Various studies have investigated the efficiency and safety of using exosomes derived from MSCs in treating patients with COVID-19. For example, Vikram et al. tested the therapeutic potential of bone marrow MSC-Exos in 24 patients infected with SARS-CoV-2 and moderate to severe ARDS.¹²²⁴ Introduction of exosomes to patients was shown to be safe and led to significant improvement in the clinical status and oxygenation. A recent clinical trial has investigated the efficacy of inhalation MSC-Exos in alleviating the post-infection symptoms of COVID-19 (NCT04276987).

It has further been shown that MSC-Exos have the ability to promote the survival of alveolar macrophages and change their phenotype from pro-inflammatory (M1) into the anti-inflammatory (M2) polarization.^{1225,1226} The implication of these findings is that exosomes can be a viable alternative to their parental cells. Different studies have found that MSC-Exos in the inflammatory environment of ARDS were responsible for reprogramming macrophages from M1-M2 polarization and also improving their phagocytosis effects and oxidative phosphorylation.¹²²⁷⁻¹²²⁹

In totality, various studies have generally concluded the following four main advantages with using exosomes for the management and treatment of patients with COVID-19:

1. Protection and proliferation of lung epithelial cells

Preclinical research conducted by many groups has shown that exosomes have a protective function. In one research article, it was shown that lung epithelial cells were protected from oxidative stress-induced cell death by miR-21-5p administration via MSC-Exos.¹²³⁰ It was also shown by a different research group that the surface of MSC-Exos expressed Alpha-1-antitrypsin (AAT). Due to its strong inhibitory properties against neutrophil-derived proteolytic enzymes, the anti-inflammatory and immunomodulatory properties of AAT plays a vital role in preserving lung epithelial cells.¹²³¹

2. Reversal of lung inflammation

The pro-inflammatory response may be a key factor in the pathophysiology of CoVs, according to data from very ill patients. Release of pro-inflammatory cytokines and chemokines is seen in macrophages, dendritic cells, and respiratory epithelial cells during the early stages of CoV infection.^{1232,1233} However, throughout the subsequent phases, these cells exhibit diminished secretion of antiviral factors, namely interferons (IFNs), while displaying elevated levels of pro-inflammatory cytokines, such as interleukin-1 β (IL-1 β) and IL-6, as well as chemokines including CCL-2, CCL-3, and CCL5. The delayed but heightened presence of pro-inflammatory substances leads to the occurrence of a "cytokine storm," which has the potential to cause organ damage. Consequently, this phenomenon represents a significant issue that contributes to the severity of the illness.¹²³⁴ Immunosuppressive drugs are given to these individuals, which is unavoidably linked to a higher risk of infection.¹²³⁵ Due to their established ability to modulate the immune system, MSCs and their exosomes that are derived from them have been studied in a number of preclinical and clinical contexts.

An interestingly comparable state is seen in lung illness linked with graft-versus-host disease (GvHD).^{1236,1237} MSC-Exos have been proven in an early clinical trial to reduce grade IV GvHD patients' symptoms.⁷⁴⁰ Despite the fact that there is little information on exosomes' powerful immunomodulatory function in a therapeutic setting. This first investigation strengthens the possibility that exosomes might help treat lung damage caused by "cytokine storms."

3. Polarization of lung macrophages

As previously reported, a cytokine storm is produced during a viral infection as a result of an inadequate immune response, which exacerbates lung damage. The lungs' pro-inflammatory macrophages play a major role in mediating this.¹²³⁸ Exosome-derived therapy may be a novel way to treat nCOV-associated pathogenicity, according to some preclinical research assessing the impact of MSC-Exos on lung macrophages in diverse lung injury models. These investigations have shown that exosome-derived proteins and a number of miRNAs, including miR-145, facilitate

the regeneration and repair of lung tissue.¹³⁰ MSC-Exos may potentially influence the phenotype and function of DCs that infiltrate the lung by promoting the production of immunosuppressive cytokines such as TGF- β and IL-10, which shields the lungs from the systemic immune response that is initiated by DCs and detrimental to macrophages.¹²³⁹

4. Reduction in pulmonary edema and lung protein permeability

Increased protein permeability and alveolar inundation, which culminate in pulmonary edema, are results of infection-induced perturbation of the lung endothelial and epithelial barrier.¹²⁴⁰ Consequently, this impairs the ability of the lungs to exchange air. In a recent study, the authors evaluated the impact of systemically delivered MSC-Exos in a mouse model of acute lung damage caused by *Escherichia coli* endotoxins.⁷⁷⁵ These exosomes were shown to lower lung water extravascularly by 43% while also lowering lung protein permeability and pulmonary edema.⁷⁷⁵ Later, using human donor lungs that were unsuitable for transplantation, the team showed that these exosomes could restore alveolar fluid clearance (AFC) in an ex vivo lung perfusion model. The capacity of MSC-Exos to internalize into injured host cells was somewhat facilitated by their CD-44-dependent mechanism.¹²⁴¹

Therefore, the injection of MSC-Exos has a great potential to repair the patient's damaged lungs via a variety of pathways, and as

such, it may be a viable therapeutic nanomedicine intervention for patients in critical condition (Figure 37).

8 | CANCER THERAPY

There is no question that the use of exosomes in the field of cancer has overwhelmingly been utilized as early detection biomarkers.¹²⁴³ Since exosomes are more stable than proteins and hormones in the blood, many research groups have favored using exosomes as early biomarkers for cancer detection and disease progression.¹²⁴³ Interestingly, since exosomes can also carry signaling molecules, they have more recently been utilized as therapeutic options. The use of exosomes as therapeutic regenerative agents is the main topic of this section. Exosomes may also be effectively used as drug delivery vehicles to treat a variety of malignancies.

8.1 | Breast cancer

In a paper titled: "Exosomes as Emerging Drug Delivery and Diagnostic Modality for Breast Cancer: Recent Advances in Isolation and Application," Kumar and colleagues shed light on the potential use of exosomes for the diagnosis and treatment of breast cancer.¹²⁴⁴ Lung cancer is the most common cancer worldwide, with breast cancer coming in second.¹²⁴⁴




FIGURE 37 Schematic representation showing the potential role of MSCs-derived exosomes in combating COVID-19 Infection. Panel (A) synergistic effect of the drug and exosomes may be utilized as an effective approach against the virus. Various hydrophobic and hydrophilic drugs with antiviral properties can be packaged into exosomes for its delivery to the target site. Panel (B) The therapeutic cargo present in Exosomes aids in the reduction in inflammation, cellular repair, alveolar fluid clearance, and other damage caused to the lung during viral infection. Reprinted with permission Pinky et al.¹²⁴²

Chemotherapy, immunotherapy, radiation, and surgery are all used to treat breast cancer. Chemotherapy has been the mainstay of cancer treatment among these medicines; nonetheless, it comes with a number of adverse effects, including nephrotoxicity, hair loss, vomiting, and neurotoxicity. Novel medication delivery techniques that avoid the mentioned negative effects and deliver the right pharmaceuticals to the intended spot have emerged as a result of these disadvantages. Exosomes are thought to be a newly developed, innovative drug delivery mechanism that offers a flexible platform for both medication administration and diagnostics.

Exosomes are being used in clinical studies to treat breast cancer. These trials are based on several reports that show enormous scientific effort that produced favorable results.¹²⁴⁴ The present human exosome experiments are mostly focused on discovering possible exosomal biomarkers in various bodily fluids (particularly blood and urine) for early cancer diagnosis, even if the notion remains far from practical reality. Proteomic analysis of 72 breast cancer patients' exosomes obtained from the cerebral fluid is part of an ongoing clinical investigation on exosomes used to diagnose leptomeningeal metastatic breast cancer.¹²⁴⁴ In a related clinical investigation; exosomes were extracted from breast cancer patient's blood and urine samples in order to measure the stress protein (HSP 70), which is thought to be a tumor biomarker.¹²⁴⁴ Exosomes are being considered for use in chemotherapy in addition to their use as biomarkers.¹²⁴⁴ To measure the expressed HER2 and HER3 dimers over the isolated exosomes, for example, has allowed scientists to identify HER2 type breast cancer. This is because the HER2 receptors are also discovered to be overexpressed on the surface of exosomes. Exosome loading may also be used to determine the impact of pembrolizumab on the tumor microenvironment in a similar manner.¹²⁴⁴

While this field is still in its infancy, there is tremendous upside currently for the use of exosomes in breast cancer therapy. Future years of research should lead to novel discoveries and will likely garnish mainstream momentum in the coming decade.

8.2 | Colorectal cancer

Globally, colorectal cancer (CRC) is the second most common cause of cancer-related deaths. It is a cancer that emerging from the colon or rectum.¹²⁴⁵ Of all CRCs, 72% are caused by colon cancer and 28% by rectal cancer.¹²⁴⁶ An adenocarcinoma originating from the colon's and/or rectum's glandular epithelial cells accounts for about 90% of colorectal cancer cases. The greater death rate associated with CRC is mostly caused by its advanced stage of diagnosis. 64% of CRC patients survive for 5 years, compared to only 12% of those with metastases.¹²⁴⁷

In a study titled: "The therapeutic potential of stem cell-derived exosomes in the ulcerative colitis and colorectal cancer," Guo et al.¹²⁴⁸ discovered that exosomes produced from different MSC sources, such as olfactory ecto-MSCs (OE-MSCs), human umbilical cord-derived MSCs (hUC-MSCs), human ADSCs, and human BMSCs, demonstrated a protective function against ulcerative colitis (UC) and colorectal cancer (CRC). It has been discovered that exosomes from OE-MSCs, ADSCs, hBMSCs, and hUC-MSCs, can improve experimental UC by suppressing inflammatory cells such as macrophages and Th1/Th17 cells, lowering the expression of pro-inflammatory cytokines, and stimulating Treg and Th2 cells' anti-inflammatory function and increasing the expression of anti-inflammatory cytokines. Furthermore, it has been shown that tumor-suppressive miRs (miR-3940-5p/miR-22-3p/miR-16-5p) included in hUC-MSC-Exo and hBMSC-Exo inhibit the growth, migration, and invasion of colorectal cancer cells via controlling the RAP2B/PI3K/AKT signaling pathway and ITGA2/ITGA6.¹²⁴⁸

Consequently, and to summarize, MSC-Exos have positive effects on UC and CRC by means of two distinct processes, namely immune response modulation and antitumor response induction, respectively. Table 37 highlights these interactions and the advantages of exosomes over their parent MSCs.

TABLE 37 Major differences between MSC and MSC-derived exosome in colorectal cancer.

	MSC	MSC-derived exosome
Therapeutic effects	Cancer immunotherapy, regeneration medicine, and immunomodulation	Cancer immunotherapy, regeneration medicine, and immunomodulation
Drug and nucleic acid delivery	Only limited drugs could be internalized by MSC, such as paclitaxel and gemcitabine, transfection efficiency is a major limitation for nucleic acid delivery	Promising carriers for all type of drugs, also for nucleic acid with increased efficiency compared to MSC
Target tissue	Injured site	Injured site
Immunogenicity	Can be allogenic for the immune system	Non-immunogenic
Clinical application in CRC	Preclinical and clinical application	Currently in preclinical applications
Production	Undergo senescence after only a few passages, expensive to have large-scale production	No senescence and easy to generate a large-scale production for clinic application

Note: Reprinted with permission from Guo and colleagues.¹²⁴⁸

8.3 | Gastric cancer

The prognosis for gastric cancer (GC), a frequent malignant tumor that affects human health, is poor, and it often has an obscure origin. It is the fourth most frequent cancer overall and the third leading cause of cancer-related deaths globally. Approximately 1.089 million new cases and 768 000 deaths of GC occur each year.¹²⁴⁹ Interestingly, various research groups have found that exosomes play an important role in the invasion and metastasis of gastric cancer and can be, therefore, used as very early detection tool for diagnosis.¹²⁵⁰ Age-standardized five-year survival rates for GC are estimated to be between 20 and 40 percent because of delayed diagnosis, a dismal prognosis, and ineffective treatment.^{1251,1252} In a paper titled: "Exosomes and Exosomal circRNAs: The Rising Stars in the Progression, Diagnosis and Prognosis of Gastric Cancer,"¹²⁵⁰ Lu and colleagues discussed the potential of exosomes as new biomarkers in the field and thoroughly point to their potential as future applications as mainly diagnosis markers but with the ability to be utilized as therapeutic options once their implication and cargo are fully understood.

8.4 | Osteosarcoma

The most frequent bone tumor that affects children and teenagers is osteosarcomas. In a study titled: "Exosomes as Efficient Nanocarriers in Osteosarcoma: Biological Functions and Potential Clinical Applications," Yang et al.¹²⁵³ overviewed the state of exosome research in the field of osteosarcomas, emphasizing both the biological roles of osteosarcoma exosomes and their use as therapeutic targets and diagnostic indicators in the disease.

Exosomes have significant roles in osteosarcomas, indicating that they may be explored as therapeutic targets as well. The osteosarcoma derived exosomal biomarkers mentioned above could also be utilized as therapeutic targets of osteosarcomas. According to Baglio et al., functional TGF- β molecules are present in the exosomes secreted by osteosarcomas, and these molecules enhance IL-6 expression, thereby promoting osteosarcoma growth and metastasis formation. Combining TGF- β inhibitors with IL-6 blocking agents has been shown to halt osteosarcoma progression while lowering drug resistance.¹²⁵⁴ Notaro et al. have recently shown that the synthetic agonist of cannabinoid receptors, WIN, significantly increased the number of exosomes released. Additionally, isolated exosomes from WIN-treated cells had strong anti-migratory effects on osteosarcoma cells that were not receiving treatment, suggesting that WIN-treated cells may provide a novel therapeutic agent for osteosarcoma therapy.¹²⁵⁵

Additionally, exosome-derived RNAs may be used as osteosarcoma therapeutic targets, according to recent research. Zhang et al.¹²⁵⁶ showed that miR-206 generated from BMSC exosomes might enter osteosarcoma cells and halt the growth of the tumor by focusing on TRA2B. Exosome-derived miR-101 has been shown in research by Zhang et al.¹²⁵⁷ to have metastasis-inhibitory

characteristics in osteosarcomas. Ye et al.¹²⁵⁸ found that patients with osteosarcomas have many dysregulated exosome-derived miRNAs. Wang et al.¹²⁵⁹ demonstrated that the exosome-derived miR-1228 might promote osteosarcoma invasion and migration by downregulating the mRNA expression of SCA1 in osteosarcomas. These studies, among others, highlight the potential for novel therapeutic targets for osteosarcomas with the use of exosomes. Furthermore, an additional review article focuses on the unique structure and relevant characteristics of exosomes as promising nanocarriers for osteosarcoma treatment.¹²⁶⁰

9 | OTHER THERAPEUTIC USES OF EXOSOMES

Lastly, there have been a number of other uses of exosomes for various illnesses that fall into random categories. Within this section, the therapeutic use of exosomes will be briefly mentioned regarding its use for the management of infertility, obesity, and sleep apnea.

9.1 | Infertility

In a study titled: "Mesenchymal Stem-Cell Derived Exosome Therapy as a Potential Future Approach for Treatment of Male Infertility Caused by Chlamydia Infection," Izadi et al.¹²⁶¹ studied the use of exosomes on male infertility following infection. Unfortunately, male fertility, sperm function, and the reproductive tract are all negatively impacted by microbiological sexually transmitted infections (STIs). Since STIs often have no symptoms and may lead to serious side effects, including fibrosis, scarring, and urogenital inflammation, optimal treatments should be performed to prevent the noxious effect of STIs on male fertility.¹²⁶¹ Chlamydia trachomatis stands out as the prevailing bacterial sexually transmitted infection (STI) that often presents without symptoms, and that may have an effect on male fertility and sperm quality. The therapeutic potential of exosomes derived from MSCs has attracted increasing attention in recent years. These exosomes are of interest because they may modulate immunity, reduce inflammation, neutralize free radicals, and promote tissue repair. Importantly, they provide these benefits without the potential drawbacks associated with traditional stem cell transplantation-based treatments. Exosome therapy, being a noninvasive modality, has shown encouraging outcomes in terms of its potential to facilitate the regeneration of impaired sperm and the treatment of asthenozoospermia.¹²⁶¹

Exosomes have shown promise in restoring spermatogenesis and sperm regeneration in animal studies. One study found that exosomes derived from amniotic fluid could restore sperm parameters like motility and concentration, as well as the number of spermatocytes and spermatogonia, and ultimately male fertility.¹²⁶² Exosomes have also been shown to increase post-thaw sperm parameters and have a protective impact against oxidative stress caused by the cryopreservation procedure and sperm cryoinjuries

(such as cell membrane damage and DNA damage).^{1263,1264} It is interesting to note that treating spermatozoa with MSC-Exos might enhance sperm sticky and fusogenic qualities by shuttling adhesion molecules, including CD44, CD29, CD54, and CD106, in addition to increase sperm characteristics after frozen-thawed sperm.¹²⁶⁵ Furthermore, a number of clinical experiments (NCT01159288, NCT03608631, and NCT01294072) have shown that exosomes may be loaded with medications or bioactive molecules for therapeutic purposes (NCT04276987, NCT03437759, NCT04213248, and NCT04602442).¹²⁶⁶

In conclusion, exosomes have been shown to have a significant impact in mitigating the effects of infection, including inflammation, cell damage, fibrogenesis inhibition, and scar formation.

9.2 | Obesity

Obesity is distinguished by the presence of low-grade, persistent inflammation, which contributes to the development of insulin resistance and diabetes.¹²⁶⁷ The presence of obesity has been associated with the onset and advancement of certain autoimmune disorders, such as thyroid autoimmunity, inflammatory bowel disease, psoriasis, psoriatic arthritis, and rheumatoid arthritis. The aforementioned issue poses a significant risk to public health, exceeding the magnitude of both the ongoing opioid crisis and the prevalence of cancer. In addition to the deposition of excess adipose tissue and a reduction in the body's resting metabolic rate, obesity is also associated with an elevated prevalence of type-2 diabetes mellitus, hypertension, atherosclerosis, hyperlipidemia, and cardiovascular disease.^{1268,1269}

By decreasing regulatory T cells (Tregs), raising Th17 and Th1 immune responses, and producing inflammatory cytokines, these illnesses are caused by a change in self-tolerance that promotes a pro-inflammatory immune response.¹²⁶⁷ Therefore, in order to create treatments that lower the risk of autoimmune disorders and other immunological issues, it becomes imperative to understand the immunological alterations that result in this low-grade inflammatory milieu. Patients that are obese and have a high BMI release a large amount of microvesicles and exosomes from their adipocytes and immune cells.¹²⁶⁷ While presently much research is focused on understanding these secreted exosomes during pathogenesis,¹²⁶⁷ eventually, exosomes could be engineered to lower inflammation and potentially reverse some of the compounding negative effects of obesity on overall health.

9.3 | Sleep apnea

In the review article titled: "The Mystery of Red Blood Cells Extracellular Vesicles in Sleep Apnea with Metabolic Dysfunction," the role of red blood cell exosomes was revealed as a contributing factor to sleep apnea and overall body inflammation.¹²⁷⁰ Humans sleep for around one-third of their lives, making it an essential

component of living. It is obvious that inadequate sleep, particularly in the form of psychiatric problems like depression or stress, may lead to poor mental performance, daytime lethargy, and decreased attentiveness.¹²⁷¹ Both genders are affected by sleep-disordered breathing (SDB), which is often linked to a broad range of co-morbid illnesses in many organ systems.¹²⁷²

OSA has become a significant public health issue, and mounting data indicates that untreated OSA may contribute to the onset of a number of illnesses, including neurological disorders as well as failing organs.^{1267,1270} Moreover, OSA may cause blood oxygenation to drop and the sleep cycle to become fragmented. Free radicals, or ROS, have the capacity to generate and combine with NO to create peroxynitrite, which reduces NO's bioavailability. The characteristic of OSA, hypoxia, is a decrease in tissue oxygen saturation that impacts several cell types, with cell-to-cell communication being essential to the result of this interaction. Although mostly known for their function as oxygen and nutrition carriers to tissues, red blood cells (RBCs) also play important roles in viscosity, blood rheology, redox regulation, and systemic NO metabolism control. It has been shown that RBCs enhance cardiac damage and cause endothelial dysfunction.¹²⁷⁰ Exosomes, which are released by RBCs in both healthy and pathological circumstances, may be essential for the identification of hypoxic situations and for facilitating their restoration. The molecular connections between alterations in RBC functional characteristics and cardiovascular disease are extensive. Thus, increasing our understanding of the pathophysiological significance of RBC-EVs and their methods of generation may aid in clarifying the nature of circulating EVs and facilitating their use in therapeutic settings.

10 | DISCUSSION AND CONCLUSIONS

The present scoping review highlights the many applications of exosomes across many fields of medicine. Exosomes are among key paracrine effectors secreted by MSCs and due to their biological cargo, which is similar to parental cells, and their ability to preserve healing properties, they are considered as an attractive candidate to replace MSCs in treating various diseases and affecting many organs (Figure 38).¹²⁷³ Nontoxicity, low immunogenicity, high stability, easy storage, and the potential to be produced as an off-the-shelf product are several benefits of exosomes compared to their cellular counterparts that led to their expanding clinical application as new therapeutic surrogates.¹²⁷⁴

Recent research conducted by Hosseini et al., the authors emphasized five significant benefits of exosomes compared to their parent cells. These included¹²⁰⁵:

- The analysis of exosomes' activity and their originating cell demonstrates that exosomes may have the same effect by either a distinct or comparable mechanism, and in several instances, they are far more potent.¹²⁷⁵ Del Fattore et al. demonstrated that MSC-Exos had superior immunomodulatory capabilities compared with the parent cells. It was shown that MSC-Exos, which




FIGURE 38 Biological mechanisms of MSC exosomes. (A) Immunomodulatory effects of MSC exosomes. (B) Reactions of MSC exosomes in response to oxidative stress. (C) Interactions between tumor cells and MSC exosomes. (D) Applications of MSC exosomes in regenerative medicine. CCL, C-C motif ligand; CXC, C-X-C motif chemokine; DCs, dendritic cells; I/R, ischemia/reperfusion; IFN- γ , interferon γ ; IL, interleukin; NK cells, natural killer cells; ROS, reactive oxygen species; TGF- β , transforming growth factor beta; Tregs, regulatory T cells. Reprinted with permission from Shen and Chen.⁷⁴⁷

likely have different characteristics than parent cells, modify the ratio of Treg/Teff cells in P-MSCs and raise the levels of anti-inflammatory cytokines, such as IL-10.¹²⁷⁶

- Exosomes, because of their diminutive size, may easily traverse narrow blood vessels and even the blood-brain barrier. In contrast, MSCs are prone to being lodged in the lungs and causing pulmonary embolism and infarction, particularly after intravenous administration.¹²⁷⁷ Because of their enormous size (and 25 mL in suspension), inosine-labeled MSCs showed a significant initial uptake when injected intravenously and trapped a substantial number of cells in the lungs.¹²⁷⁸
- Exosomes generated from MSCs, whether autologous or allogeneic, pose no threat since they do not include MHC class I or II on their surface. A recent prospective non-randomized cohort trial was conducted to assess the safety and therapeutic effectiveness of exosomes produced from allogeneic BMSCs in COVID-19 patients.¹²²⁴ All safety objectives were satisfied after 72h of a single intravenous injection of exosomes, and no adverse effects were noted. While 24 individuals were observed, 17 (or 71% of the total) made a full recovery. The ability to restore patients' oxygenation further improved under these simple clinical settings. An increase in lymphocyte and neutrophil counts, as well as a

decrease in acute phase reactants, C-reactive protein, and ferritin, were all indicators of immune reconstitution in the laboratory results. Along with that discovery, scientists also announced that exosomes produced from MSCs are a promising new treatment option for severe COVID-19.¹²²⁴ As mentioned before, there is still cause for worry about MSCs' aberrant differentiation and spontaneous transformation.

- Exosomes are harvested much simpler and much more affordable to produce and freeze-store grade-clinical exosomes in accordance with GMP standards.¹²⁷⁹ As an example, quality control for exosome manufacture might begin with the donor competence analysis and be used throughout the whole process. Another example of a closed-loop system would be the possibility of completely automating some manufacturing operations, such as final concentration and vial filling. Every batch of exosomes is tested for size, homogeneity, number, and positive and negative indicators in the final product.¹²⁸⁰
- Effortless manipulation of exosomes and the ability to design them enables the creation of a diverse array of products customized for specific purposes using a variety of direct and indirect methods. Enhancing the therapeutic efficacy of exosomes may be achieved by including medicinal medicines within the exosomes




FIGURE 39 Stem cell extracellular vesicles (EVs) for clinical applications. EVs are designed, manufactured, and quality controlled beforehand and stored as off-the-shelf medications to be infused in patients as needed. Other EVs may also be customized accordingly for the individual patient by harvesting autologous stem cells, expanding, and modifying them, producing EVs, and infusing them back to the same patient. In this process, the stem cells can also be genetically modified, and their EVs may undergo further modification by loading them with therapeutic molecules. Finally, EVs need to undergo quality control and be stored for future administration. Reprinted with permission from Riazifar et al.¹²⁸²

are deriving them from various sources depending on the clinical indications.^{129,1281}

To date, over 600 clinical trials (registered, ongoing, and completed) worldwide using EVs in diagnostic and therapeutic capacities will pave way to their future use in standard clinical practice. Many open questions remain including the exact signaling molecules required in various exosomes for the treatment of various illnesses. For instance, exosome content may be vastly different in signaling molecules to treat Parkinson's disease versus osteoarthritis of the knee. While these open questions remain, positive outcomes from various MSC-sourced exosomes have shown benefit across practically all fields of medicine. Figure 39 depicts the future clinical challenges toward commercializing and utilizing exosomes in every day clinical practice. The number of studies (over 5000 publications yearly) and human clinical trials will certainly pave way toward a tremendous future of clinical practice using exosomes.

ACKNOWLEDGMENT

Open access funding provided by Universitat Bern.

CONFLICT OF INTEREST STATEMENT

None.

DATA AVAILABILITY STATEMENT

Data sharing is not applicable to this article as no data sets were generated or analyzed during the current study.

ORCID

Yufeng Zhang  <https://orcid.org/0000-0001-8702-5291>

REFERENCES

1. Ghosh A, Davey M, Chute IC, et al. Rapid isolation of extracellular vesicles from cell culture and biological fluids using a synthetic peptide with specific affinity for heat shock proteins. *PLoS One*. 2014;9:e110443.
2. Tan A, Rajadas J, Seifalian AM. Exosomes as nano-theranostic delivery platforms for gene therapy. *Adv Drug Deliv Rev*. 2013;65:357-367.
3. Chen J-j, Zhao B, Zhao J, Li S. Potential roles of exosomal microRNAs as diagnostic biomarkers and therapeutic application in Alzheimer's disease. *Neural Plast*. 2017;2017:7027380.
4. Ha D, Yang N, Nadihe V. Exosomes as therapeutic drug carriers and delivery vehicles across biological membranes: current perspectives and future challenges. *Acta Pharm Sin B*. 2016;6:287-296.

5. Zhuang X, Xiang X, Grizzle W, et al. Treatment of brain inflammatory diseases by delivering exosome encapsulated anti-inflammatory drugs from the nasal region to the brain. *Mol Ther*. 2011;19:1769-1779.
6. Fayazi N, Sheykhasan M, Soleimani Asl S, Najafi R. Stem cell-derived exosomes: a new strategy of neurodegenerative disease treatment. *Mol Neurobiol*. 2021;58:3494-3514.
7. Andreone BJ, Larhammar M, Lewcock JW. Cell death and neurodegeneration. *Cold Spring Harb Perspect Biol*. 2020;12:a036434.
8. Hou Y, Dan X, Babbar M, et al. Ageing as a risk factor for neurodegenerative disease. *Nat Rev Neurol*. 2019;15:565-581.
9. Agosta F, Galantucci S, Filippi M. Advanced magnetic resonance imaging of neurodegenerative diseases. *Neurol Sci*. 2017;38:41-51.
10. Erkinen MG, Kim M-O, Geschwind MD. Clinical neurology and epidemiology of the major neurodegenerative diseases. *Cold Spring Harb Perspect Biol*. 2018;10:a033118.
11. Chen X, Pan W. The treatment strategies for neurodegenerative diseases by integrative medicine. *Integr Med Int*. 2015;1:223-225.
12. Huang M, Gu X, Gao X. Nanotherapeutic strategies for the treatment of neurodegenerative diseases. *Brain Targeted Drug Delivery System*. Elsevier; 2019:321-356.
13. Jankovic J. Dopamine depleters in the treatment of hyperkinetic movement disorders. *Expert Opin Pharmacother*. 2016;17:2461-2470.
14. Raschetti R, Albanese E, Vanacore N, Maggini M. Cholinesterase inhibitors in mild cognitive impairment: a systematic review of randomised trials. *PLoS Med*. 2007;4:e338.
15. Chaudhuri KR, Schapira AH. Non-motor symptoms of Parkinson's disease: dopaminergic pathophysiology and treatment. *Lancet Neurol*. 2009;8:464-474.
16. Umemura A, Jaggi JL, Hurtig HI, et al. Deep brain stimulation for movement disorders: morbidity and mortality in 109 patients. *J Neurosurg*. 2003;98:779-784.
17. Cotzias G, Papavasiliou P, Gellene R. Experimental treatment of parkinsonism with L-Dopa. *Neurology*. 1968;18:276-277.
18. Yahr M, Duvoisin R, Hoehn M, Scheer M, Barrett R. L-Dopa (L-3,4-dihydroxyphenylamine)—its clinical effects in parkinsonism. *Trans Am Neurol Assoc*. 1968;93:56-63.
19. Comfort A. Cholinesterase inhibition in treatment of Alzheimer's dementia. *Lancet*. 1978;311:659-660.
20. Rabins PV, Lyketsos CG. Cholinesterase inhibitors and memantine have a role in the treatment of Alzheimer's disease. *Nat Clin Pract Neurol*. 2006;2:578-579.
21. Marks WJ, Ostrem JL, Verhagen L, et al. Safety and tolerability of intraputamenal delivery of CERE-120 (adeno-associated virus serotype 2-neurturin) to patients with idiopathic Parkinson's disease: an open-label, phase I trial. *Lancet Neurol*. 2008;7:400-408.
22. Stewart WF, Kawa S, Corrada M, Metter EJ. Risk of Alzheimer's disease and duration of NSAID use. *Neurology*. 1997;48:626-632.
23. Traynor B, Bruijn L, Conwit R, et al. Neuroprotective agents for clinical trials in ALS: a systematic assessment. *Neurology*. 2006;67:20-27.
24. Ristori G, Romano S, Visconti A, et al. Riluzole in cerebellar ataxia: a randomized, double-blind, placebo-controlled pilot trial. *Neurology*. 2010;74:839-845.
25. Armstrong MJ, Miyasaki JM. Evidence-based guideline: pharmacologic treatment of chorea in Huntington disease: report of the guideline development subcommittee of the American Academy of Neurology. *Neurology*. 2012;79:597-603.
26. Meldolesi J. News about therapies of Alzheimer's disease: extracellular vesicles from stem cells exhibit advantages compared to other treatments. *Biomedicine*. 2022;10:105.
27. Ryan ST, Hosseini-Beheshti E, Afrose D, et al. Extracellular vesicles from mesenchymal stromal cells for the treatment of inflammation-related conditions. *Int J Mol Sci*. 2021;22:3023.
28. Wang Y, Lai X, Wu D, Liu B, Wang N, Rong L. Umbilical mesenchymal stem cell-derived exosomes facilitate spinal cord functional recovery through the miR-199a-3p/145-5p-mediated NGF/TrkA signaling pathway in rats. *Stem Cell Res Ther*. 2021;12:117.
29. Ma Y, Xu X, Li C, et al. Induced neural progenitor cell-derived extracellular vesicles promote neural progenitor cell survival via extracellular signal-regulated kinase pathway. *CNS Neurosci Ther*. 2021;27:1605-1609.
30. Han M, Cao Y, Xue H, et al. Neuroprotective effect of mesenchymal stromal cell-derived extracellular vesicles against cerebral ischemia-reperfusion-induced neural functional injury: a pivotal role for AMPK and JAK2/STAT3/NF- κ B signaling pathway modulation. *Drug Des Devel Ther*. 2020;14:2865-2876.
31. Otero-Ortega L, Laso-García F, Frutos MC, et al. Low dose of extracellular vesicles identified that promote recovery after ischemic stroke. *Stem Cell Res Ther*. 2020;11:70.
32. Feng Y, Guo M, Zhao H, Han S, Dong Q, Cui M. Mesenchymal-stem-cell-derived extracellular vesicles mitigate trained immunity in the brain. *Front Bioeng Biotechnol*. 2020;8:599058.
33. Jiang D, Gong F, Ge X, et al. Neuron-derived exosomes-transmitted miR-124-3p protect traumatically injured spinal cord by suppressing the activation of neurotoxic microglia and astrocytes. *J Nanobiotechnol*. 2020;18:105.
34. Garcia-Contreras M, Thakor AS. Human adipose tissue-derived mesenchymal stem cells and their extracellular vesicles modulate lipopolysaccharide activated human microglia. *Cell Death Discov*. 2021;7:98.
35. Wang X, Yang G. Bone marrow mesenchymal stem cells-derived exosomes reduce A β deposition and improve cognitive function recovery in mice with Alzheimer's disease by activating sphingosine kinase/sphingosine-1-phosphate signaling pathway. *Cell Biol Int*. 2021;45:775-784.
36. Xiang C, Yang K, Liang Z, et al. Sphingosine-1-phosphate mediates the therapeutic effects of bone marrow mesenchymal stem cell-derived microvesicles on articular cartilage defect. *Transl Res*. 2018;193:42-53.
37. Ma X, Huang M, Zheng M, et al. ADSCs-derived extracellular vesicles alleviate neuronal damage, promote neurogenesis and rescue memory loss in mice with Alzheimer's disease. *J Control Release*. 2020;327:688-702.
38. Wang H, Liu Y, Li J, et al. Tail-vein injection of MSC-derived small extracellular vesicles facilitates the restoration of hippocampal neuronal morphology and function in APP/PS1 mice. *Cell Death Discov*. 2021;7:230.
39. Kaminski N, Köster C, Mouloud Y, et al. Mesenchymal stromal cell-derived extracellular vesicles reduce neuroinflammation, promote neural cell proliferation and improve oligodendrocyte maturation in neonatal hypoxic-ischemic brain injury. *Front Cell Neurosci*. 2020;14:601176.
40. Soliman HM, Ghonaim GA, Gharib SM, et al. Exosomes in Alzheimer's disease: from being pathological players to potential diagnostics and therapeutics. *Int J Mol Sci*. 2021;22:10794.
41. Zhang T, Ma S, Lv J, et al. The emerging role of exosomes in Alzheimer's disease. *Ageing Res Rev*. 2021;68:101321.
42. Apodaca LA, Baddour AAD, Garcia C, et al. Human neural stem cell-derived extracellular vesicles mitigate hallmarks of Alzheimer's disease. *Alzheimers Res Ther*. 2021;13:57.
43. Xun C, Ge L, Tang F, et al. Insight into the proteomic profiling of exosomes secreted by human OM-MSCs reveals a new potential therapy. *Biomed Pharmacother*. 2020;131:110584.
44. Kaur S, Verma H, Dhiman M, et al. Brain exosomes: friend or foe in Alzheimer's disease? *Mol Neurobiol*. 2021;58:6610-6624.
45. Cui G-h, Zhu J, Wang Y-c, Wu J, Liu J-r, Guo H-d. Effects of exosomal miRNAs in the diagnosis and treatment of Alzheimer's disease. *Mech Ageing Dev*. 2021;200:111593.

46. Liu W-L, Lin H-W, Lin M-R, et al. Emerging blood exosome-based biomarkers for preclinical and clinical Alzheimer's disease: a meta-analysis and systematic review. *Neural Regen Res*. 2022;17:2381.
47. Li C-C, Hsu W-F, Wo AM. Exosomes-potential for blood-based marker in Alzheimer's disease. *Acta Neurol Taiwanica*. 2022;31:1-6.
48. Soares Martins T, Trindade D, Vaz M, et al. Diagnostic and therapeutic potential of exosomes in Alzheimer's disease. *J Neurochem*. 2021;156:162-181.
49. Sharma S. Role of extracellular vesicles in Alzheimer's disease: current advances. *Curr Mol Med*. 2022;22:85-97.
50. Dickson DW. Parkinson's disease and parkinsonism: neuropathology. *Cold Spring Harb Perspect Med*. 2012;2:a009258.
51. Esposito E, Di Matteo V, Di Giovanni G. Death in the substantia nigra: a motor tragedy. *Expert Rev Neurother*. 2007;7:677-697.
52. Bourdy R, Sánchez-Catalán M-J, Kaufling J, et al. Control of the nigrostriatal dopamine neuron activity and motor function by the tail of the ventral tegmental area. *Neuropsychopharmacology*. 2014;39:2788-2798.
53. Karlsen K, Larsen JP, Tandberg E, Jørgensen K. Fatigue in patients with Parkinson's disease. *Mov Disord*. 1999;14:237-241.
54. Schrag A, Horsfall L, Walters K, Noyce A, Petersen I. Prediagnostic presentations of Parkinson's disease in primary care: a case-control study. *Lancet Neurol*. 2015;14:57-64.
55. Tan LC. Mood disorders in Parkinson's disease. *Parkinsonism Relat Disord*. 2012;18:S74-S76.
56. Jarmalavičiūtė A, Tunaitis V, Pivoraitė U, Venalis A, Pivoriūnas A. Exosomes from dental pulp stem cells rescue human dopaminergic neurons from 6-hydroxy-dopamine-induced apoptosis. *Cytotherapy*. 2015;17:932-939.
57. Narbutė K, Pilipenko V, Pupure J, et al. Intranasal administration of extracellular vesicles derived from human teeth stem cells improves motor symptoms and normalizes tyrosine hydroxylase expression in the substantia nigra and striatum of the 6-hydroxydopamine-treated rats. *Stem Cells Transl Med*. 2019;8:490-499.
58. Chen Y-R, Lai P-L, Chien Y, et al. Improvement of impaired motor functions by human dental exfoliated deciduous teeth stem cell-derived factors in a rat model of Parkinson's disease. *Int J Mol Sci*. 2020;21:3807.
59. Chen H-X, Liang F-C, Gu P, et al. Exosomes derived from mesenchymal stem cells repair a Parkinson's disease model by inducing autophagy. *Cell Death Dis*. 2020;11:288.
60. Haney MJ, Zhao Y, Harrison EB, et al. Specific transfection of inflamed brain by macrophages: a new therapeutic strategy for neurodegenerative diseases. *PLoS One*. 2013;8:e61852.
61. Upadhyay R, Shetty AK. Extracellular vesicles for the diagnosis and treatment of Parkinson's disease. *Aging Dis*. 2021;12:1438-1450.
62. Yang X, Ma Y, Xie H, et al. Extracellular vesicles in the treatment of Parkinson's disease: a review. *Curr Med Chem*. 2021;28:6375-6394.
63. Zhang P, Rasheed M, Liang J, Wang C, Feng L, Chen Z. Emerging potential of exosomal non-coding RNA in Parkinson's disease: a review. *Front Aging Neurosci*. 2022;14:819836.
64. Li K-L, Huang H-Y, Ren H, Yang X-L. Role of exosomes in the pathogenesis of inflammation in Parkinson's disease. *Neural Regen Res*. 2022;17:1898.
65. Ouerdane Y, Hassaballah MY, Nagah A, et al. Exosomes in Parkinson: revisiting their pathologic role and potential applications. *Pharmaceuticals*. 2022;15:76.
66. Manu MS, Hohjoh H, Yamamura T. Extracellular vesicles as pro- and anti-inflammatory mediators, biomarkers and potential therapeutic agents in multiple sclerosis. *Aging Dis*. 2021;12:1451-1461.
67. Gandhi R, Laroni A, Weiner HL. Role of the innate immune system in the pathogenesis of multiple sclerosis. *J Neuroimmunol*. 2010;221:7-14.
68. El Behi M, Dubucquoi S, Lefranc D, et al. New insights into cell responses involved in experimental autoimmune encephalomyelitis and multiple sclerosis. *Immunol Lett*. 2005;96:11-26.
69. Van Kaer L, Postoak JL, Wang C, Yang G, Wu L. Innate, innate-like and adaptive lymphocytes in the pathogenesis of MS and EAE. *Cell Mol Immunol*. 2019;16:531-539.
70. Ponath G, Park C, Pitt D. The role of astrocytes in multiple sclerosis. *Front Immunol*. 2018;9:217.
71. Traiffort E, Kassoussi A, Zahaf A, Laouarem Y. Astrocytes and microglia as major players of myelin production in normal and pathological conditions. *Front Cell Neurosci*. 2020;14:79.
72. Ludwin SK, Rao VT, Moore CS, Antel JP. Astrocytes in multiple sclerosis. *Mult Scler J*. 2016;22:1114-1124.
73. Lecuyer M-A, Kebir H, Prat A. Glial influences on BBB functions and molecular players in immune cell trafficking. *Biochim Biophys Acta*. 2016;1862:472-482.
74. Laso-García F, Ramos-Cejudo J, Carrillo-Salinas FJ, et al. Therapeutic potential of extracellular vesicles derived from human mesenchymal stem cells in a model of progressive multiple sclerosis. *PLoS One*. 2018;13:e0202590.
75. Riazifar M, Mohammadi MR, Pone EJ, et al. Stem cell-derived exosomes as nanotherapeutics for autoimmune and neurodegenerative disorders. *ACS Nano*. 2019;13:6670-6688.
76. Li Z, Liu F, He X, Yang X, Shan F, Feng J. Exosomes derived from mesenchymal stem cells attenuate inflammation and demyelination of the central nervous system in EAE rats by regulating the polarization of microglia. *Int Immunopharmacol*. 2019;67:268-280.
77. Gomzikova MO, James V, Rizvanov AA. Therapeutic application of mesenchymal stem cells derived extracellular vesicles for immunomodulation. *Front Immunol*. 2019;10:2663.
78. Casella G, Colombo F, Finardi A, et al. Extracellular vesicles containing IL-4 modulate neuroinflammation in a mouse model of multiple sclerosis. *Mol Ther*. 2018;26:2107-2118.
79. Casella G, Rasouli J, Boehm A, et al. Oligodendrocyte-derived extracellular vesicles as antigen-specific therapy for autoimmune neuroinflammation in mice. *Sci Transl Med*. 2020;12:eaba0599.
80. Clark K, Zhang S, Barthe S, et al. Placental mesenchymal stem cell-derived extracellular vesicles promote myelin regeneration in an animal model of multiple sclerosis. *Cells*. 2019;8:1497.
81. Confavreux C, Hutchinson M, Hours MM, Cortinovis-Tourniaire P, Moreau T, Group PiMS. Rate of pregnancy-related relapse in multiple sclerosis. *New Engl J Med*. 1998;339:285-291.
82. Williams JL, Gatson NN, Smith KM, Almad A, McTigue DM, Whitacre CC. Serum exosomes in pregnancy-associated immune modulation and neuroprotection during CNS autoimmunity. *Clin Immunol*. 2013;149:236-243.
83. Silva AK. Exosomes: a novel therapeutic paradigm for treatment of depression. *Curr Drug Targets*. 2020;21. doi:10.2174/1389450121999201006193005
84. Heninger G, Delgado P, Charney D. The revised monoamine theory of depression: a modulatory role for monoamines, based on new findings from monoamine depletion experiments in humans. *Pharmacopsychiatry*. 1996;29:2-11.
85. Al-Harbi KS. Treatment-resistant depression: therapeutic trends, challenges, and future directions. *Patient Prefer Adherence*. 2012;6:369-388.
86. Bhatt S, Radhakrishnan M, Jindal A, Devadoss T, Dhar AK. Neuropharmacological evaluation of a novel 5-HT₃ receptor antagonist (6g) on chronic unpredictable mild stress-induced changes in behavioural and brain oxidative stress parameters in mice. *Indian J Pharmacol*. 2014;46:191-196.
87. Keller J, Gomez R, Williams G, et al. HPA axis in major depression: cortisol, clinical symptomatology and genetic variation predict cognition. *Mol Psychiatry*. 2017;22:527-536.
88. Mot YY, Moses EJ, Mohd Yusoff N, Ling K-H, Yong YK, Tan JJ. Mesenchymal stromal cells-derived exosome and the roles in the treatment of traumatic brain injury. *Cell Mol Neurobiol*. 2023;43:469-489.

89. Zhang Z, Zhang Z, Guan L, Zhang K, Zhang Q, Dai L. A combined procedure to deliver autologous mesenchymal stromal cells to patients with traumatic brain injury. *Cytotherapy*. 2008;10:134-139.
90. Li Y, Yang Y-Y, Ren J-L, Xu F, Chen F-M, Li A. Exosomes secreted by stem cells from human exfoliated deciduous teeth contribute to functional recovery after traumatic brain injury by shifting microglia M1/M2 polarization in rats. *Stem Cell Res Ther*. 2017;8:198.
91. Gupta S, Guleria RS, Szabo YZ. MicroRNAs as biomarker and novel therapeutic target for posttraumatic stress disorder in veterans. *Psychiatry Res*. 2021;305:114252.
92. Jaffee CMS, Meyer KS. A brief overview of traumatic brain injury (TBI) and post-traumatic stress disorder (PTSD) within the department of defense. *Clin Neuropsychol*. 2009;23:1291-1298.
93. Tanev KS, Pentel KZ, Kredlow MA, Charney ME. PTSD and TBI co-morbidity: scope, clinical presentation and treatment options. *Brain Inj*. 2014;28:261-270.
94. Moore BA, Brock MS, Brager A, Collen J, LoPresti M, Mysliwiec V. Posttraumatic stress disorder, traumatic brain injury, sleep, and performance in military personnel. *Sleep Med Clin*. 2020;15:87-100.
95. Howard JT, Sosnov JA, Janak JC, et al. Associations of initial injury severity and posttraumatic stress disorder diagnoses with long-term hypertension risk after combat injury. *Hypertension*. 2018;71:824-832.
96. Dyball D, Evans S, Boos CJ, Stevelink SA, Fear NT. The association between PTSD and cardiovascular disease and its risk factors in male veterans of the Iraq/Afghanistan conflicts: a systematic review. *Int Rev Psychiatry*. 2019;31:34-48.
97. Levine AB, Levine LM, Levine TB. Posttraumatic stress disorder and cardiometabolic disease. *Cardiology*. 2013;127:1-19.
98. Millner AJ, Ursano RJ, Hwang I, et al. Prior mental disorders and lifetime suicidal behaviors among US Army soldiers in the Army study to assess risk and resilience in Servicemembers (Army STARRS). *Suicide Life Threat Behav*. 2019;49:3-22.
99. Toblin RL, Quartana PJ, Riviere LA, Walper KC, Hoge CW. Chronic pain and opioid use in US soldiers after combat deployment. *JAMA Intern Med*. 2014;174:1400-1401.
100. Eraly SA, Nievergelt CM, Maihofer AX, et al. Assessment of plasma C-reactive protein as a biomarker of posttraumatic stress disorder risk. *JAMA Psychiatry*. 2014;71:423-431.
101. Rauch SA, King A, Kim HM, et al. Cortisol awakening response in PTSD treatment: predictor or mechanism of change. *Psychoneuroendocrinology*. 2020;118:104714.
102. Carvalho C, Wrobel S, Meyer C, et al. Gellan gum-based luminal fillers for peripheral nerve regeneration: an in vivo study in the rat sciatic nerve repair model. *Biomater Sci*. 2018;6:1059-1075.
103. Carvalho CR, Oliveira JM, Reis RL. Modern trends for peripheral nerve repair and regeneration: beyond the hollow nerve guidance conduit. *Front Bioeng Biotechnol*. 2019;7:337.
104. Nocera G, Jacob C. Mechanisms of Schwann cell plasticity involved in peripheral nerve repair after injury. *Cell Mol Life Sci*. 2020;77:3977-3989.
105. Wang Y, Zhang Y, Li X, Zhang Q. The progress of biomaterials in peripheral nerve repair and regeneration. *J Neurorestoratol*. 2020;8:252-269.
106. Philips C, Campos F, Roosens A, Sánchez-Quevedo MC, Declercq H, Carriel V. Qualitative and quantitative evaluation of a novel detergent-based method for decellularization of peripheral nerves. *Ann Biomed Eng*. 2018;46:1921-1937.
107. Jia Y, Yang W, Zhang K, et al. Nanofiber arrangement regulates peripheral nerve regeneration through differential modulation of macrophage phenotypes. *Acta Biomater*. 2019;83:291-301.
108. Yu T, Xu Y, Jia X, Ao Q. Intraluminal guiding structure of nerve conduits for peripheral nerve regeneration. *Sci Adv Mater*. 2020;12:56-65.
109. Lin T, Liu S, Chen S, et al. Hydrogel derived from porcine decellularized nerve tissue as a promising biomaterial for repairing peripheral nerve defects. *Acta Biomater*. 2018;73:326-338.
110. Chiono V, Tonda-Turo C. Trends in the design of nerve guidance channels in peripheral nerve tissue engineering. *Prog Neurobiol*. 2015;131:87-104.
111. Bucan V, Vaslaitis D, Peck C-T, Strauß S, Vogt PM, Radtke C. Effect of exosomes from rat adipose-derived mesenchymal stem cells on neurite outgrowth and sciatic nerve regeneration after crush injury. *Mol Neurobiol*. 2019;56:1812-1824.
112. Zhao J, Ding Y, He R, et al. Dose-effect relationship and molecular mechanism by which BMSC-derived exosomes promote peripheral nerve regeneration after crush injury. *Stem Cell Res Ther*. 2020;11:360.
113. Chen J, Ren S, Duschner D, et al. Exosomes from human adipose-derived stem cells promote sciatic nerve regeneration via optimizing Schwann cell function. *J Cell Physiol*. 2019;234:23097-23110.
114. Zhang B, Wu X, Zhang X, et al. Human umbilical cord mesenchymal stem cell exosomes enhance angiogenesis through the Wnt4/ β -catenin pathway. *Stem Cells Transl Med*. 2015;4:513-522.
115. Pascucci L, Alessandri G, Dall'Aglio C, et al. Membrane vesicles mediate pro-angiogenic activity of equine adipose-derived mesenchymal stromal cells. *Vet J*. 2014;202:361-366.
116. Bian S, Zhang L, Duan L, Wang X, Min Y, Yu H. Extracellular vesicles derived from human bone marrow mesenchymal stem cells promote angiogenesis in a rat myocardial infarction model. *J Mol Med*. 2014;92:387-397.
117. Zhang Y, Chopp M, Meng Y, et al. Effect of exosomes derived from multipotential mesenchymal stromal cells on functional recovery and neurovascular plasticity in rats after traumatic brain injury. *J Neurosurg*. 2015;122:856-867.
118. Yu T, Xu Y, Ahmad MA, Javed R, Hagiwara H, Tian X. Exosomes as a promising therapeutic strategy for peripheral nerve injury. *Curr Neuropharmacol*. 2021;19:2141-2151.
119. Okin D, Medzhitov R. Evolution of inflammatory diseases. *Curr Biol*. 2012;22:R733-R740.
120. Suh JH, Joo HS, Hong EB, Lee HJ, Lee JM. Therapeutic application of exosomes in inflammatory diseases. *Int J Mol Sci*. 2021;22:1144.
121. Schett G, Neurath MF. Resolution of chronic inflammatory disease: universal and tissue-specific concepts. *Nat Commun*. 2018;9:3261.
122. Dominguez-Villar M, Hafler DA. Regulatory T cells in autoimmune disease. *Nat Immunol*. 2018;19:665-673.
123. Eccleston C, Cooper TE, Fisher E, Anderson B, Wilkinson NM. Non-steroidal anti-inflammatory drugs (NSAIDs) for chronic non-cancer pain in children and adolescents. *Cochrane Database Syst Rev*. 2017;8(8):CD012537.
124. Totzke J, Gurbani D, Raphemot R, et al. Takinib, a selective TAK1 inhibitor, broadens the therapeutic efficacy of TNF- α inhibition for cancer and autoimmune disease. *Cell Chem Biol*. 2017;24:1029-1039. e1027.
125. Swart JF, Delemarre EM, Van Wijk F, et al. Haematopoietic stem cell transplantation for autoimmune diseases. *Nat Rev Rheumatol*. 2017;13:244-256.
126. Schwarzenbach H, Gahan PB. Exosomes in immune regulation. *Non-coding RNA*. 2021;7:4.
127. Qian X, An N, Ren Y, Yang C, Zhang X, Li L. Immunosuppressive effects of mesenchymal stem cells-derived exosomes. *Stem Cell Rev Rep*. 2021;17:411-427.
128. Ti D, Hao H, Tong C, et al. LPS-preconditioned mesenchymal stromal cells modify macrophage polarization for resolution of chronic inflammation via exosome-shuttled let-7b. *J Transl Med*. 2015;13:308.
129. Wang B, Yao K, Huuskes BM, et al. Mesenchymal stem cells deliver exogenous microRNA-let7c via exosomes to attenuate renal fibrosis. *Mol Ther*. 2016;24:1290-1301.

130. Hao Q, Gudapati V, Monsel A, et al. Mesenchymal stem cell-derived extracellular vesicles decrease lung injury in mice. *J Immunol*. 2019;203:1961-1972.
131. Grabiec AM, Hussell T. The role of airway macrophages in apoptotic cell clearance following acute and chronic lung inflammation. *Seminars in Immunopathology*. Vol. 38. Springer; 2016:409-423.
132. Wu H, Fan H, Shou Z, et al. Extracellular vesicles containing miR-146a attenuate experimental colitis by targeting TRAF6 and IRAK1. *Int Immunopharmacol*. 2019;68:204-212.
133. Zhang R, Zhu Y, Li Y, et al. Human umbilical cord mesenchymal stem cell exosomes alleviate sepsis-associated acute kidney injury via regulating microRNA-146b expression. *Biotechnol Lett*. 2020;42:669-679.
134. Liu Y, Lou G, Li A, et al. AMSC-derived exosomes alleviate lipopolysaccharide/d-galactosamine-induced acute liver failure by miR-17-mediated reduction of TXNIP/NLRP3 inflammasome activation in macrophages. *EBioMedicine*. 2018;36:140-150.
135. Li X, Liu L, Yang J, et al. Exosome derived from human umbilical cord mesenchymal stem cell mediates MiR-181c attenuating burn-induced excessive inflammation. *EBioMedicine*. 2016;8:72-82.
136. Wang X, Gu H, Qin D, et al. Exosomal miR-223 contributes to mesenchymal stem cell-elicited cardioprotection in polymicrobial sepsis. *Sci Rep*. 2015;5:13721.
137. Shao M, Xu Q, Wu Z, et al. Exosomes derived from human umbilical cord mesenchymal stem cells ameliorate IL-6-induced acute liver injury through miR-455-3p. *Stem Cell Res Ther*. 2020;11:37.
138. Liu Y, Zou R, Wang Z, Wen C, Zhang F, Lin F. Exosomal KLF3-AS1 from hMSCs promoted cartilage repair and chondrocyte proliferation in osteoarthritis. *Biochem J*. 2018;475:3629-3638.
139. Xie K, Liu L, Chen J, Liu F. Exosomal miR-1246 derived from human umbilical cord blood mesenchymal stem cells attenuates hepatic ischemia reperfusion injury by modulating T helper 17/regulatory T balance. *IUBMB Life*. 2019;71:2020-2030.
140. Fujii S, Miura Y, Fujishiro A, et al. Graft-versus-host disease amelioration by human bone marrow mesenchymal stromal/stem cell-derived extracellular vesicles is associated with peripheral preservation of naive T cell populations. *Stem Cells*. 2018;36:434-445.
141. Zhang W, Wang Y, Kong Y. Exosomes derived from mesenchymal stem cells modulate miR-126 to ameliorate hyperglycemia-induced retinal inflammation via targeting HMGB1. *Invest Ophthalmol Vis Sci*. 2019;60:294-303.
142. Zhuansun Y, Du Y, Huang F, et al. MSCs exosomal miR-1470 promotes the differentiation of CD4⁺ CD25⁺ FOXP3⁺ Tregs in asthmatic patients by inducing the expression of P27KIP1. *Int Immunopharmacol*. 2019;77:105981.
143. Katz G, Pobezinsky LA, Jeurling S, Shinzawa M, Van Laethem F, Singer A. T cell receptor stimulation impairs IL-7 receptor signaling by inducing expression of the microRNA miR-17 to target Janus kinase 1. *Sci Signal*. 2014;7:ra83.
144. Xin H, Katakowski M, Wang F, et al. MicroRNA-17-92 cluster in exosomes enhance neuroplasticity and functional recovery after stroke in rats. *Stroke*. 2017;48:747-753.
145. Wei Z, Qiao S, Zhao J, et al. miRNA-181a over-expression in mesenchymal stem cell-derived exosomes influenced inflammatory response after myocardial ischemia-reperfusion injury. *Life Sci*. 2019;232:116632.
146. Ma M, Chen S, Liu Z, et al. miRNA-221 of exosomes originating from bone marrow mesenchymal stem cells promotes oncogenic activity in gastric cancer. *Onco Targets Ther*. 2017;10:4161-4171.
147. Lu F-B, Chen D-Z, Chen L, et al. Attenuation of experimental autoimmune hepatitis in mice with bone mesenchymal stem cell-derived exosomes carrying microRNA-223-3p. *Mol Cells*. 2019;42:906.
148. Hou K, Li G, Zhao J, et al. Bone mesenchymal stem cell derived-exosomal microRNA-29b-3p ameliorates hypoxic-ischemic brain injury by inhibiting apoptosis and promoting angiogenesis through PTEN and Akt signaling pathway. Available at SSRN 3421597; 2019.
149. Mao G, Zhang Z, Hu S, et al. Exosomes derived from miR-92a-3p-overexpressing human mesenchymal stem cells enhance chondrogenesis and suppress cartilage degradation via targeting WNT5A. *Stem Cell Res Ther*. 2018;9:247.
150. Harrell CR, Jankovic MG, Fellabaum C, et al. Molecular mechanisms responsible for anti-inflammatory and immunosuppressive effects of mesenchymal stem cell-derived factors. *Adv Exp Med Biol*. 2019;1084:187-206.
151. Shin H, Ryu HH, Kwon O, Park BS, Jo SJ. Clinical use of conditioned media of adipose tissue-derived stem cells in female pattern hair loss: a retrospective case series study. *Int J Dermatol*. 2015;54:730-735.
152. La Greca A, Solari C, Furmento V, et al. Extracellular vesicles from pluripotent stem cell-derived mesenchymal stem cells acquire a stromal modulatory proteomic pattern during differentiation. *Exp Mol Med*. 2018;50:1-12.
153. Whiteside TL. Exosome and mesenchymal stem cell cross-talk in the tumor microenvironment. *Seminars in Immunology*. Elsevier; 2018.
154. Chen W, Huang Y, Han J, et al. Immunomodulatory effects of mesenchymal stromal cells-derived exosome. *Immunol Res*. 2016;64:831-840.
155. Kilpinen L, Impola U, Sankkila L, et al. Extracellular membrane vesicles from umbilical cord blood-derived MSC protect against ischemic acute kidney injury, a feature that is lost after inflammatory conditioning. *J Extracell Vesicles*. 2013;2:21927.
156. Zhang B, Yin Y, Lai RC, Tan SS, Choo ABH, Lim SK. Mesenchymal stem cells secrete immunologically active exosomes. *Stem Cells Dev*. 2014;23:1233-1244.
157. Lankford KL, Arroyo EJ, Nazimek K, Bryniarski K, Askenase PW, Kocsis JD. Intravenously delivered mesenchymal stem cell-derived exosomes target M2-type macrophages in the injured spinal cord. *PLoS One*. 2018;13:e0190358.
158. Huldani H, Jasim SA, Bokov DO, et al. Application of extracellular vesicles derived from mesenchymal stem cells as potential therapeutic tools in autoimmune and rheumatic diseases. *Int Immunopharmacol*. 2022;106:108634.
159. Shen Z, Huang W, Liu J, Tian J, Wang S, Rui K. Effects of mesenchymal stem cell-derived exosomes on autoimmune diseases. *Front Immunol*. 2021;12:749192.
160. Zou J, Peng H, Liu Y. The roles of exosomes in immunoregulation and autoimmune thyroid diseases. *Front Immunol*. 2021;12:757674.
161. Peng J, Liang Q, Xu Z, et al. Current understanding of exosomal MicroRNAs in glioma immune regulation and therapeutic responses. *Front Immunol*. 2022;12:813747.
162. Zhang X, Xu D, Song Y, Wang T. Research progress in the application of exosomes in immunotherapy. *Front Immunol*. 2022;13:731516.
163. Matheakakis A, Batsali A, Papadaki HA, Pontikoglou CG. Therapeutic implications of mesenchymal stromal cells and their extracellular vesicles in autoimmune diseases: from biology to clinical applications. *Int J Mol Sci*. 2021;22:10132.
164. Martinez-Arroyo O, Ortega A, Forner MJ, Cortes R. Mesenchymal stem cell-derived extracellular vesicles as non-coding RNA therapeutic vehicles in autoimmune diseases. *Pharmaceutics*. 2022;14:733.
165. Wang L-T, Liu K-J, Sytwu H-K, Yen M-L, Yen BL. Advances in mesenchymal stem cell therapy for immune and inflammatory diseases: use of cell-free products and human pluripotent stem cell-derived mesenchymal stem cells. *Stem Cells Transl Med*. 2021;10:1288-1303.
166. Mirzaei R, Zamani F, Hajibaba M, et al. The pathogenic, therapeutic and diagnostic role of exosomal microRNA in the autoimmune diseases. *J Neuroimmunol*. 2021;358:577640.

167. Liu H, Chen Y, Yin G, Xie Q. Therapeutic prospects of MicroRNAs carried by mesenchymal stem cells-derived extracellular vesicles in autoimmune diseases. *Life Sci.* 2021;277:119458.
168. Engeroff P, Vogel M. The potential of exosomes in allergy immunotherapy. *Vaccine.* 2022;10:133.
169. Paul WE, Zhu J. How are TH2-type immune responses initiated and amplified? *Nat Rev Immunol.* 2010;10:225-235.
170. Wu LC, Zarrin AA. The production and regulation of IgE by the immune system. *Nat Rev Immunol.* 2014;14:247-259.
171. Galli SJ, Tsai M, Piliponsky AM. The development of allergic inflammation. *Nature.* 2008;454:445-454.
172. Wernersson S, Pejler G. Mast cell secretory granules: armed for battle. *Nat Rev Immunol.* 2014;14:478-494.
173. Strunk RC, Bloomberg GR. Omalizumab for asthma. *New Engl J Med.* 2006;354:2689-2695.
174. Hovhannisyan L, Czechowska E, Gutowska-Owsiak D. The role of non-immune cell-derived extracellular vesicles in allergy. *Front Immunol.* 2021;12:702381.
175. Willart M, Hammad H. Lung dendritic cell-epithelial cell crosstalk in Th2 responses to allergens. *Curr Opin Immunol.* 2011;23:772-777.
176. Haj-Salem I, Plante S, Gounni A, Rouabhia M, Chakir J. Fibroblast-derived exosomes promote epithelial cell proliferation through TGF- β 2 signalling pathway in severe asthma. *Allergy.* 2018;73:178-186.
177. Khanna K, Chaudhuri R, Aich J, et al. Secretory inositol polyphosphate 4-phosphatase protects against airway inflammation and remodeling. *Am J Respir Cell Mol Biol.* 2019;60:399-412.
178. Gupta R, Radicioni G, Abdelwahab S, et al. Intercellular communication between airway epithelial cells is mediated by exosome-like vesicles. *Am J Respir Cell Mol Biol.* 2019;60:209-220.
179. Zhang M, Yu Q, Tang W, et al. Epithelial exosomal contactin-1 promotes monocyte-derived dendritic cell-dominant T-cell responses in asthma. *J Allergy Clin Immunol.* 2021;148:1545-1558.
180. Prado N, Marazuela EG, Segura E, et al. Exosomes from bronchoalveolar fluid of tolerized mice prevent allergic reaction. *J Immunol.* 2008;181:1519-1525.
181. Kulshreshtha A, Ahmad T, Agrawal A, Ghosh B. Proinflammatory role of epithelial cell-derived exosomes in allergic airway inflammation. *J Allergy Clin Immunol.* 2013;131:1194-1203. e1114.
182. Gon Y, Maruoka S, Inoue T, et al. Selective release of miRNAs via extracellular vesicles is associated with house-dust mite allergen-induced airway inflammation. *Clin Exp Allergy.* 2017;47:1586-1598.
183. Cho BS, Kim JO, Ha DH, Yi YW. Exosomes derived from human adipose tissue-derived mesenchymal stem cells alleviate atopic dermatitis. *Stem Cell Res Ther.* 2018;9:187.
184. Guo L, Lai P, Wang Y, et al. Extracellular vesicles from mesenchymal stem cells prevent contact hypersensitivity through the suppression of Tc1 and Th1 cells and expansion of regulatory T cells. *Int Immunopharmacol.* 2019;74:105663.
185. Fang S-B, Zhang H-Y, Wang C, et al. Small extracellular vesicles derived from human mesenchymal stromal cells prevent group 2 innate lymphoid cell-dominant allergic airway inflammation through delivery of miR-146a-5p. *J Extracell Vesicles.* 2020;9:1723260.
186. Shin K-O, Ha DH, Kim JO, et al. Exosomes from human adipose tissue-derived mesenchymal stem cells promote epidermal barrier repair by inducing de novo synthesis of ceramides in atopic dermatitis. *Cells.* 2020;9:680.
187. Nazimek K, Bustos-Morán E, Blas-Rus N, et al. Syngeneic red blood cell-induced extracellular vesicles suppress delayed-type hypersensitivity to self-antigens in mice. *Clin Exp Allergy.* 2019;49:1487-1499.
188. Ren J, Liu Y, Yao Y, et al. Intranasal delivery of MSC-derived exosomes attenuates allergic asthma via expanding IL-10 producing lung interstitial macrophages in mice. *Int Immunopharmacol.* 2021;91:107288.
189. Salimi M, Barlow JL, Saunders SP, et al. A role for IL-25 and IL-33-driven type-2 innate lymphoid cells in atopic dermatitis. *J Exp Med.* 2013;210:2939-2950.
190. Moy AP, Murali M, Kroshinsky D, Duncan LM, Nazarian RM. Immunologic overlap of helper T-cell subtypes 17 and 22 in erythrodermic psoriasis and atopic dermatitis. *JAMA Dermatol.* 2015;151:753-760.
191. Uter W, Werfel T, White IR, Johansen JD. Contact allergy: a review of current problems from a clinical perspective. *Int J Environ Res Public Health.* 2018;15:1108.
192. Hadžavdić SL, Pustišek N, Žužul K, Švigir A. Contact allergy: an update. *G Ital Dermatol Venereol.* 2018;153:419-428.
193. Simonsen AB, Johansen J, Deleuran M, Mortz CG, Sommerlund M. Contact allergy in children with atopic dermatitis: a systematic review. *Br J Dermatol.* 2017;177:395-405.
194. Justiz Vaillant A, Goyal A, Bansal P, Varacallo M. *Systemic Lupus Erythematosus.* StatPearls Publishing; 2021.
195. Yan S, Wang W, Gao G, et al. Key genes and functional coexpression modules involved in the pathogenesis of systemic lupus erythematosus. *J Cell Physiol.* 2018;233:8815-8825.
196. Tsokos GC, Lo MS, Reis PC, Sullivan KE. New insights into the immunopathogenesis of systemic lupus erythematosus. *Nat Rev Rheumatol.* 2016;12:716-730.
197. Yu F, Haas M, Glasscock R, Zhao M-H. Redefining lupus nephritis: clinical implications of pathophysiologic subtypes. *Nat Rev Nephrol.* 2017;13:483-495.
198. Pons-Estel GJ, Alarcón GS, Scofield L, Reinlib L, Cooper GS. Understanding the epidemiology and progression of systemic lupus erythematosus. *Seminars in Arthritis and Rheumatism.* Vol. 39. Elsevier; 2010:257-268.
199. Yang C, Sun J, Yang J, et al. Immunomodulatory effect of MSCs and MSCs-derived extracellular vesicles in systemic lupus erythematosus. *Front Immunol.* 2021;12:714832.
200. Cush JJ. Rheumatoid arthritis: early diagnosis and treatment. *Rheum Dis Clin.* 2022;48:537-547.
201. Smolen JS, Landewé RB, Bijlsma JW, et al. EULAR recommendations for the management of rheumatoid arthritis with synthetic and biological disease-modifying antirheumatic drugs: 2019 update. *Ann Rheum Dis.* 2020;79:685-699.
202. Bykerk VP, Shadick N, Frits M, et al. Flares in rheumatoid arthritis: frequency and management. A report from the BRASS registry. *J Rheumatol.* 2014;41:227-234.
203. Markuse IM, Dirven L, Gerards AH, et al. Disease flares in rheumatoid arthritis are associated with joint damage progression and disability: 10-year results from the BeSt study. *Arthritis Res Ther.* 2015;17:232.
204. Scherer HU, Häupl T, Burmester GR. The etiology of rheumatoid arthritis. *J Autoimmun.* 2020;110:102400.
205. Lin Y-J, Anzaghe M, Schülke S. Update on the pathomechanism, diagnosis, and treatment options for rheumatoid arthritis. *Cells.* 2020;9:880.
206. Ghoryani M, Shariati-Sarabi Z, Tavakkol-Afshari J, Mohammadi M. The sufficient immunoregulatory effect of autologous bone marrow-derived mesenchymal stem cell transplantation on regulatory T cells in patients with refractory rheumatoid arthritis. *J Immunol Res.* 2020;2020:3562753.
207. Ghoryani M, Shariati-Sarabi Z, Tavakkol-Afshari J, Ghasemi A, Poursamimi J, Mohammadi M. Amelioration of clinical symptoms of patients with refractory rheumatoid arthritis following treatment with autologous bone marrow-derived mesenchymal stem cells: a successful clinical trial in Iran. *Biomed Pharmacother.* 2019;109:1834-1840.
208. Cosenza S, Toupet K, Maumus M, et al. Mesenchymal stem cells-derived exosomes are more immunosuppressive than microparticles in inflammatory arthritis. *Theranostics.* 2018;8:1399-1410.

209. Zhang J, Rong Y, Luo C, Cui W. Bone marrow mesenchymal stem cell-derived exosomes prevent osteoarthritis by regulating synovial macrophage polarization. *Aging (Albany NY)*. 2020;12:25138-25152.
210. Cosenza S, Ruiz M, Toupet K, Jorgensen C, Noël D. Mesenchymal stem cells derived exosomes and microparticles protect cartilage and bone from degradation in osteoarthritis. *Sci Rep*. 2017;7:16214.
211. Cherif H, Bisson DG, Mannarino M, Rabau O, Ouellet JA, Haglund L. Senotherapeutic drugs for human intervertebral disc degeneration and low back pain. *elife*. 2020;9:e54693.
212. Hu Z-L, Li H-Y, Chang X, et al. Exosomes derived from stem cells as an emerging therapeutic strategy for intervertebral disc degeneration. *World J Stem Cells*. 2020;12:803-813.
213. Wu H, Zhou X, Wang X, et al. miR-34a in extracellular vesicles from bone marrow mesenchymal stem cells reduces rheumatoid arthritis inflammation via the cyclin I/ATM/ATR/p53 axis. *J Cell Mol Med*. 2021;25:1896-1910.
214. Zheng J, Zhu L, In II, Chen Y, Jia N, Zhu W. Bone marrow-derived mesenchymal stem cells-secreted exosomal microRNA-192-5p delays inflammatory response in rheumatoid arthritis. *Int Immunopharmacol*. 2020;78:105985.
215. Meng Q, Qiu B. Exosomal microRNA-320a derived from mesenchymal stem cells regulates rheumatoid arthritis fibroblast-like synoviocyte activation by suppressing CXCL9 expression. *Front Physiol*. 2020;11:441.
216. Chen Z, Wang H, Xia Y, Yan F, Lu Y. Therapeutic potential of mesenchymal cell-derived miRNA-150-5p-expressing exosomes in rheumatoid arthritis mediated by the modulation of MMP14 and VEGF. *J Immunol*. 2018;201:2472-2482.
217. Su Y, Liu Y, Ma C, Guan C, Ma X, Meng S. Mesenchymal stem cell-originated exosomal lncRNA HAND2-AS1 impairs rheumatoid arthritis fibroblast-like synoviocyte activation through miR-143-3p/TNFAIP3/NF- κ B pathway. *J Orthop Surg Res*. 2021;16:116.
218. Tsujimaru K, Takanashi M, Sudo K, et al. Extracellular microvesicles that originated adipose tissue derived mesenchymal stem cells have the potential ability to improve rheumatoid arthritis on mice. *Regen Ther*. 2020;15:305-311.
219. Wang L, Huang S, Li S, et al. Efficacy and safety of umbilical cord mesenchymal stem cell therapy for rheumatoid arthritis patients: a prospective phase I/II study. *Drug Des Devel Ther*. 2019;13:4331-4340.
220. Park EH, Lim H-s, Lee S, et al. Intravenous infusion of umbilical cord blood-derived mesenchymal stem cells in rheumatoid arthritis: a phase Ia clinical trial. *Stem Cells Transl Med*. 2018;7:636-642.
221. Miao H-b, Wang F, Lin S, Chen Z. Update on the role of extracellular vesicles in rheumatoid arthritis. *Expert Rev Mol Med*. 2022;24:e12.
222. Arden NK, Leyland K. Osteoarthritis year 2013 in review: clinical. *Osteoarthr Cartil*. 2013;21:1409-1413.
223. Kim SJ, Shetty AA, Kurian NM, et al. Articular cartilage repair using autologous collagen-induced chondrogenesis (ACIC): a pragmatic and cost-effective enhancement of a traditional technique. *Knee Surg Sports Traumatol Arthrosc*. 2020;28:2598-2603.
224. Kong L, Zheng L-Z, Qin L, Ho KK. Role of mesenchymal stem cells in osteoarthritis treatment. *J Orthop Translat*. 2017;9:89-103.
225. Zhang K, Zhao X, Chen X, et al. Enhanced therapeutic effects of mesenchymal stem cell-derived exosomes with an injectable hydrogel for hindlimb ischemia treatment. *ACS Appl Mater Interfaces*. 2018;10:30081-30091.
226. Liu X, Yang Y, Li Y, et al. Integration of stem cell-derived exosomes with in situ hydrogel glue as a promising tissue patch for articular cartilage regeneration. *Nanoscale*. 2017;9:4430-4438.
227. Li S, Stöckl S, Lukas C, et al. Curcumin-primed human BMSC-derived extracellular vesicles reverse IL-1 β -induced catabolic responses of OA chondrocytes by upregulating miR-126-3p. *Stem Cell Res Ther*. 2021;12:252.
228. Vonk LA, van Dooremalen SF, Liv N, et al. Mesenchymal stromal/stem cell-derived extracellular vesicles promote human cartilage regeneration in vitro. *Theranostics*. 2018;8:906-920.
229. Zhu Y, Wang Y, Zhao B, et al. Comparison of exosomes secreted by induced pluripotent stem cell-derived mesenchymal stem cells and synovial membrane-derived mesenchymal stem cells for the treatment of osteoarthritis. *Stem Cell Res Ther*. 2017;8:64.
230. Tao S-C, Yuan T, Zhang Y-L, Yin W-J, Guo S-C, Zhang C-Q. Exosomes derived from miR-140-5p-overexpressing human synovial mesenchymal stem cells enhance cartilage tissue regeneration and prevent osteoarthritis of the knee in a rat model. *Theranostics*. 2017;7:180-195.
231. Duan A, Shen K, Li B, et al. Extracellular vesicles derived from LPS-preconditioned human synovial mesenchymal stem cells inhibit extracellular matrix degradation and prevent osteoarthritis of the knee in a mouse model. *Stem Cell Res Ther*. 2021;12:427.
232. Wang Z, Yan K, Ge G, et al. Exosomes derived from miR-155-5p-overexpressing synovial mesenchymal stem cells prevent osteoarthritis via enhancing proliferation and migration, attenuating apoptosis, and modulating extracellular matrix secretion in chondrocytes. *Cell Biol Toxicol*. 2021;37:85-96.
233. Wang Y, Yu D, Liu Z, et al. Exosomes from embryonic mesenchymal stem cells alleviate osteoarthritis through balancing synthesis and degradation of cartilage extracellular matrix. *Stem Cell Res Ther*. 2017;8:189.
234. He L, He T, Xing J, et al. Bone marrow mesenchymal stem cell-derived exosomes protect cartilage damage and relieve knee osteoarthritis pain in a rat model of osteoarthritis. *Stem Cell Res Ther*. 2020;11:276.
235. Jin Z, Ren J, Qi S. Human bone mesenchymal stem cells-derived exosomes overexpressing microRNA-26a-5p alleviate osteoarthritis via down-regulation of PTGS2. *Int Immunopharmacol*. 2020;78:105946.
236. Zhou X, Liang H, Hu X, et al. BMSC-derived exosomes from congenital polydactyly tissue alleviate osteoarthritis by promoting chondrocyte proliferation. *Cell Death Discov*. 2020;6:142.
237. Liu Y, Lin L, Zou R, Wen C, Wang Z, Lin F. MSC-derived exosomes promote proliferation and inhibit apoptosis of chondrocytes via lncRNA-KLF3-AS1/miR-206/GIT1 axis in osteoarthritis. *Cell Cycle*. 2018;17:2411-2422.
238. Zhang S, Teo KYW, Chuah SJ, Lai RC, Lim SK, Toh WS. MSC exosomes alleviate temporomandibular joint osteoarthritis by attenuating inflammation and restoring matrix homeostasis. *Biomaterials*. 2019;200:35-47.
239. Chen P, Zheng L, Wang Y, et al. Desktop-stereolithography 3D printing of a radially oriented extracellular matrix/mesenchymal stem cell exosome bioink for osteochondral defect regeneration. *Theranostics*. 2019;9:2439-2459.
240. Wu J, Kuang L, Chen C, et al. miR-100-5p-abundant exosomes derived from infrapatellar fat pad MSCs protect articular cartilage and ameliorate gait abnormalities via inhibition of mTOR in osteoarthritis. *Biomaterials*. 2019;206:87-100.
241. Tofiño-Vian M, Guillén MI, Pérez del Caz MD, Castejón MA, Alcaraz MJ. Extracellular vesicles from adipose-derived mesenchymal stem cells downregulate senescence features in osteoarthritic osteoblasts. *Oxidative Med Cell Longev*. 2017;2017:7197598.
242. Shang X, Fang Y, Xin W, You H. The application of extracellular vesicles mediated miRNAs in osteoarthritis: current knowledge and perspective. *J Inflamm Res*. 2022;15:2583-2599.
243. Tao S-C, Huang J-Y, Gao Y, et al. Small extracellular vesicles in combination with sleep-related circRNA3503: a targeted therapeutic agent with injectable thermosensitive hydrogel to prevent osteoarthritis. *Bioact Mater*. 2021;6:4455-4469.
244. Qiu M, Liu D, Fu Q. MiR-129-5p shuttled by human synovial mesenchymal stem cell-derived exosomes relieves IL-1 β induced osteoarthritis via targeting HMGB1. *Life Sci*. 2021;269:118987.

245. Rong Y, Zhang J, Jiang D, et al. Hypoxic pretreatment of small extracellular vesicles mediates cartilage repair in osteoarthritis by delivering miR-216a-5p. *Acta Biomater.* 2021;122:325-342.
246. Wang K, Li F, Yuan Y, et al. Synovial mesenchymal stem cell-derived EV-packaged miR-31 downregulates histone demethylase KDM2A to prevent knee osteoarthritis. *Mol Ther Nucleic Acids.* 2020;22:1078-1091.
247. Chen X, Shi Y, Xue P, Ma X, Li J, Zhang J. Mesenchymal stem cell-derived exosomal microRNA-136-5p inhibits chondrocyte degeneration in traumatic osteoarthritis by targeting ELF3. *Arthritis Res Ther.* 2020;22:256.
248. Qiu B, Xu X, Yi P, Hao Y. Curcumin reinforces MSC-derived exosomes in attenuating osteoarthritis via modulating the miR-124/NF- κ B and miR-143/ROCK1/TLR9 signalling pathways. *J Cell Mol Med.* 2020;24:10855-10865.
249. Ragni E, Perucca Orfei C, De Luca P, et al. Inflammatory priming enhances mesenchymal stromal cell secretome potential as a clinical product for regenerative medicine approaches through secreted factors and EV-miRNAs: the example of joint disease. *Stem Cell Res Ther.* 2020;11:165.
250. Zhao C, Chen JY, Peng WM, Yuan B, Bi Q, Xu YJ. Exosomes from adipose-derived stem cells promote chondrogenesis and suppress inflammation by upregulating miR-145 and miR-221. *Mol Med Rep.* 2020;21:1881-1889.
251. Jing H, Zhang X, Luo K, et al. miR-381-abundant small extracellular vesicles derived from kartogenin-preconditioned mesenchymal stem cells promote chondrogenesis of MSCs by targeting TAOX1. *Biomaterials.* 2020;231:119682.
252. Luo P, Jiang C, Ji P, Wang M, Xu J. Exosomes of stem cells from human exfoliated deciduous teeth as an anti-inflammatory agent in temporomandibular joint chondrocytes via miR-100-5p/mTOR. *Stem Cell Res Ther.* 2019;10:216.
253. Sun H, Hu S, Zhang Z, Lun J, Liao W, Zhang Z. Expression of exosomal microRNAs during chondrogenic differentiation of human bone mesenchymal stem cells. *J Cell Biochem.* 2019;120:171-181.
254. Wang R, Xu B, Xu H. TGF- β 1 promoted chondrocyte proliferation by regulating Sp1 through MSC-exosomes derived miR-135b. *Cell Cycle.* 2018;17:2756-2765.
255. Shang X, Böker KO, Taheri S, Lehmann W, Schilling AF. Extracellular vesicles allow epigenetic mechanotransduction between chondrocytes and osteoblasts. *Int J Mol Sci.* 2021;22:13282.
256. Peng S, Yan Y, Li R, Dai H, Xu J. Extracellular vesicles from M1-polarized macrophages promote inflammation in the temporomandibular joint via miR-1246 activation of the Wnt/ β -catenin pathway. *Ann N Y Acad Sci.* 2021;1503:48-59.
257. Zhou Y, Ming J, Li Y, et al. Exosomes derived from miR-126-3p-overexpressing synovial fibroblasts suppress chondrocyte inflammation and cartilage degradation in a rat model of osteoarthritis. *Cell Death Discov.* 2021;7:37.
258. Wu X, Crawford R, Xiao Y, Mao X, Prasad I. Osteoarthritic subchondral bone release exosomes that promote cartilage degeneration. *Cells.* 2021;10:251.
259. Dai J, Dong R, Han X, et al. Osteoclast-derived exosomal let-7a-5p targets Smad2 to promote the hypertrophic differentiation of chondrocytes. *Am J Physiol Cell Physiol.* 2020;319:C21-C33.
260. Li Z, Wang Y, Xiang S, et al. Chondrocytes-derived exosomal miR-8485 regulated the Wnt/ β -catenin pathways to promote chondrogenic differentiation of BMSCs. *Biochem Biophys Res Commun.* 2020;523:506-513.
261. Ni Z, Kuang L, Chen H, et al. The exosome-like vesicles from osteoarthritic chondrocyte enhanced mature IL-1 β production of macrophages and aggravated synovitis in osteoarthritis. *Cell Death Dis.* 2019;10:522.
262. Mao G, Hu S, Zhang Z, et al. Exosomal miR-95-5p regulates chondrogenesis and cartilage degradation via histone deacetylase 2/8. *J Cell Mol Med.* 2018;22:5354-5366.
263. Xing Y, Chen X, Cao Y, Huang J, Xie X, Wei Y. Expression of Wnt and notch signaling pathways in inflammatory bowel disease treated with mesenchymal stem cell transplantation: evaluation in a rat model. *Stem Cell Res Ther.* 2015;6:101.
264. Song W-J, Li Q, Ryu M-O, et al. TSG-6 released from intraperitoneally injected canine adipose tissue-derived mesenchymal stem cells ameliorate inflammatory bowel disease by inducing M2 macrophage switch in mice. *Stem Cell Res Ther.* 2018;9:91.
265. Heidari N, Abbasi-Kenarsari H, Namaki S, et al. Adipose-derived mesenchymal stem cell-secreted exosome alleviates dextran sulfate sodium-induced acute colitis by Treg cell induction and inflammatory cytokine reduction. *J Cell Physiol.* 2021;236:5906-5920.
266. Yu H, Yang X, Xiao X, et al. Human adipose mesenchymal stem cell-derived exosomes protect mice from DSS-induced inflammatory bowel disease by promoting intestinal-stem-cell and epithelial regeneration. *Aging Dis.* 2021;12:1423-1437.
267. Li Y, Altemus J, Lightner AL. Mesenchymal stem cells and acellular products attenuate murine induced colitis. *Stem Cell Res Ther.* 2020;11:515.
268. Ma ZJ, Wang YH, Li ZG, et al. Immunosuppressive effect of exosomes from mesenchymal stromal cells in defined medium on experimental colitis. *Int J Stem Cells.* 2019;12:440-448.
269. Wu Y, Qiu W, Xu X, et al. Exosomes derived from human umbilical cord mesenchymal stem cells alleviate inflammatory bowel disease in mice through ubiquitination. *Am J Transl Res.* 2018;10:2026-2036.
270. Mao F, Wu Y, Tang X, et al. Exosomes derived from human umbilical cord mesenchymal stem cells relieve inflammatory bowel disease in mice. *Biomed Res Int.* 2017;2017:5356760.
271. Tolomeo AM, Castagliuolo I, Piccoli M, et al. Extracellular vesicles secreted by mesenchymal stromal cells exert opposite effects to their cells of origin in murine sodium dextran sulfate-induced colitis. *Front Immunol.* 2021;12:627605.
272. Cao L, Xu H, Wang G, Liu M, Tian D, Yuan Z. Extracellular vesicles derived from bone marrow mesenchymal stem cells attenuate dextran sodium sulfate-induced ulcerative colitis by promoting M2 macrophage polarization. *Int Immunopharmacol.* 2019;72:264-274.
273. Liu H, Liang Z, Wang F, et al. Exosomes from mesenchymal stromal cells reduce murine colonic inflammation via a macrophage-dependent mechanism. *JCI Insight.* 2019;4:e131273.
274. Tian J, Zhu Q, Zhang Y, et al. Olfactory ecto-mesenchymal stem cell-derived exosomes ameliorate experimental colitis via modulating Th1/Th17 and Treg cell responses. *Front Immunol.* 2020;11:598322.
275. Duan L, Huang H, Zhao X, et al. Extracellular vesicles derived from human placental mesenchymal stem cells alleviate experimental colitis in mice by inhibiting inflammation and oxidative stress. *Int J Mol Med.* 2020;46:1551-1561.
276. Gómez-Ferrer M, Amaro-Prellezo E, Dorronsoro A, et al. Hif-overexpression and pro-inflammatory priming in human mesenchymal stromal cells improves the healing properties of extracellular vesicles in experimental crohn's disease. *Int J Mol Sci.* 2021;22:11269.
277. Krzewska A, Ben-Skowronek I. Effect of associated autoimmune diseases on type 1 diabetes mellitus incidence and metabolic control in children and adolescents. *Biomed Res Int.* 2016;2016:6219730.
278. Esposito S, Toni G, Tascini G, Santi E, Berio MG, Principi N. Environmental factors associated with type 1 diabetes. *Front Endocrinol.* 2019;10:592.
279. Roep BO, Thomaidou S, van Tienhoven R, Zaldumbide A. Type 1 diabetes mellitus as a disease of the β -cell (do not blame the immune system?). *Nat Rev Endocrinol.* 2021;17:150-161.
280. Nakano M, Nagaishi K, Konari N, et al. Bone marrow-derived mesenchymal stem cells improve diabetes-induced cognitive impairment by exosome transfer into damaged neurons and astrocytes. *Sci Rep.* 2016;6:24805.

281. Nojehdehi S, Soudi S, Hesampour A, Rasouli S, Soleimani M, Hashemi SM. Immunomodulatory effects of mesenchymal stem cell-derived exosomes on experimental type-1 autoimmune diabetes. *J Cell Biochem*. 2018;119:9433-9443.
282. Shigemoto-Kuroda T, Oh JY, Kim D-k, et al. MSC-derived extracellular vesicles attenuate immune responses in two autoimmune murine models: type 1 diabetes and uveoretinitis. *Stem Cell Reports*. 2017;8:1214-1225.
283. Lin JT, Lane JM. Osteoporosis: a review. *Clin Orthop Relat Res*. 2004;425:126-134.
284. Qiu M, Zhai S, Fu Q, Liu D. Bone marrow mesenchymal stem cells-derived exosomal microRNA-150-3p promotes osteoblast proliferation and differentiation in osteoporosis. *Hum Gene Ther*. 2021;32:717-729.
285. Yang B-c, Kuang M-j, Kang J-y, Zhao J, Ma J-x, Ma X-l. Human umbilical cord mesenchymal stem cell-derived exosomes act via the miR-1263/Mob1/hippo signaling pathway to prevent apoptosis in disuse osteoporosis. *Biochem Biophys Res Commun*. 2020;524:883-889.
286. Yahao G, Xinjia W. The role and mechanism of exosomes from umbilical cord mesenchymal stem cells in inducing osteogenesis and preventing osteoporosis. *Cell Transplant*. 2021;30:9636897211057465.
287. Zhang L, Wang Q, Su H, Cheng J. Exosomes from adipose derived mesenchymal stem cells alleviate diabetic osteoporosis in rats through suppressing NLRP3 inflammasome activation in osteoclasts. *J Biosci Bioeng*. 2021;131:671-678.
288. Zhang X, Wang Y, Zhao H, et al. Extracellular vesicle-encapsulated miR-22-3p from bone marrow mesenchymal stem cell promotes osteogenic differentiation via FTO inhibition. *Stem Cell Res Ther*. 2020;11:227.
289. Xiao F, Zuo B, Tao B, et al. Exosomes derived from cyclic mechanical stretch-exposed bone marrow mesenchymal stem cells inhibit RANKL-induced osteoclastogenesis through the NF- κ B signaling pathway. *Ann Transl Med*. 2021;9:798.
290. Li X, Chen R, Li Y, et al. miR-27a-5p—abundant small extracellular vesicles derived from Epimedium-preconditioned bone mesenchymal stem cells stimulate osteogenesis by targeting Atg4B-mediated autophagy. *Front Cell Dev Biol*. 2021;2334.
291. Peng Z, Lu S, Lou Z, et al. Retracted Article: exosomes from bone marrow mesenchymal stem cells promoted osteogenic differentiation by delivering miR-196a that targeted Dickkopf-1 to activate Wnt/ β -catenin pathway. *Bioengineered*. 2023 Dec 3;14(1):1996015.
292. Li L, Zhou X, Zhang J-t, et al. Exosomal miR-186 derived from BMSCs promote osteogenesis through hippo signaling pathway in postmenopausal osteoporosis. *J Orthop Surg Res*. 2021;16:23.
293. Liu W, Huang J, Chen F, et al. MSC-derived small extracellular vesicles overexpressing miR-20a promoted the osteointegration of porous titanium alloy by enhancing osteogenesis via targeting BAMBI. *Stem Cell Res Ther*. 2021;12:348.
294. Fu M, Fang L, Xiang X, Fan X, Wu J, Wang J. Microarray analysis of circRNAs sequencing profile in exosomes derived from bone marrow mesenchymal stem cells in postmenopausal osteoporosis patients. *J Clin Lab Anal*. 2022;36:e23916.
295. Zhao P, Xiao L, Peng J, Qian Y, Huang C. Exosomes derived from bone marrow mesenchymal stem cells improve osteoporosis through promoting osteoblast proliferation via MAPK pathway. *Eur Rev Med Pharmacol Sci*. 2018;22:3962-3970.
296. Cai X, Zhang Z-Y, Yuan J-T, et al. hucMSC-derived exosomes attenuate colitis by regulating macrophage pyroptosis via the miR-378a-5p/NLRP3 axis. *Stem Cell Res Ther*. 2021;12:416.
297. Wang G, Yuan J, Cai X, et al. HucMSC-exosomes carrying miR-326 inhibit neddylation to relieve inflammatory bowel disease in mice. *Clin Transl Med*. 2020;10:e113.
298. Yang S, Liang X, Song J, et al. A novel therapeutic approach for inflammatory bowel disease by exosomes derived from human umbilical cord mesenchymal stem cells to repair intestinal barrier via TSG-6. *Stem Cell Res Ther*. 2021;12:315.
299. Li S, Liu J, Liu S, Jiao W, Wang X. Chitosan oligosaccharides packaged into rat adipose mesenchymal stem cells-derived extracellular vesicles facilitating cartilage injury repair and alleviating osteoarthritis. *J Nanobiotechnol*. 2021;19:343.
300. Huang B, Su Y, Shen E, Song M, Liu D, Qi H. Extracellular vesicles from GPNMB-modified bone marrow mesenchymal stem cells attenuate bone loss in an ovariectomized rat model. *Life Sci*. 2021;272:119208.
301. Baharlou H, Nouraei Z, Azimi M, et al. Umbilical cord mesenchymal stem cells as well as their released exosomes suppress proliferation of activated PBMCs in multiple sclerosis. *Scand J Immunol*. 2021;93:e13013.
302. Shamili FH, Alibolandi M, Rafatpanah H, et al. Immunomodulatory properties of MSC-derived exosomes armed with high affinity aptamer toward myelin as a platform for reducing multiple sclerosis clinical score. *J Control Release*. 2019;299:149-164.
303. Giunti D, Marini C, Parodi B, et al. Role of miRNAs shuttled by mesenchymal stem cell-derived small extracellular vesicles in modulating neuroinflammation. *Sci Rep*. 2021;11:1740.
304. Li B, Xing Y, Gan Y, He J, Hua H. Labial gland-derived mesenchymal stem cells and their exosomes ameliorate murine Sjögren's syndrome by modulating the balance of Treg and Th17 cells. *Stem Cell Res Ther*. 2021;12:478.
305. Rui K, Hong Y, Zhu Q, et al. Olfactory ecto-mesenchymal stem cell-derived exosomes ameliorate murine Sjögren's syndrome by modulating the function of myeloid-derived suppressor cells. *Cell Mol Immunol*. 2021;18:440-451.
306. Chen X, Wei Q, Sun H, et al. Exosomes derived from human umbilical cord mesenchymal stem cells regulate macrophage polarization to attenuate systemic lupus erythematosus-associated diffuse alveolar hemorrhage in mice. *Int J Stem Cells*. 2021;14:331-340.
307. Dou R, Zhang X, Xu X, Wang P, Yan B. Mesenchymal stem cell exosomal tsRNA-21109 alleviate systemic lupus erythematosus by inhibiting macrophage M1 polarization. *Mol Immunol*. 2021;139:106-114.
308. Rozier P, Maumus M, Maria ATJ, et al. Lung fibrosis is improved by extracellular vesicles from IFN γ -primed mesenchymal stromal cells in murine systemic sclerosis. *Cells*. 2021;10:2727.
309. Rozier P, Maumus M, Maria ATJ, et al. Mesenchymal stromal cells-derived extracellular vesicles alleviate systemic sclerosis via miR-29a-3p. *J Autoimmun*. 2021;121:102660.
310. Baral H, Uchiyama A, Yokoyama Y, et al. Antifibrotic effects and mechanisms of mesenchymal stem cell-derived exosomes in a systemic sclerosis mouse model: possible contribution of miR-196b-5p. *J Dermatol Sci*. 2021;104:39-47.
311. Jin J, Ou Q, Wang Z, et al. BMSC-derived extracellular vesicles intervened the pathogenic changes of scleroderma in mice through miRNAs. *Stem Cell Res Ther*. 2021;12:327.
312. Li M, Zhang H-P, Wang X-Y, Chen Z-G, Lin X-F, Zhu W. Mesenchymal stem cell-derived exosomes ameliorate dermal fibrosis in a murine model of bleomycin-induced scleroderma. *Stem Cells Dev*. 2021;30:981-990.
313. Guo L, Lai P, Wang Y, et al. Extracellular vesicles derived from mesenchymal stem cells prevent skin fibrosis in the cGVHD mouse model by suppressing the activation of macrophages and B cells immune response. *Int Immunopharmacol*. 2020;84:106541.
314. Soundara Rajan T, Giacompo S, Diomedea F, Bramanti P, Trubiani O, Mazzon E. Human periodontal ligament stem cells secretome from multiple sclerosis patients suppresses NALP3 inflammasome activation in experimental autoimmune encephalomyelitis. *Int J Immunopathol Pharmacol*. 2017;30:238-252.

315. Meng H-Y, Chen L-Q, Chen L-H. The inhibition by human MSCs-derived miRNA-124a overexpression exosomes in the proliferation and migration of rheumatoid arthritis-related fibroblast-like synovioocyte cell. *BMC Musculoskelet Disord*. 2020;21:150.
316. Mahdipour E, Salmasi Z, Sabeti N. Potential of stem cell-derived exosomes to regenerate β islets through Pdx-1 dependent mechanism in a rat model of type 1 diabetes. *J Cell Physiol*. 2019;234:20310-20321.
317. Zhu L, Huang X, Yu W, Chen H, Chen Y, Dai Y. Transplantation of adipose tissue-derived stem cell-derived exosomes ameliorates erectile function in diabetic rats. *Andrologia*. 2018;50:e12871.
318. Chen C, Wang D, Moshaverinia A, et al. Mesenchymal stem cell transplantation in tight-skin mice identifies miR-151-5p as a therapeutic target for systemic sclerosis. *Cell Res*. 2017;27:559-577.
319. Bai L, Shao H, Wang H, et al. Author correction: effects of mesenchymal stem cell-derived exosomes on experimental autoimmune uveitis. *Sci Rep*. 2018;8:9889.
320. Chen L, Lu F-b, Chen D-z, et al. BMSCs-derived miR-223-containing exosomes contribute to liver protection in experimental autoimmune hepatitis. *Mol Immunol*. 2018;93:38-46.
321. Shpilsky D, Bambs C, Kip K, et al. Association between ideal cardiovascular health and markers of subclinical cardiovascular disease. *Clin Cardiol*. 2018;41:1593-1599.
322. Hu SS. Report on cardiovascular health and diseases burden in China: an updated summary of 2020. *Chin Circ J*. 2021;36:521-545.
323. Chen Q, Liu Y, Ding X, et al. Bone marrow mesenchymal stem cell-secreted exosomes carrying microRNA-125b protect against myocardial ischemia reperfusion injury via targeting SIRT7. *Mol Cell Biochem*. 2020;465:103-114.
324. Cui M, Han Y, Yang J, Li G, Yang C. A narrative review of the research status of exosomes in cardiovascular disease. *Ann Palliat Med*. 2022;11:363-377.
325. Virani SS, Alonso A, Aparicio HJ, et al. Heart disease and stroke statistics-2021 update: a report from the American Heart Association. *Circulation*. 2021;143:e254-e743.
326. Ye L, D'Agostino G, Loo SJ, et al. Early regenerative capacity in the porcine heart. *Circulation*. 2018;138:2798-2808.
327. Porrello ER, Mahmoud AI, Simpson E, et al. Transient regenerative potential of the neonatal mouse heart. *Science*. 2011;331:1078-1080.
328. Yan H, Rao X, Wang R, Zhu S, Liu R, Zheng X. Cell cycle withdrawal limit the regenerative potential of neonatal cardiomyocytes. *Cardiovasc Eng Technol*. 2021;12:475-484.
329. Nie S, Wang X, Sivakumaran P, et al. Biologically active constituents of the secretome of human W8B2⁺ cardiac stem cells. *Sci Rep*. 2018;8:1579.
330. Wagner T, Traxler D, Simader E, et al. Different pro-angiogenic potential of γ -irradiated PBMC-derived secretome and its subfractions. *Sci Rep*. 2018;8:18016.
331. Korf-Klingebiel M, Reboll MR, Klede S, et al. Myeloid-derived growth factor (C19orf10) mediates cardiac repair following myocardial infarction. *Nat Med*. 2015;21:140-149.
332. Selvasandran K, Makhoul G, Jaiswal PK, et al. A tumor necrosis factor- α and hypoxia-induced secretome therapy for myocardial repair. *Ann Thorac Surg*. 2018;105:715-723.
333. Waters R, Alam P, Pacelli S, Chakravarti AR, Ahmed RP, Paul A. Stem cell-inspired secretome-rich injectable hydrogel to repair injured cardiac tissue. *Acta Biomater*. 2018;69:95-106.
334. Li H, Zuo S, He Z, et al. Paracrine factors released by GATA-4 overexpressed mesenchymal stem cells increase angiogenesis and cell survival. *Am J Physiol Heart Circ Physiol*. 2010;299:H1772-H1781.
335. Kishore R, Khan M. More than tiny sacks: stem cell exosomes as cell-free modality for cardiac repair. *Circ Res*. 2016;118:330-343.
336. Wang W, Zheng H. Myocardial infarction: the protective role of MiRNAs in myocardium pathology. *Front Cardiovasc Med*. 2021;8:631817.
337. Zhao J, Li X, Hu J, et al. Mesenchymal stromal cell-derived exosomes attenuate myocardial ischaemia-reperfusion injury through miR-182-regulated macrophage polarization. *Cardiovasc Res*. 2019;115:1205-1216.
338. Youn S-W, Li Y, Kim Y-M, et al. Modification of cardiac progenitor cell-derived exosomes by miR-322 provides protection against myocardial infarction through nox2-dependent angiogenesis. *Antioxidants*. 2019;8:18.
339. Chen L, Wang Y, Pan Y, et al. Cardiac progenitor-derived exosomes protect ischemic myocardium from acute ischemia/reperfusion injury. *Biochem Biophys Res Commun*. 2013;431:566-571.
340. Wang Y, Zhang L, Li Y, et al. Exosomes/microvesicles from induced pluripotent stem cells deliver cardioprotective miRNAs and prevent cardiomyocyte apoptosis in the ischemic myocardium. *Int J Cardiol*. 2015;192:61-69.
341. Zhang J, Zhang X. Ischaemic preconditioning-induced serum exosomes protect against myocardial ischaemia/reperfusion injury in rats by activating the PI3K/AKT signalling pathway. *Cell Biochem Funct*. 2021;39:287-295.
342. Dai Y, Wang S, Chang S, et al. M2 macrophage-derived exosomes carry microRNA-148a to alleviate myocardial ischemia/reperfusion injury via inhibiting TXNIP and the TLR4/NF- κ B/NLRP3 inflammasome signaling pathway. *J Mol Cell Cardiol*. 2020;142:65-79.
343. Tang J, Jin L, Liu Y, et al. Exosomes derived from mesenchymal stem cells protect the myocardium against ischemia/reperfusion injury through inhibiting pyroptosis. *Drug Des Devel Ther*. 2020;14:3765-3775.
344. Luo H, Li X, Li T, et al. microRNA-423-3p exosomes derived from cardiac fibroblasts mediates the cardioprotective effects of ischemic post-conditioning. *Cardiovasc Res*. 2019;115:1189-1204.
345. Peng Y, Zhao J-L, Peng Z-Y, Xu W-F, Yu G-L. Exosomal miR-25-3p from mesenchymal stem cells alleviates myocardial infarction by targeting pro-apoptotic proteins and EZH2. *Cell Death Dis*. 2020;11:317.
346. Zhu D, Wang Y, Thomas M, et al. Exosomes from adipose-derived stem cells alleviate myocardial infarction via microRNA-31/FIHL1/HIF-1 α pathway. *J Mol Cell Cardiol*. 2022;162:10-19.
347. Pan J, Alimujiang M, Chen Q, Shi H, Luo X. Exosomes derived from miR-146a-modified adipose-derived stem cells attenuate acute myocardial infarction-induced myocardial damage via downregulation of early growth response factor 1. *J Cell Biochem*. 2019;120:4433-4443.
348. Gallet R, Dawkins J, Valle J, et al. Exosomes secreted by cardiosphere-derived cells reduce scarring, attenuate adverse remodelling, and improve function in acute and chronic porcine myocardial infarction. *Eur Heart J*. 2017;38:201-211.
349. Li S, Jiang J, Yang Z, Li Z, Ma X, Li X. Cardiac progenitor cell-derived exosomes promote H9C2 cell growth via Akt/mTOR activation. *Int J Mol Med*. 2018;42:1517-1525.
350. Agarwal U, George A, Bhutani S, et al. Experimental, systems, and computational approaches to understanding the microRNA-mediated reparative potential of cardiac progenitor cell-derived exosomes from pediatric patients. *Circ Res*. 2017;120:701-712.
351. Sahoo S, Klychko E, Thorne T, et al. Exosomes from human CD34⁺ stem cells mediate their proangiogenic paracrine activity. *Circ Res*. 2011;109:724-728.
352. Shao L, Zhang Y, Lan B, et al. MiRNA-sequence indicates that mesenchymal stem cells and exosomes have similar mechanism to enhance cardiac repair. *Biomed Res Int*. 2017;2017:4150705.
353. Zhu J, Lu K, Zhang N, et al. Myocardial reparative functions of exosomes from mesenchymal stem cells are enhanced by hypoxia treatment of the cells via transferring microRNA-210 in an nSMase2-dependent way. *Artif Cells Nanomed Biotechnol*. 2018;46:1659-1670.
354. Feng Y, Huang W, Wani M, Yu X, Ashraf M. Ischemic preconditioning potentiates the protective effect of stem cells through

- secretion of exosomes by targeting Mecp2 via miR-22. *PLoS One*. 2014;9:e88685.
355. Zhu L-P, Tian T, Wang J-Y, et al. Hypoxia-elicited mesenchymal stem cell-derived exosomes facilitates cardiac repair through miR-125b-mediated prevention of cell death in myocardial infarction. *Theranostics*. 2018;8:6163-6177.
 356. Gonzalez-King H, García NA, Ontoria-Oviedo I, Ciria M, Montero JA, Sepúlveda P. Hypoxia inducible factor-1 α potentiates jagged 1-mediated angiogenesis by mesenchymal stem cell-derived exosomes. *Stem Cells*. 2017;35:1747-1759.
 357. Zhang Z, Yang J, Yan W, Li Y, Shen Z, Asahara T. Pretreatment of cardiac stem cells with exosomes derived from mesenchymal stem cells enhances myocardial repair. *J Am Heart Assoc*. 2016;5:e002856.
 358. Zhou B, Bentham J, Di Cesare M, et al. Worldwide trends in blood pressure from 1975 to 2015: a pooled analysis of 1479 population-based measurement studies with 19.1 million participants. *Lancet*. 2017;389:37-55.
 359. Mills K, Stefanescu A, He J. The global epidemiology of hypertension. *Nat Rev Nephrol*. 2020;16:223-237.
 360. Liu ZZ, Jose PA, Yang J, Zeng C. Importance of extracellular vesicles in hypertension. *Exp Biol Med*. 2021;246:342-353.
 361. Tong Y, Ye C, Ren X-S, et al. Exosome-mediated transfer of ACE (angiotensin-converting enzyme) from adventitial fibroblasts of spontaneously hypertensive rats promotes vascular smooth muscle cell migration. *Hypertension*. 2018;72:881-888.
 362. Preston RA, Jy W, Jimenez JJ, et al. Effects of severe hypertension on endothelial and platelet microparticles. *Hypertension*. 2003;41:211-217.
 363. Erdbrügger U, Le TH. Extracellular vesicles in renal diseases: more than novel biomarkers? *J Am Soc Nephrol*. 2016;27:12-26.
 364. Wang J, Su C, Wang Y, et al. Elevated circulating endothelial microparticles and brachial-ankle pulse wave velocity in well-controlled hypertensive patients. *J Hum Hypertens*. 2009;23:307-315.
 365. Huang P-H, Huang S-S, Chen Y-H, et al. Increased circulating CD31+/annexin V+ apoptotic microparticles and decreased circulating endothelial progenitor cell levels in hypertensive patients with microalbuminuria. *J Hypertens*. 2010;28:1655-1665.
 366. Moriya H, Kobayashi S, Ohtake T, et al. Aliskiren, a direct renin inhibitor, improves vascular endothelial function in patients on hemodialysis independent of antihypertensive effect~ a pilot study~. *Kidney Blood Press Res*. 2013;37:190-198.
 367. Baldan-Martin M, Mourino-Alvarez L, Gonzalez-Calero L, et al. Plasma molecular signatures in hypertensive patients with renin-angiotensin system suppression: new predictors of renal damage and de novo albuminuria indicators. *Hypertension*. 2016;68:157-166.
 368. Gilead JJ, Carlson JM, Schoeffel CD, Carey RM, Felder RA. Urinary exosome miRNome analysis and its applications to salt sensitivity of blood pressure. *Clin Biochem*. 2013;46:1131-1134.
 369. Zhu X, Badawi M, Pomeroy S, et al. Comprehensive toxicity and immunogenicity studies reveal minimal effects in mice following sustained dosing of extracellular vesicles derived from HEK293T cells. *J Extracell Vesicles*. 2017;6:1324730.
 370. de Abreu RC, Fernandes H, da Costa Martins PA, Sahoo S, Emanueli C, Ferreira L. Native and bioengineered extracellular vesicles for cardiovascular therapeutics. *Nat Rev Cardiol*. 2020;17:685-697.
 371. Kalluri R, LeBleu VS. The biology, function, and biomedical applications of exosomes. *Science*. 2020;367:eaau6977.
 372. Good ME, Musante L, La Salvia S, et al. Circulating extracellular vesicles in normotension restrain vasodilation in resistance arteries. *Hypertension*. 2020;75:218-228.
 373. Shaihov-Teper O, Ram E, Ballan N, et al. Extracellular vesicles from epicardial fat facilitate atrial fibrillation. *Circulation*. 2021;143:2475-2493.
 374. Kornej J, Börschel CS, Benjamin EJ, Schnabel RB. Epidemiology of atrial fibrillation in the 21st century: novel methods and new insights. *Circ Res*. 2020;127:4-20.
 375. Wijesurendra RS, Casadei B. Mechanisms of atrial fibrillation. *Heart*. 2019;105:1860-1867.
 376. Krijthe BP, Kunst A, Benjamin EJ, et al. Projections on the number of individuals with atrial fibrillation in the European Union, from 2000 to 2060. *Eur Heart J*. 2013;34:2746-2751.
 377. Clauss S, Sinner MF, Käb S, Wakili R. The role of microRNAs in antiarrhythmic therapy for atrial fibrillation. *Arrhythmia Electrophysiol Rev*. 2015;4:146-155.
 378. Ng FS, Handa BS, Li X, Peters NS. Toward mechanism-directed electrophenotype-based treatments for atrial fibrillation. *Front Physiol*. 2020;11:987.
 379. Philips T, Taghji P, El Haddad M, et al. Improving procedural and one-year outcome after contact force-guided pulmonary vein isolation: the role of interlesion distance, ablation index, and contact force variability in the 'CLOSE'-protocol. *Europace*. 2018;20:f419-f427.
 380. Wei Z, Bing Z, Shaohuan Q, et al. Expression of miRNAs in plasma exosomes derived from patients with atrial fibrillation. *Clin Cardiol*. 2020;43:1450-1459.
 381. Mun D, Kim H, Kang JY, et al. Expression of miRNAs in circulating exosomes derived from patients with persistent atrial fibrillation. *FASEB J*. 2019;33:5979-5989.
 382. Members WC, Hunt SA, Abraham WT, et al. ACC/AHA 2005 guideline update for the diagnosis and management of chronic heart failure in the adult—summary article: a report of the American College of Cardiology/American Heart Association task force on practice guidelines (writing committee to update the 2001 guidelines for the evaluation and Management of Heart Failure): developed in collaboration with the American College of Chest Physicians and the International Society for Heart and Lung Transplantation: endorsed by the Heart Rhythm Society. *Circulation*. 2005;112:1825-1852.
 383. Benjamin EJ, Muntner P, Alonso A, et al. Heart disease and stroke statistics—2019 update: a report from the American Heart Association. *Circulation*. 2019;139:e56-e528.
 384. Huffman MD, Berry JD, Ning H, et al. Lifetime risk for heart failure among white and black Americans: cardiovascular lifetime risk pooling project. *J Am Coll Cardiol*. 2013;61:1510-1517.
 385. Heidenreich PA, Albert NM, Allen LA, et al. Forecasting the impact of heart failure in the United States: a policy statement from the American Heart Association. *Circ Heart Fail*. 2013;6:606-619.
 386. Chen F, Li X, Zhao J, Geng J, Xie J, Xu B. Bone marrow mesenchymal stem cell-derived exosomes attenuate cardiac hypertrophy and fibrosis in pressure overload induced remodeling. *In Vitro Cell Dev Biol Anim*. 2020;56:567-576.
 387. Kobiyama K, Ley K. Atherosclerosis: a chronic inflammatory disease with an autoimmune component. *Circ Res*. 2018;123:1118-1120.
 388. Davies PF. Hemodynamic shear stress and the endothelium in cardiovascular pathophysiology. *Nat Clin Pract Cardiovasc Med*. 2009;6:16-26.
 389. Ding N, Sang Y, Chen J, et al. Cigarette smoking, smoking cessation, and long-term risk of 3 major atherosclerotic diseases. *J Am Coll Cardiol*. 2019;74:498-507.
 390. Furukawa S, Fujita T, Shimabukuro M, et al. Increased oxidative stress in obesity and its impact on metabolic syndrome. *J Clin Invest*. 2017;114:1752-1761.
 391. Poznyak A, Grechko AV, Poggio P, Myasoedova VA, Alfieri V, Orekhov AN. The diabetes mellitus-atherosclerosis connection: the role of lipid and glucose metabolism and chronic inflammation. *Int J Mol Sci*. 2020;21:1835.
 392. Danenberg HD, Welt FG, Walker M III, Seifert P, Toegel GS, Edelman ER. Systemic inflammation induced by lipopolysaccharide increases neointimal formation after balloon and stent injury in rabbits. *Circulation*. 2002;105:2917-2922.

393. Dzau VJ, Braun-Dullaeus RC, Sedding DG. Vascular proliferation and atherosclerosis: new perspectives and therapeutic strategies. *Nat Med*. 2002;8:1249-1256.
394. Libby P, Ridker PM, Hansson GK. Progress and challenges in translating the biology of atherosclerosis. *Nature*. 2011;473:317-325.
395. Davis GE, Senger DR. Endothelial extracellular matrix: biosynthesis, remodeling, and functions during vascular morphogenesis and neovessel stabilization. *Circ Res*. 2005;97:1093-1107.
396. Yao Y, Sun W, Sun Q, et al. Platelet-derived exosomal MicroRNA-25-3p inhibits coronary vascular endothelial cell inflammation through Adam10 via the NF- κ B signaling pathway in ApoE^{-/-} mice. *Front Immunol*. 2019;10:2205.
397. Zhong X, Gao W, Wu R, Liu H, Ge J. Dendritic cell exosome-shuttled miRNA146a regulates exosome-induced endothelial cell inflammation by inhibiting Irak-1: a feedback control mechanism. *Mol Med Rep*. 2019;20:5315-5323.
398. Bouchareychas L, Duong P, Covarrubias S, et al. Macrophage exosomes resolve atherosclerosis by regulating hematopoiesis and inflammation via MicroRNA cargo. *Cell Rep*. 2020;32:107881.
399. Li J, Xue H, Li T, et al. Exosomes derived from mesenchymal stem cells attenuate the progression of atherosclerosis in ApoE^{-/-} mice via miR-let7 mediated infiltration and polarization of M2 macrophage. *Biochem Biophys Res Commun*. 2019;510:565-572.
400. Zhang Y-G, Song Y, Guo X-L, et al. Exosomes derived from oxLDL-stimulated macrophages induce neutrophil extracellular traps to drive atherosclerosis. *Cell Cycle*. 2019;18:2672-2682.
401. Chen L, Hu L, Li Q, Ma J, Li H. Exosome-encapsulated miR-505 from ox-LDL-treated vascular endothelial cells aggravates atherosclerosis by inducing NET formation. *Acta Biochim Biophys Sin*. 2019;51:1233-1241.
402. Lin B, Xie W, Zeng C, et al. Transfer of exosomal microRNA-203-3p from dendritic cells to bone marrow-derived macrophages reduces development of atherosclerosis by downregulating Ctss in mice. *Aging (Albany NY)*. 2021;13:15638-15658.
403. Liu Y, Sun Y, Lin X, et al. Perivascular adipose-derived exosomes reduce foam cell formation by regulating expression of cholesterol transporters. *Front Cardiovasc Med*. 2021;8:697510.
404. Joladarashi D, Kishore R. Mesenchymal stromal cell exosomes in cardiac repair. *Curr Cardiol Rep*. 2022;24:405-417.
405. Zhou M, Wang H, Zeng X, et al. Mortality, morbidity, and risk factors in China and its provinces, 1990-2017: a systematic analysis for the global burden of disease study 2017. *Lancet*. 2019;394:1145-1158.
406. Benjamin EJ, Blaha MJ, Chiuve SE, et al. Heart disease and stroke statistics—2017 update: a report from the American Heart Association. *Circulation*. 2017;135:e146-e603.
407. Shen F, Jiang L, Han F, Degos V, Chen S, Su H. Increased inflammatory response in old mice is associated with more severe neuronal injury at the acute stage of ischemic stroke. *Aging Dis*. 2019;10:12.
408. Jayaraj RL, Azimullah S, Beiram R, Jalal FY, Rosenberg GA. Neuroinflammation: friend and foe for ischemic stroke. *J Neuroinflammation*. 2019;16:142.
409. Rust R, Grönnert L, Schwab ME. Inflammation after stroke: a local rather than systemic response? *Trends Neurosci*. 2018;41:877-879.
410. Nakamura K, Shichita T. Cellular and molecular mechanisms of sterile inflammation in ischaemic stroke. *J Biochem*. 2019;165:459-464.
411. Tsuyama J, Nakamura A, Ooboshi H, Yoshimura A, Shichita T. Pivotal role of innate myeloid cells in cerebral post-ischemic sterile inflammation. *Seminars in Immunopathology*. Vol. 40. Springer; 2018:523-538.
412. Ma Y, Liu Y, Zhang Z, Yang G-Y. Significance of complement system in ischemic stroke: a comprehensive review. *Aging Dis*. 2019;10:429-462.
413. Deng L, Peng Q, Wang H, et al. Intrathecal injection of allogenic bone marrow-derived mesenchymal stromal cells in treatment of patients with severe ischemic stroke: study protocol for a randomized controlled observer-blinded trial. *Transl Stroke Res*. 2019;10:170-177.
414. Zhang ZG, Buller B, Chopp M. Exosomes—beyond stem cells for restorative therapy in stroke and neurological injury. *Nat Rev Neurol*. 2019;15:193-203.
415. Chen J, Chopp M. Exosome therapy for stroke. *Stroke*. 2018;49:1083-1090.
416. Li Y, Tang Y, Yang G-Y. Therapeutic application of exosomes in ischaemic stroke. *Stroke Vasc Neurol*. 2021;6:483-495.
417. Xin H, Li Y, Cui Y, Yang JJ, Zhang ZG, Chopp M. Systemic administration of exosomes released from mesenchymal stromal cells promote functional recovery and neurovascular plasticity after stroke in rats. *J Cereb Blood Flow Metab*. 2013;33:1711-1715.
418. Doeppner TR, Herz J, Görgens A, et al. Extracellular vesicles improve post-stroke neuroregeneration and prevent postischemic immunosuppression. *Stem Cells Transl Med*. 2015;4:1131-1143.
419. Hu B, Chen S, Zou M, He Z, Shao S, Liu B. Effect of extracellular vesicles on neural functional recovery and immunologic suppression after rat cerebral apoplexy. *Cell Physiol Biochem*. 2016;40:155-162.
420. Lee JY, Kim E, Choi S-M, et al. Microvesicles from brain-extract-treated mesenchymal stem cells improve neurological functions in a rat model of ischemic stroke. *Sci Rep*. 2016;6:33038.
421. Ophelders DR, Wolfs TG, Jellema RK, et al. Mesenchymal stromal cell-derived extracellular vesicles protect the fetal brain after hypoxia-ischemia. *Stem Cells Transl Med*. 2016;5:754-763.
422. Zhang H, Wu J, Wu J, et al. Exosome-mediated targeted delivery of miR-210 for angiogenic therapy after cerebral ischemia in mice. *J Nanobiotechnol*. 2019;17:1-13.
423. Yang J, Zhang X, Chen X, Wang L, Yang G. Exosome mediated delivery of miR-124 promotes neurogenesis after ischemia. *Mol Ther Nucleic Acids*. 2017;7:278-287.
424. Otero-Ortega L, Laso-García F, Gómez-de Frutos MC, et al. White matter repair after extracellular vesicles administration in an experimental animal model of subcortical stroke. *Sci Rep*. 2017;7:44433.
425. Deng M, Xiao H, Zhang H, et al. Mesenchymal stem cell-derived extracellular vesicles ameliorates hippocampal synaptic impairment after transient global ischemia. *Front Cell Neurosci*. 2017;11:205.
426. Nalamolu KR, Venkatesh I, Mohandass A, et al. Exosomes secreted by the cocultures of normal and oxygen-glucose-deprived stem cells improve post-stroke outcome. *Neuromolecular Med*. 2019;21:529-539.
427. Thomi G, Joerger-Messerli M, Haesler V, Muri L, Surbek D, Schoeberlein A. Intranasally administered exosomes from umbilical cord stem cells have preventive neuroprotective effects and contribute to functional recovery after perinatal brain injury. *Cells*. 2019;8:855.
428. Xin H, Wang F, Li Y, et al. Secondary release of exosomes from astrocytes contributes to the increase in neural plasticity and improvement of functional recovery after stroke in rats treated with exosomes harvested from microRNA 133b-overexpressing multipotent mesenchymal stromal cells. *Cell Transplant*. 2017;26:243-257.
429. Moore T, Bowley B, Pessina M, et al. Mesenchymal derived exosomes enhance recovery of motor function in a monkey model of cortical injury. *Restor Neurol Neurosci*. 2019;37:347-362.
430. Go V, Bowley BG, Pessina MA, et al. Extracellular vesicles from mesenchymal stem cells reduce microglial-mediated neuroinflammation after cortical injury in aged rhesus monkeys. *Geroscience*. 2020;42:1-17.
431. Medalla M, Chang W, Calderazzo SM, et al. Treatment with mesenchymal-derived extracellular vesicles reduces injury-related pathology in pyramidal neurons of monkey perilesional ventral premotor cortex. *J Neurosci*. 2020;40:3385-3407.
432. Liu W, Bai X, Zhang A, Huang J, Xu S, Zhang J. Role of exosomes in central nervous system diseases. *Front Mol Neurosci*. 2019;12:240.

433. Thomi G, Surbek D, Haesler V, Joerger-Messerli M, Schoeberlein A. Exosomes derived from umbilical cord mesenchymal stem cells reduce microglia-mediated neuroinflammation in perinatal brain injury. *Stem Cell Res Ther.* 2019;10:105.
434. Zhang Y, Chopp M, Liu XS, et al. Exosomes derived from mesenchymal stromal cells promote axonal growth of cortical neurons. *Mol Neurobiol.* 2017;54:2659-2673.
435. Xiao Y, Geng F, Wang G, Li X, Zhu J, Zhu W. Bone marrow-derived mesenchymal stem cells-derived exosomes prevent oligodendrocyte apoptosis through exosomal miR-134 by targeting caspase-8. *J Cell Biochem.* 2019;120:2109-2118.
436. Xin H, Li Y, Liu Z, et al. MiR-133b promotes neural plasticity and functional recovery after treatment of stroke with multipotent mesenchymal stromal cells in rats via transfer of exosome-enriched extracellular particles. *Stem Cells.* 2013;31:2737-2746.
437. Joerger-Messerli MS, Oppliger B, Spinelli M, et al. Extracellular vesicles derived from Wharton's jelly mesenchymal stem cells prevent and resolve programmed cell death mediated by perinatal hypoxia-ischemia in neuronal cells. *Cell Transplant.* 2018;27:168-180.
438. Tran PH, Xiang D, Tran TT, et al. Exosomes and nanoengineering: a match made for precision therapeutics. *Adv Mater.* 2020;32:1904040.
439. Luan X, Sansanaphongpricha K, Myers I, Chen H, Yuan H, Sun D. Engineering exosomes as refined biological nanoplatforms for drug delivery. *Acta Pharmacol Sin.* 2017;38:754-763.
440. Xiong Y, Song J, Huang X, et al. Exosomes derived from mesenchymal stem cells: novel effects in the treatment of ischemic stroke. *Front Neurosci.* 2022;16:899887.
441. Webb RL, Kaiser EE, Scoville SL, et al. Human neural stem cell extracellular vesicles improve tissue and functional recovery in the murine thromboembolic stroke model. *Transl Stroke Res.* 2018;9:530-539.
442. Sun X, Jung J-H, Arvola O, et al. Stem cell-derived exosomes protect astrocyte cultures from in vitro ischemia and decrease injury as post-stroke intravenous therapy. *Front Cell Neurosci.* 2019;13:394.
443. Webb RL, Kaiser EE, Jurgielewicz BJ, et al. Human neural stem cell extracellular vesicles improve recovery in a porcine model of ischemic stroke. *Stroke.* 2018;49:1248-1256.
444. Hong P, Yang H, Wu Y, Li K, Tang Z. The functions and clinical application potential of exosomes derived from adipose mesenchymal stem cells: a comprehensive review. *Stem Cell Res Ther.* 2019;10:242.
445. Yang Y, Cai Y, Zhang Y, Liu J, Xu Z. Exosomes secreted by adipose-derived stem cells contribute to angiogenesis of brain microvascular endothelial cells following oxygen-glucose deprivation in vitro through microRNA-181b/TRPM7 axis. *J Mol Neurosci.* 2018;65:74-83.
446. Geng W, Tang H, Luo S, et al. Exosomes from miRNA-126-modified ADSCs promotes functional recovery after stroke in rats by improving neurogenesis and suppressing microglia activation. *Am J Transl Res.* 2019;11:780-792.
447. Jiang M, Wang H, Jin M, et al. Exosomes from MiR-30d-5p-ADSCs reverse acute ischemic stroke-induced, autophagy-mediated brain injury by promoting M2 microglial/macrophage polarization. *Cell Physiol Biochem.* 2018;47:864-878.
448. Huang X, Ding J, Li Y, et al. Exosomes derived from PEDF modified adipose-derived mesenchymal stem cells ameliorate cerebral ischemia-reperfusion injury by regulation of autophagy and apoptosis. *Exp Cell Res.* 2018;371:269-277.
449. Chen K-H, Chen C-H, Wallace CG, et al. Intravenous administration of xenogenic adipose-derived mesenchymal stem cells (ADMSC) and ADMSC-derived exosomes markedly reduced brain infarct volume and preserved neurological function in rat after acute ischemic stroke. *Oncotarget.* 2016;7:74537-74556.
450. Pei X, Li Y, Zhu L, Zhou Z. Astrocyte-derived exosomes suppress autophagy and ameliorate neuronal damage in experimental ischemic stroke. *Exp Cell Res.* 2019;382:111474.
451. Xu L, Cao H, Xie Y, et al. Exosome-shuttled miR-92b-3p from ischemic preconditioned astrocytes protects neurons against oxygen and glucose deprivation. *Brain Res.* 2019;1717:66-73.
452. Hira K, Ueno Y, Tanaka R, et al. Astrocyte-derived exosomes treated with a semaphorin 3A inhibitor enhance stroke recovery via prostaglandin D2 synthase. *Stroke.* 2018;49:2483-2494.
453. Pan Q, He C, Liu H, et al. Microvascular endothelial cells-derived microvesicles imply in ischemic stroke by modulating astrocyte and blood brain barrier function and cerebral blood flow. *Mol Brain.* 2016;9:63.
454. Venkat P, Cui C, Chopp M, et al. MiR-126 mediates brain endothelial cell exosome treatment-induced neurorestorative effects after stroke in type 2 diabetes mellitus mice. *Stroke.* 2019;50:2865-2874.
455. Zheng Y, He R, Wang P, Shi Y, Zhao L, Liang J. Exosomes from LPS-stimulated macrophages induce neuroprotection and functional improvement after ischemic stroke by modulating microglial polarization. *Biomater Sci.* 2019;7:2037-2049.
456. Song Y, Li Z, He T, et al. M2 microglia-derived exosomes protect the mouse brain from ischemia-reperfusion injury via exosomal miR-124. *Theranostics.* 2019;9:2910-2923.
457. Wang X, Wang J, Shi X, et al. Proteomic analyses identify a potential mechanism by which extracellular vesicles aggravate ischemic stroke. *Life Sci.* 2019;231:116527.
458. Xiao B, Chai Y, Lv S, et al. Endothelial cell-derived exosomes protect SH-SY5Y nerve cells against ischemia/reperfusion injury. *Int J Mol Med.* 2017;40:1201-1209.
459. Njock M-S, Cheng HS, Dang LT, et al. Endothelial cells suppress monocyte activation through secretion of extracellular vesicles containing antiinflammatory microRNAs. *Blood.* 2015;125:3202-3212.
460. Tian Y, Zhu P, Liu S, et al. IL-4-polarized BV2 microglia cells promote angiogenesis by secreting exosomes. *Adv Clin Exp Med.* 2019;28:421-430.
461. Wan T, Huang Y, Gao X, Wu W, Guo W. Microglia polarization: a novel target of exosome for stroke treatment. *Front Cell Dev Biol.* 2022;10:842320.
462. Zha Z, Gao Y-F, Ji J, et al. Bu Shen Yi sui capsule alleviates neuroinflammation and demyelination by promoting microglia toward M2 polarization, which correlates with changes in miR-124 and miR-155 in experimental autoimmune encephalomyelitis. *Oxidative Med Cell Longev.* 2021;2021:5521503.
463. Zhang H, Lin S, McElroy CL, et al. Circulating pro-inflammatory exosomes worsen stroke outcomes in aging. *Circ Res.* 2021;129:e121-e140.
464. Zang J, Wu Y, Su X, et al. Inhibition of PDE1-B by vinpocetine regulates microglial exosomes and polarization through enhancing autophagic flux for neuroprotection against ischemic stroke. *Front Cell Dev Biol.* 2021;8:616590.
465. Lin S, Yin Q, Zhong Q, et al. Heme activates TLR4-mediated inflammatory injury via MyD88/TRIF signaling pathway in intracerebral hemorrhage. *J Neuroinflammation.* 2012;9:46.
466. Zhang Q, Chen Z, Zhang Z, Jin Z, Muratoglu OK, Varadarajan KM. Leveraging subject-specific musculoskeletal modeling to assess effect of anterior cruciate ligament retaining total knee arthroplasty during walking gait. *Proc Inst Mech Eng H.* 2020;234:1445-1456.
467. Zhang Q, He L, Chen M, et al. PSD-93 mediates the crosstalk between neuron and microglia and facilitates acute ischemic stroke injury by binding to CX3CL1. *J Neurochem.* 2021;157:2145-2157.
468. Shu ZM, Shu XD, Li HQ, et al. Ginkgolide B protects against ischemic stroke via modulating microglia polarization in mice. *CNS Neurosci Ther.* 2016;22:729-739.
469. Zhang Z, Zou X, Zhang R, et al. Human umbilical cord mesenchymal stem cell-derived exosomal miR-146a-5p reduces microglial-mediated neuroinflammation via suppression of the IRAK1/TRAF6 signaling pathway after ischemic stroke. *Aging (Albany NY).* 2021;13:3060-3079.

470. Zhao B, Fei Y, Zhu J, Yin Q, Fang W, Li Y. PAF receptor inhibition attenuates neuronal pyroptosis in cerebral ischemia/reperfusion injury. *Mol Neurobiol*. 2021;58:6520-6539.
471. Zhang M, Gillaspay AF, Gipson JR, et al. Neuroinvasive listeria monocytogenes infection triggers IFN-activation of microglia and upregulates microglial miR-155. *Front Immunol*. 2018;9:2751.
472. Zhang R, Xu M, Wang Y, Xie F, Zhang G, Qin X. Nrf2—a promising therapeutic target for defending against oxidative stress in stroke. *Mol Neurobiol*. 2017;54:6006-6017.
473. Zhang Y, Hu Y-W, Zheng L, Wang Q. Characteristics and roles of exosomes in cardiovascular disease. *DNA Cell Biol*. 2017;36:202-211.
474. Zhang L, Lu X, Gong L, et al. Tetramethylpyrazine protects blood-spinal cord barrier integrity by modulating microglia polarization through activation of STAT3/SOCS3 and inhibition of NF- κ B signaling pathways in experimental autoimmune encephalomyelitis mice. *Cell Mol Neurobiol*. 2021;41:717-731.
475. Zhao S-c, Ma L-s, Chu Z-h, Xu H, Wu W-q, Liu F. Regulation of microglial activation in stroke. *Acta Pharmacol Sin*. 2017;38:445-458.
476. Lai CP, Mardini O, Ericsson M, et al. Dynamic biodistribution of extracellular vesicles in vivo using a multimodal imaging reporter. *ACS Nano*. 2014;8:483-494.
477. Betzer O, Perets N, Angel A, et al. In vivo neuroimaging of exosomes using gold nanoparticles. *ACS Nano*. 2017;11:10883-10893.
478. Sonoda T, Matsuzaki J, Yamamoto Y, et al. Serum microRNA-based risk prediction for stroke. *Stroke*. 2019;50:1510-1518.
479. Xu Y, Hu Y, Xu S, Liu F, Gao Y. Exosomal microRNAs as potential biomarkers and therapeutic agents for acute ischemic stroke: new expectations. *Front Neurol*. 2022;12:747380.
480. Hadem J, Tacke F, Bruns T, et al. Etiologies and outcomes of acute liver failure in Germany. *Clin Gastroenterol Hepatol*. 2012;10:664-669.e2.
481. Lee W. *Acute Liver Failure. Seminars in Respiratory and Critical Care Medicine*. Thieme Medical Publishers; 2012.
482. Blackmore L, Bernal W. Acute liver failure. *Clin Med*. 2015;15:468-472.
483. Patton H, Misel M, Gish RG. Acute liver failure in adults: an evidence-based management protocol for clinicians. *Gastroenterol Hepatol*. 2012;8:161-212.
484. Lee WM. Acute liver failure in the United States. *Seminars in Liver Disease*. Thieme Medical Publishers, Inc.; 2003.
485. Shokravi S, Borisov V, Zaman BA, et al. Mesenchymal stromal cells (MSCs) and their exosome in acute liver failure (ALF): a comprehensive review. *Stem Cell Res Ther*. 2022;13:192.
486. Yan Y, Jiang W, Tan Y, et al. hucMSC exosome-derived GPX1 is required for the recovery of hepatic oxidant injury. *Mol Ther*. 2017;25:465-479.
487. Jun JH, Kim JY, Choi JH, Lim J-Y, Kim K, Kim GJ. Exosomes from placenta-derived mesenchymal stem cells are involved in liver regeneration in hepatic failure induced by bile duct ligation. *Stem Cells Int*. 2020;2020:5485738.
488. Damania A, Jaiman D, Teotia AK, Kumar A. Mesenchymal stromal cell-derived exosome-rich fractionated secretome confers a hepatoprotective effect in liver injury. *Stem Cell Res Ther*. 2018;9:31.
489. Zhang S, Jiang L, Hu H, et al. Pretreatment of exosomes derived from hUCMSCs with TNF- α ameliorates acute liver failure by inhibiting the activation of NLRP3 in macrophage. *Life Sci*. 2020;246:117401.
490. Chen L, Xiang B, Wang X, Xiang C. Exosomes derived from human menstrual blood-derived stem cells alleviate fulminant hepatic failure. *Stem Cell Res Ther*. 2017;8:9.
491. Zheng J, Lu T, Zhou C, et al. Extracellular vesicles derived from human umbilical cord mesenchymal stem cells protect liver ischemia/reperfusion injury by reducing CD154 expression on CD4+ T cells via CCT2. *Adv Sci*. 2020;7:1903746.
492. Huang B, Cheng X, Wang H, et al. Mesenchymal stem cells and their secreted molecules predominantly ameliorate fulminant hepatic failure and chronic liver fibrosis in mice respectively. *J Transl Med*. 2016;14:45.
493. Parekkadan B, van Poll D, Suganuma K, et al. Mesenchymal stem cell-derived molecules reverse fulminant hepatic failure. *PLoS One*. 2007;2:e941.
494. Lotfinia M, Kadivar M, Piryaee A, et al. Effect of secreted molecules of human embryonic stem cell-derived mesenchymal stem cells on acute hepatic failure model. *Stem Cells Dev*. 2016;25:1898-1908.
495. Zhang J, Gao J, Lin D, et al. Potential networks regulated by MSCs in acute-on-chronic liver failure: exosomal miRNAs and intracellular target genes. *Front Genet*. 2021;12:650536.
496. Chen L, Zhang J, Yang L, Zhang G, Wang Y, Zhang S. The effects of conditioned medium derived from mesenchymal stem cells cocultured with hepatocytes on damaged hepatocytes and acute liver failure in rats. *Stem Cells Int*. 2018;2018:9156560.
497. Foroutan T, Kabiri F, Motamedi E. Silica magnetic graphene oxide improves the effects of stem cell-conditioned medium on acute liver failure. *ACS Omega*. 2021;6:21194-21206.
498. Zagoura DS, Roubelakis MG, Bitsika V, et al. Therapeutic potential of a distinct population of human amniotic fluid mesenchymal stem cells and their secreted molecules in mice with acute hepatic failure. *Gut*. 2012;61:894-906.
499. Lee WM, Larrey D, Olsson R, et al. Hepatic findings in long-term clinical trials of ximelagatran. *Drug Saf*. 2005;28:351-370.
500. Thakkar S, Li T, Liu Z, Wu L, Roberts R, Tong W. Drug-induced liver injury severity and toxicity (DILIst): binary classification of 1279 drugs by human hepatotoxicity. *Drug Discov Today*. 2020;25:201-208.
501. Walker PA, Ryder S, Lavado A, Dilworth C, Riley RJ. The evolution of strategies to minimise the risk of human drug-induced liver injury (DILI) in drug discovery and development. *Arch Toxicol*. 2020;94:2559-2585.
502. Sgro C, Clinard F, Ouazir K, et al. Incidence of drug-induced hepatic injuries: a French population-based study. *Hepatology*. 2002;36:451-455.
503. Kim KA. Current status of liver diseases in Korea: toxic and alcoholic liver diseases. *Korean J Hepatol*. 2009;15:S29-S33.
504. Li T, Yan Y, Wang B, et al. Exosomes derived from human umbilical cord mesenchymal stem cells alleviate liver fibrosis. *Stem Cells Dev*. 2013;22:845-854.
505. Tan CY, Lai RC, Wong W, Dan YY, Lim S-K, Ho HK. Mesenchymal stem cell-derived exosomes promote hepatic regeneration in drug-induced liver injury models. *Stem Cell Res Ther*. 2014;5:76.
506. Jiang W, Tan Y, Cai M, et al. Human umbilical cord MSC-derived exosomes suppress the development of CCl 4-induced liver injury through antioxidant effect. *Stem Cells Int*. 2018;2018:6079642.
507. Di Rocco G, Baldari S, Toietta G. Exosomes and other extracellular vesicles-mediated microRNA delivery for cancer therapy. *Transl Cancer Res*. 2017;6:S1321-S1330.
508. Villa F, Quarto R, Tasso R. Extracellular vesicles as natural, safe and efficient drug delivery systems. *Pharmaceutics*. 2019;11:557.
509. Fonsato V, Collino F, Herrera MB, et al. Human liver stem cell-derived microvesicles inhibit hepatoma growth in SCID mice by delivering antitumor microRNAs. *Stem Cells*. 2012;30:1985-1998.
510. Lou G, Song X, Yang F, et al. Exosomes derived from miR-122-modified adipose tissue-derived MSCs increase chemosensitivity of hepatocellular carcinoma. *J Hematol Oncol*. 2015;8:122.
511. Wang F, Li L, Piontek K, Sakaguchi M, Selaru FM. Exosome miR-335 as a novel therapeutic strategy in hepatocellular carcinoma. *Hepatology*. 2018;67:940-954.
512. Baldari S, Di Rocco G, Magenta A, Picozza M, Toietta G. Extracellular vesicles-encapsulated MicroRNA-125b produced in genetically modified mesenchymal stromal cells inhibits hepatocellular carcinoma cell proliferation. *Cells*. 2019;8:1560.

513. Ma B, Jiang H, Jia J, et al. Murine bone marrow stromal cells pulsed with homologous tumor-derived exosomes inhibit proliferation of liver cancer cells. *Clin Transl Oncol*. 2012;14:764-773.
514. Son SH, Gangadaran P, Ahn B-C. A novel strategy of transferring NIS protein to cells using extracellular vesicles leads to increase in iodine uptake and cytotoxicity. *Int J Nanomedicine*. 2019;14:1779-1787.
515. Chen W, Quan Y, Fan S, et al. Exosome-transmitted circular RNA hsa_circ_0051443 suppresses hepatocellular carcinoma progression. *Cancer Lett*. 2020;475:119-128.
516. Matsuda A, Ishiguro K, Yan IK, Patel T. Extracellular vesicle-based therapeutic targeting of β -catenin to modulate anticancer immune responses in hepatocellular cancer. *Hepatol Commun*. 2019;3:525-541.
517. Ishiguro K, Yan IK, Lewis-Tuffin L, Patel T. Targeting liver cancer stem cells using engineered biological nanoparticles for the treatment of hepatocellular cancer. *Hepatol Commun*. 2020;4:298-313.
518. Zhang J, Shan W-f, Jin T-t, et al. Propofol exerts anti-hepatocellular carcinoma by microvesicle-mediated transfer of miR-142-3p from macrophage to cancer cells. *J Transl Med*. 2014;12:279.
519. Rao Q, Zuo B, Lu Z, et al. Tumor-derived exosomes elicit tumor suppression in murine hepatocellular carcinoma models and humans in vitro. *Hepatology*. 2016;64:456-472.
520. Nimitrungtaewee N, Inmutto N, Chattipakorn SC, Chattipakorn N. Extracellular vesicles as a new hope for diagnosis and therapeutic intervention for hepatocellular carcinoma. *Cancer Med*. 2021;10:8253-8271.
521. Zhou L, Shen M, Fan X, Liu Y, Yang L. Pathogenic and potential therapeutic roles of exosomes derived from immune cells in liver diseases. *Front Immunol*. 2022;13:810300.
522. Bonventre JV, Yang L. Cellular pathophysiology of ischemic acute kidney injury. *J Clin Invest*. 2011;121:4210-4221.
523. Schäfer R, Spohn G, Baer PC. Mesenchymal stem/stromal cells in regenerative medicine: can preconditioning strategies improve therapeutic efficacy. *Transfus Med Hemother*. 2016;43:256-267.
524. Prockop DJ, Kota DJ, Bazhanov N, Reger RL. Evolving paradigms for repair of tissues by adult stem/progenitor cells (MSCs). *J Cell Mol Med*. 2010;14:2190-2199.
525. Squillaro T, Peluso G, Galderisi U. Clinical trials with mesenchymal stem cells: an update. *Cell Transplant*. 2016;25:829-848.
526. Fais S, O'Driscoll L, Borrás FE, et al. Evidence-based clinical use of nanoscale extracellular vesicles in nanomedicine. *ACS Nano*. 2016;10:3886-3899.
527. Oh S, Kwon S-H. Extracellular vesicles in acute kidney injury and clinical applications. *Int J Mol Sci*. 2021;22:8913.
528. Kosanović M, Milutinović B, Glamočlija S, Morlans IM, Ortiz A, Bozic M. Extracellular vesicles and acute kidney injury: potential therapeutic avenue for renal repair and regeneration. *Int J Mol Sci*. 2022;23:3792.
529. Bruno S, Grange C, Deregibus MC, et al. Mesenchymal stem cell-derived microvesicles protect against acute tubular injury. *J Am Soc Nephrol*. 2009;20:1053-1067.
530. Gatti S, Bruno S, Deregibus MC, et al. Microvesicles derived from human adult mesenchymal stem cells protect against ischaemia-reperfusion-induced acute and chronic kidney injury. *Nephrol Dial Transplant*. 2011;26:1474-1483.
531. Shen B, Liu J, Zhang F, et al. CCR2 positive exosome released by mesenchymal stem cells suppresses macrophage functions and alleviates ischemia/reperfusion-induced renal injury. *Stem Cells Int*. 2016;2016:1240301.
532. Bruno S, Grange C, Collino F, et al. Microvesicles derived from mesenchymal stem cells enhance survival in a lethal model of acute kidney injury. *PLoS One*. 2012;7:e33115.
533. Reis LA, Borges FT, Simoes MJ, Borges AA, Sinigaglia-Coimbra R, Schor N. Bone marrow-derived mesenchymal stem cells repaired but did not prevent gentamicin-induced acute kidney injury through paracrine effects in rats. *PLoS One*. 2012;7:e44092.
534. Collino F, Bruno S, Incarnato D, et al. AKI recovery induced by mesenchymal stromal cell-derived extracellular vesicles carrying microRNAs. *J Am Soc Nephrol*. 2015;26:2349-2360.
535. Zhu G, Pei L, Lin F, et al. Exosomes from human-bone-marrow-derived mesenchymal stem cells protect against renal ischemia/reperfusion injury via transferring miR-199a-3p. *J Cell Physiol*. 2019;234:23736-23749.
536. Zhou Y, Xu H, Xu W, et al. Exosomes released by human umbilical cord mesenchymal stem cells protect against cisplatin-induced renal oxidative stress and apoptosis in vivo and in vitro. *Stem Cell Res Ther*. 2013;4:34.
537. Zou X, Zhang G, Cheng Z, et al. Microvesicles derived from human Wharton's jelly mesenchymal stromal cells ameliorate renal ischemia-reperfusion injury in rats by suppressing CX3CL1. *Stem Cell Res Ther*. 2014;5:40.
538. Ju G-q, Cheng J, Zhong L, et al. Microvesicles derived from human umbilical cord mesenchymal stem cells facilitate tubular epithelial cell dedifferentiation and growth via hepatocyte growth factor induction. *PLoS One*. 2015;10:e0121534.
539. Gu D, Zou X, Ju G, Zhang G, Bao E, Zhu Y. Mesenchymal stromal cells derived extracellular vesicles ameliorate acute renal ischemia reperfusion injury by inhibition of mitochondrial fission through miR-30. *Stem Cells Int*. 2016;2016:2093940.
540. Zhang G, Zou X, Miao S, et al. The anti-oxidative role of microvesicles derived from human Wharton-Jelly mesenchymal stromal cells through NOX2/gp91 (phox) suppression in alleviating renal ischemia-reperfusion injury in rats. *PLoS One*. 2014;9:e92129.
541. Zhang G, Zou X, Huang Y, et al. Mesenchymal stromal cell-derived extracellular vesicles protect against acute kidney injury through anti-oxidation by enhancing Nrf2/ARE activation in rats. *Kidney Blood Press Res*. 2016;41:119-128.
542. Zou X, Gu D, Xing X, et al. Human mesenchymal stromal cell-derived extracellular vesicles alleviate renal ischemic reperfusion injury and enhance angiogenesis in rats. *Am J Transl Res*. 2016;8:4289-4299.
543. Wang B, Jia H, Zhang B, et al. Pre-incubation with hucMSC-exosomes prevents cisplatin-induced nephrotoxicity by activating autophagy. *Stem Cell Res Ther*. 2017;8:75.
544. Zhang Z-Y, Hou Y-P, Zou X-Y, et al. Oct-4 enhanced the therapeutic effects of mesenchymal stem cell-derived extracellular vesicles in acute kidney injury. *Kidney Blood Press Res*. 2020;45:95-108.
545. Cao J-Y, Wang B, Tang T-T, et al. Exosomal miR-125b-5p deriving from mesenchymal stem cells promotes tubular repair by suppression of p53 in ischemic acute kidney injury. *Theranostics*. 2021;11:5248-5266.
546. Lin K-C, Yip H-K, Shao P-L, et al. Combination of adipose-derived mesenchymal stem cells (ADMSC) and ADMSC-derived exosomes for protecting kidney from acute ischemia-reperfusion injury. *Int J Cardiol*. 2016;216:173-185.
547. Gao F, Zuo B, Wang Y, Li S, Yang J, Sun D. Protective function of exosomes from adipose tissue-derived mesenchymal stem cells in acute kidney injury through SIRT1 pathway. *Life Sci*. 2020;255:117719.
548. Liu Y, Cui J, Wang H, et al. Enhanced therapeutic effects of MSC-derived extracellular vesicles with an injectable collagen matrix for experimental acute kidney injury treatment. *Stem Cell Res Ther*. 2020;11:161.
549. Zhang C, Shang Y, Chen X, et al. Supramolecular nanofibers containing arginine-glycine-aspartate (RGD) peptides boost therapeutic efficacy of extracellular vesicles in kidney repair. *ACS Nano*. 2020;14:12133-12147.
550. Choi HY, Moon SJ, Ratliff BB, et al. Microparticles from kidney-derived mesenchymal stem cells act as carriers of proangiogenic

- signals and contribute to recovery from acute kidney injury. *PLoS One*. 2014;9:e87853.
551. Ranghino A, Bruno S, Bussolati B, et al. The effects of glomerular and tubular renal progenitors and derived extracellular vesicles on recovery from acute kidney injury. *Stem Cell Res Ther*. 2017;8:24.
 552. Sanchez MBH, Bruno S, Grange C, et al. Human liver stem cells and derived extracellular vesicles improve recovery in a murine model of acute kidney injury. *Stem Cell Res Ther*. 2014;5:124.
 553. Grange C, Papadimitriou E, Dimuccio V, et al. Urinary extracellular vesicles carrying klotho improve the recovery of renal function in an acute tubular injury model. *Mol Ther*. 2020;28:490-502.
 554. Du J, Sun Q, Wang Z, et al. Tubular epithelial cells derived-exosomes containing CD26 protects mice against renal ischemia/reperfusion injury by maintaining proliferation and dissipating inflammation. *Biochem Biophys Res Commun*. 2021;553:134-140.
 555. Li X, Liao J, Su X, et al. Human urine-derived stem cells protect against renal ischemia/reperfusion injury in a rat model via exosomal miR-146a-5p which targets IRAK1. *Theranostics*. 2020;10:9561-9578.
 556. Tang T-T, Wang B, Wu M, et al. Extracellular vesicle-encapsulated IL-10 as novel nanotherapeutics against ischemic AKI. *Sci Adv*. 2020;6:eaz0748.
 557. Cantaluppi V, Gatti S, Medica D, et al. Microvesicles derived from endothelial progenitor cells protect the kidney from ischemia-reperfusion injury by microRNA-dependent reprogramming of resident renal cells. *Kidney Int*. 2012;82:412-427.
 558. He Z, Wang H, Yue L. Endothelial progenitor cells-secreted extracellular vesicles containing microRNA-93-5p confer protection against sepsis-induced acute kidney injury via the KDM6B/H3K27me3/TNF- α axis. *Exp Cell Res*. 2020;395:112173.
 559. Gregorini M, Corradetti V, Pattonieri EF, et al. Perfusion of isolated rat kidney with mesenchymal stromal cells/extracellular vesicles prevents ischaemic injury. *J Cell Mol Med*. 2017;21:3381-3393.
 560. Nassar W, El-Ansary M, Sabry D, et al. Umbilical cord mesenchymal stem cells derived extracellular vesicles can safely ameliorate the progression of chronic kidney diseases. *Biomater Res*. 2016;20:21.
 561. Webster AC, Nagler EV, Morton RL, Masson P. Chronic kidney disease. *Lancet*. 2017;389:1238-1252.
 562. Duffield JS. Cellular and molecular mechanisms in kidney fibrosis. *J Clin Invest*. 2014;124:2299-2306.
 563. Gregorini M, Corradetti V, Rocca C, et al. Mesenchymal stromal cells prevent renal fibrosis in a rat model of unilateral ureteral obstruction by suppressing the renin-angiotensin system via HuR. *PLoS One*. 2016;11:e0148542.
 564. Gregorini M, Maccario R, Avanzini MA, et al. Antineutrophil cytoplasmic antibody-associated renal vasculitis treated with autologous mesenchymal stromal cells: evaluation of the contribution of immune-mediated mechanisms. *Mayo Clinic Proceedings*. Vol. 88. Elsevier; 2013:1174-1179.
 565. Reidy K, Kang HM, Hostetter T, Susztak K. Molecular mechanisms of diabetic kidney disease. *J Clin Invest*. 2014;124:2333-2340.
 566. Badal SS, Danesh FR. New insights into molecular mechanisms of diabetic kidney disease. *Am J Kidney Dis*. 2014;63:S63-S83.
 567. Cao Q, Huang C, Chen X-M, Pollock CA. Mesenchymal stem cell-derived exosomes: toward cell-free therapeutic strategies in chronic kidney disease. *Front Med*. 2022;9:816656.
 568. Filomeni G, De Zio D, Cecconi F. Oxidative stress and autophagy: the clash between damage and metabolic needs. *Cell Death Differ*. 2015;22:377-388.
 569. Ebrahim N, Ahmed IA, Hussien NI, et al. Mesenchymal stem cell-derived exosomes ameliorated diabetic nephropathy by autophagy induction through the mTOR signaling pathway. *Cells*. 2018;7:226.
 570. Daehn IS, Duffield JS. The glomerular filtration barrier: a structural target for novel kidney therapies. *Nat Rev Drug Discov*. 2021;20:770-788.
 571. Kasztan M, Pollock DM. Impact of ET-1 and sex in glomerular hyperfiltration in humanized sickle cell mice. *Clin Sci*. 2019;133:1475-1486.
 572. Nagaishi K, Mizue Y, Chikenji T, et al. Mesenchymal stem cell therapy ameliorates diabetic nephropathy via the paracrine effect of renal trophic factors including exosomes. *Sci Rep*. 2016;6:34842.
 573. Ahlborg G, Lindström J. Insulin sensitivity and big ET-1 conversion to ET-1 after ETA-or ETB-receptor blockade in humans. *J Appl Physiol*. 2002;93:2112-2121.
 574. Hao Y, Miao J, Liu W, Cai K, Huang X, Peng L. Mesenchymal stem cell-derived exosomes carry MicroRNA-125a to protect against diabetic nephropathy by targeting histone deacetylase 1 and downregulating endothelin-1. *Diabetes Metab Syndr Obes*. 2021;14:1405-1418.
 575. Meng X-m, Nikolic-Paterson DJ, Lan HY. TGF- β : the master regulator of fibrosis. *Nat Rev Nephrol*. 2016;12:325-338.
 576. Martínez-Klimova E, Aparicio-Trejo OE, Tapia E, Pedraza-Chaverri J. Unilateral ureteral obstruction as a model to investigate fibrosis-attenuating treatments. *Biomolecules*. 2019;9:141.
 577. Ratliff BB, Abdulmahdi W, Pawar R, Wolin MS. Oxidant mechanisms in renal injury and disease. *Antioxid Redox Signal*. 2016;25:119-146.
 578. Ishiy CSRA, Ormanji MS, Maquigussa E, Ribeiro RS, da Silva NA, Boim MA. Comparison of the effects of mesenchymal stem cells with their extracellular vesicles on the treatment of kidney damage induced by chronic renal artery stenosis. *Stem Cells Int*. 2020;2020:8814574.
 579. Dendooven A, Ishola DA Jr, Nguyen TQ, et al. Oxidative stress in obstructive nephropathy. *Int J Exp Pathol*. 2011;92:202-210.
 580. Jin J, Qian F, Zheng D, He W, Gong J, He Q. Mesenchymal stem cells attenuate renal fibrosis via exosomes-mediated delivery of microRNA let-7i-5p antagomir. *Int J Nanomedicine*. 2021;16:3565-3578.
 581. Ji C, Zhang J, Zhu Y, et al. Exosomes derived from hucMSC attenuate renal fibrosis through CK1 δ / β -TRCP-mediated YAP degradation. *Cell Death Dis*. 2020;11:327.
 582. Kumar S, Liu J, Pang P, et al. Sox9 activation highlights a cellular pathway of renal repair in the acutely injured mammalian kidney. *Cell Rep*. 2015;12:1325-1338.
 583. Gao M, Yu Z, Yao D, Qian Y, Wang Q, Jia R. Mesenchymal stem cells therapy: a promising method for the treatment of uterine scars and premature ovarian failure. *Tissue Cell*. 2022;74:101676.
 584. Yang M, Lin L, Sha C, et al. Bone marrow mesenchymal stem cell-derived exosomal miR-144-5p improves rat ovarian function after chemotherapy-induced ovarian failure by targeting PTEN. *Lab Invest*. 2020;100:342-352.
 585. Tan Q, Xia D, Ying X. miR-29a in exosomes from bone marrow mesenchymal stem cells inhibit fibrosis during endometrial repair of intrauterine adhesion. *Int J Stem Cells*. 2020;13:414-423.
 586. Xiao B, Zhu Y, Huang J, Wang T, Wang F, Sun S. Exosomal transfer of bone marrow mesenchymal stem cell-derived miR-340 attenuates endometrial fibrosis. *Biol Open*. 2019;8:bio039958.
 587. Cui X, Zhu L, Zhai R, Zhang B, Zhang F. Mesenchymal stem cell-derived exosomes: a promising vector in treatment for diabetes and its microvascular complications. *Am J Transl Res*. 2021;13:3942-3953.
 588. Fluit MB, Mohit N, Gambhir KK, Nunlee-Bland G. To the future: the role of exosome-derived microRNAs as markers, mediators, and therapies for endothelial dysfunction in type 2 diabetes mellitus. *J Diabetes Res*. 2022;2022:5126968.
 589. Zheng Y, Ley SH, Hu FB. Global aetiology and epidemiology of type 2 diabetes mellitus and its complications. *Nat Rev Endocrinol*. 2018;14:88-98.
 590. Chatterjee S, Khunti K, Davies MJ. Type 2 diabetes. *Lancet*. 2017;389:2239-2251.
 591. Zhang Y, Liu Y, Liu H, Tang WH. Exosomes: biogenesis, biologic function and clinical potential. *Cell Biosci*. 2019;9:19.

592. Zampetaki A, Kiechl S, Drozdov I, et al. Plasma microRNA profiling reveals loss of endothelial miR-126 and other microRNAs in type 2 diabetes. *Circ Res*. 2010;107:810-817.
593. Wang S, Aurora AB, Johnson BA, et al. The endothelial-specific microRNA miR-126 governs vascular integrity and angiogenesis. *Dev Cell*. 2008;15:261-271.
594. Jansen F, Yang X, Hoelscher M, et al. Endothelial microparticle-mediated transfer of microRNA-126 promotes vascular endothelial cell repair via SPRED1 and is abrogated in glucose-damaged endothelial microparticles. *Circulation*. 2013;128:2026-2038.
595. Jansen F, Wang H, Przybilla D, et al. Vascular endothelial microparticles-incorporated microRNAs are altered in patients with diabetes mellitus. *Cardiovasc Diabetol*. 2016;15:49.
596. Eissa S, Matboli M, Bekhet MM. Clinical verification of a novel urinary microRNA panel: 133b, -342 and -30 as biomarkers for diabetic nephropathy identified by bioinformatics analysis. *Biomed Pharmacother*. 2016;83:92-99.
597. Shao X, Kong W-X, Li Y-T. MiR-133 inhibits kidney injury in rats with diabetic nephropathy via MAPK/ERK pathway. *Eur Rev Med Pharmacol Sci*. 2019;23:10957-10963.
598. Jiang ZH, Tang YZ, Song HN, Yang M, Li B, Ni CL. miRNA-342 suppresses renal interstitial fibrosis in diabetic nephropathy by targeting SOX6. *Int J Mol Med*. 2020;45:45-52.
599. Yang X, Yang M, Chen Y, et al. miR-30a-5p targets Becn1 to ameliorate high-glucose-induced glomerular podocyte injury in immortalized rat podocyte cell line. *Am J Transl Res*. 2021;13:1516-1525.
600. Valadi H, Ekström K, Bossios A, Sjöstrand M, Lee JJ, Lötvald JO. Exosome-mediated transfer of mRNAs and microRNAs is a novel mechanism of genetic exchange between cells. *Nat Cell Biol*. 2007;9:654-659.
601. Katayama M, Wiklander OP, Fritz T, et al. Circulating exosomal miR-20b-5p is elevated in type 2 diabetes and could impair insulin action in human skeletal muscle. *Diabetes*. 2019;68:515-526.
602. Olivieri F, Spazzafumo L, Bonafè M, et al. MiR-21-5p and miR-126a-3p levels in plasma and circulating angiogenic cells: relationship with type 2 diabetes complications. *Oncotarget*. 2015;6:35372-35382.
603. Zuo K, Li M, Zhang X, et al. MiR-21 suppresses endothelial progenitor cell proliferation by activating the TGFβ signaling pathway via downregulation of WWP1. *Int J Clin Exp Pathol*. 2015;8:414-422.
604. Fu Q, Jiang H, Wang Z, et al. Injury factors alter miRNAs profiles of exosomes derived from islets and circulation. *Aging (Albany NY)*. 2018;10:3986-3999.
605. Li M, Liu Q, Lei J, Wang X, Chen X, Ding Y. MiR-362-3p inhibits the proliferation and migration of vascular smooth muscle cells in atherosclerosis by targeting ADAMTS1. *Biochem Biophys Res Commun*. 2017;493:270-276.
606. Fluit MB, Kumari N, Nunlee-Bland G, Nekhai S, Gambhir KK. miRNA-15a, miRNA-15b, and miRNA-499 are reduced in erythrocytes of pre-diabetic African-American adults. *Jacobs J Diabetes Endocrinol*. 2016;2:014.
607. Kamalden TA, Macgregor-Das AM, Kannan SM, et al. Exosomal microRNA-15a transfer from the pancreas augments diabetic complications by inducing oxidative stress. *Antioxid Redox Signal*. 2017;27:913-930.
608. Wan S, Wang J, Wang J, et al. Increased serum miR-7 is a promising biomarker for type 2 diabetes mellitus and its microvascular complications. *Diabetes Res Clin Pract*. 2017;130:171-179.
609. Santovito D, Toto L, De Nardis V, et al. Plasma microRNA signature associated with retinopathy in patients with type 2 diabetes. *Sci Rep*. 2021;11:4136.
610. Li Q, Kim Y-R, Vikram A, et al. P66Shc-induced microRNA-34a causes diabetic endothelial dysfunction by downregulating Sirtuin1. *Arterioscler Thromb Vasc Biol*. 2016;36:2394-2403.
611. Yin C, Lin X, Sun Y, Ji X. Dysregulation of miR-210 is involved in the development of diabetic retinopathy and serves a regulatory role in retinal vascular endothelial cell proliferation. *Eur J Med Res*. 2020;25:20.
612. Zhou Z, Collado A, Sun C, et al. Downregulation of erythrocyte miR-210 induces endothelial dysfunction in type 2 diabetes. *Diabetes*. 2022;71:285-297.
613. Luo Y-F, Wan X-X, Zhao L-L, et al. MicroRNA-139-5p upregulation is associated with diabetic endothelial cell dysfunction by targeting c-jun. *Aging (Albany NY)*. 2021;13:1186-1211.
614. Mohan A, Singh RS, Kumari M, et al. Urinary exosomal microRNA-451-5p is a potential early biomarker of diabetic nephropathy in rats. *PLoS One*. 2016;11:e0154055.
615. Fluit MB, Shivapurkar N, Kumari M, et al. Systemic inhibition of miR-451 increases fibrotic signaling and diminishes autophagic response to exacerbate renal damage in tattyho/Jng mice. *Am J Physiol Renal Physiol*. 2020;319:F476-F486.
616. Kumari M, Mohan A, Ecelbarger CM, et al. miR-451 loaded exosomes are released by the renal cells in response to injury and associated with reduced kidney function in human. *Front Physiol*. 2020;11:234.
617. Li W, Yang S, Qiao R, Zhang J. Potential value of urinary exosome-derived let-7c-5p in the diagnosis and progression of type II diabetic nephropathy. *Clin Lab*. 2018;64:709-718.
618. Jia Y, Guan M, Zheng Z, et al. miRNAs in urine extracellular vesicles as predictors of early-stage diabetic nephropathy. *J Diabetes Res*. 2016;2016:7932765.
619. Liu F, Zhang Z-P, Xin G-D, Guo L-H, Jiang Q, Wang Z-X. miR-192 prevents renal tubulointerstitial fibrosis in diabetic nephropathy by targeting Egr1. *Eur Rev Med Pharmacol Sci*. 2018;22:4252-4260.
620. Zang J, Maxwell AP, Simpson DA, McKay GJ. Differential expression of urinary exosomal microRNAs miR-21-5p and miR-30b-5p in individuals with diabetic kidney disease. *Sci Rep*. 2019;9:10900.
621. Lin Z, Ge J, Wang Z, et al. Let-7e modulates the inflammatory response in vascular endothelial cells through ceRNA crosstalk. *Sci Rep*. 2017;7:42498.
622. Kong L, Du X, Hu N, et al. Downregulation of let-7e-5p contributes to endothelial progenitor cell dysfunction in deep vein thrombosis via targeting FASLG. *Thromb Res*. 2016;138:30-36.
623. Eissa S, Matboli M, Aboushabha R, Bekhet MM, Soliman Y. Urinary exosomal microRNA panel unravels novel biomarkers for diagnosis of type 2 diabetic kidney disease. *J Diabetes Complicat*. 2016;30:1585-1592.
624. Zhang J, Li S, Li L, et al. Exosome and exosomal microRNA: trafficking, sorting, and function. *Genom Proteom Bioinform*. 2015;13:17-24.
625. Barile L, Moccetti T, Marbán E, Vassalli G. Roles of exosomes in cardioprotection. *Eur Heart J*. 2017;38:1372-1379.
626. Delić D, Eisele C, Schmid R, et al. Urinary exosomal miRNA signature in type II diabetic nephropathy patients. *PLoS One*. 2016;11:e0150154.
627. Prabu P, Rome S, Sathishkumar C, et al. MicroRNAs from urinary extracellular vesicles are non-invasive early biomarkers of diabetic nephropathy in type 2 diabetes patients with the 'Asian Indian phenotype'. *Diabetes Metab*. 2019;45:276-285.
628. Sun Y, Tao Q, Wu X, Zhang L, Liu Q, Wang L. The utility of exosomes in diagnosis and therapy of diabetes mellitus and associated complications. *Front Endocrinol*. 2021;12:756581.
629. Sabatier F, Darmon P, Hugel B, et al. Type 1 and type 2 diabetic patients display different patterns of cellular microparticles. *Diabetes*. 2002;51:2840-2845.
630. Li S, Wei J, Zhang C, et al. Cell-derived microparticles in patients with type 2 diabetes mellitus: a systematic review and meta-analysis. *Cell Physiol Biochem*. 2016;39:2439-2450.
631. Burger D, Turner M, Xiao F, Munkonda MN, Akbari S, Burns KD. High glucose increases the formation and pro-oxidative activity of endothelial microparticles. *Diabetologia*. 2017;60:1791-1800.

632. Duan Y, Luo Q, Wang Y, et al. Adipose mesenchymal stem cell-derived extracellular vesicles containing microRNA-26a-5p target TLR4 and protect against diabetic nephropathy. *J Biol Chem*. 2020;295:12868-12884.
633. Sun A-l, Deng J-t, Guan G-j, et al. Dipeptidyl peptidase-IV is a potential molecular biomarker in diabetic kidney disease. *Diab Vasc Dis Res*. 2012;9:301-308.
634. Abe H, Sakurai A, Ono H, et al. Urinary exosomal mRNA of WT1 as diagnostic and prognostic biomarker for diabetic nephropathy. *J Med Invest*. 2018;65:208-215.
635. Zubiri I, Posada-Ayala M, Sanz-Maroto A, et al. Diabetic nephropathy induces changes in the proteome of human urinary exosomes as revealed by label-free comparative analysis. *J Proteome*. 2014;96:92-102.
636. Barutta F, Tricarico M, Corbelli A, et al. Urinary exosomal microRNAs in incipient diabetic nephropathy. *PLoS One*. 2013;8:e73798.
637. Sharma K, Karl B, Mathew AV, et al. Metabolomics reveals signature of mitochondrial dysfunction in diabetic kidney disease. *J Am Soc Nephrol*. 2013;24:1901-1912.
638. Sakurai A, Ono H, Ochi A, et al. Involvement of Elf3 on Smad3 activation-dependent injuries in podocytes and excretion of urinary exosome in diabetic nephropathy. *PLoS One*. 2019;14:e0216788.
639. Lange T, Stracke S, Rettig R, et al. Identification of miR-16 as an endogenous reference gene for the normalization of urinary exosomal miRNA expression data from CKD patients. *PLoS One*. 2017;12:e0183435.
640. Gudehithlu KP, Garcia-Gomez I, Vernik J, et al. In diabetic kidney disease urinary exosomes better represent kidney specific protein alterations than whole urine. *Am J Nephrol*. 2016;42:418-424.
641. Davidson SM, Riquelme JA, Takov K, et al. Cardioprotection mediated by exosomes is impaired in the setting of type II diabetes but can be rescued by the use of non-diabetic exosomes in vitro. *J Cell Mol Med*. 2018;22:141-151.
642. Wang X, Gu H, Huang W, et al. Hsp20-mediated activation of exosome biogenesis in cardiomyocytes improves cardiac function and angiogenesis in diabetic mice. *Diabetes*. 2016;65:3111-3128.
643. Wang X, Huang W, Liu G, et al. Cardiomyocytes mediate anti-angiogenesis in type 2 diabetic rats through the exosomal transfer of miR-320 into endothelial cells. *J Mol Cell Cardiol*. 2014;74:139-150.
644. Prattichizzo F, Giuliani A, De Nigris V, et al. Extracellular microRNAs and endothelial hyperglycaemic memory: a therapeutic opportunity? *Diabetes Obes Metab*. 2016;18:855-867.
645. Schara K, Štukelj R, Krek J, et al. A study of extracellular vesicle concentration in active diabetic Charcot neuroarthropathy. *Eur J Pharm Sci*. 2017;98:58-63.
646. Ju R, Zhuang ZW, Zhang J, et al. Angiopoietin-2 secretion by endothelial cell exosomes: regulation by the phosphatidylinositol 3-kinase (PI3K)/Akt/endothelial nitric oxide synthase (eNOS) and syndecan-4/syntenin pathways. *J Biol Chem*. 2014;289:510-519.
647. Nair S, Jayabalan N, Guanzon D, et al. Human placental exosomes in gestational diabetes mellitus carry a specific set of miRNAs associated with skeletal muscle insulin sensitivity. *Clin Sci*. 2018;132:2451-2467.
648. Smith SM, Boppana A, Traupman JA, et al. Impaired glucose metabolism in patients with diabetes, prediabetes, and obesity is associated with severe COVID-19. *J Med Virol*. 2021;93:409-415.
649. Song L, Yuan J, Liu Y, et al. Ghrelin system is involved in improvements in glucose metabolism mediated by hyperbaric oxygen treatment in a streptozotocin-induced type 1 diabetes mouse model. *Mol Med Rep*. 2020;22:3767-3776.
650. Miras AD, Kamocka A, Pérez-Pevida B, et al. Diabetes Care Publish Ahead of Print, published online November, 6, 2020; 2020.
651. Zhang L, Chu W, Zheng L, et al. Zinc oxide nanoparticles from *Cyperus rotundus* attenuates diabetic retinopathy by inhibiting NLRP3 inflammasome activation in STZ-induced diabetic rats. *J Biochem Mol Toxicol*. 2020;34:e22583.
652. Wysham C, Shubrook J. Beta-cell failure in type 2 diabetes: mechanisms, markers, and clinical implications. *Postgrad Med*. 2020;132:676-686.
653. Xiao X, Liang S, Zhao Y, et al. Core-shell structured 5-FU@ ZIF-90@ ZnO as a biodegradable nanopatform for synergistic cancer therapy. *Nanoscale*. 2020;12:3846-3854.
654. Tschen S-l, Dhawan S, Gurlo T, Bhushan A. Age-dependent decline in β -cell proliferation restricts the capacity of β -cell regeneration in mice. *Diabetes*. 2009;58:1312-1320.
655. Atkinson MA, Eisenbarth GS, Michels AW. Type 1 diabetes. *Lancet*. 2014;383:69-82.
656. Löhr M, Klöppel G. Residual insulin positivity and pancreatic atrophy in relation to duration of chronic type 1 (insulin-dependent) diabetes mellitus and microangiopathy. *Diabetologia*. 1987;30:757-762.
657. Meier J, Bhushan A, Butler A, Rizza R, Butler P. Sustained beta cell apoptosis in patients with long-standing type 1 diabetes: indirect evidence for islet regeneration? *Diabetologia*. 2005;48:2221-2228.
658. Mastrototaro L, Roden M. Insulin resistance and insulin sensitizing agents. *Metabolism*. 2021;125:154892.
659. Sędzikowska A, Szablewski L. Insulin and insulin resistance in Alzheimer's disease. *Int J Mol Sci*. 2021;22:9987.
660. Galli M, Hameed A, Żbikowski A, Zabielski P. Aquaporins in insulin resistance and diabetes: more than channels! *Redox Biol*. 2021;44:102027.
661. Zhu Q, Zhu R, Jin J. Neutral ceramidase-enriched exosomes prevent palmitic acid-induced insulin resistance in H4 IIEC 3 hepatocytes. *FEBS Open Bio*. 2016;6:1078-1084.
662. Chen M-T, Zhao Y-t, Zhou L-y, et al. Exosomes derived from human umbilical cord mesenchymal stem cells enhance insulin sensitivity in insulin resistant human adipocytes. *Curr Med Sci*. 2021;41:87-93.
663. Liu J, Jiang F, Jiang Y, et al. Roles of exosomes in ocular diseases. *Int J Nanomedicine*. 2020;15:10519-10538.
664. Tellez-Gabriel M, Knutsen E, Perander M. Current status of circulating tumor cells, circulating tumor DNA, and exosomes in breast cancer liquid biopsies. *Int J Mol Sci*. 2020;21:9457.
665. Ashrafizadeh M, Kumar AP, Aref AR, Zarrabi A, Mostafavi E. Exosomes as promising nanostructures in diabetes mellitus: from insulin sensitivity to ameliorating diabetic complications. *Int J Nanomedicine*. 2022;17:1229-1253.
666. Ji Y, Luo Z, Gao H, et al. Hepatocyte-derived exosomes from early onset obese mice promote insulin sensitivity through miR-3075. *Nat Metab*. 2021;3:1163-1174.
667. Sun Y, Shi H, Yin S, et al. Human mesenchymal stem cell derived exosomes alleviate type 2 diabetes mellitus by reversing peripheral insulin resistance and relieving β -cell destruction. *ACS Nano*. 2018;12:7613-7628.
668. Li F, Li H, Jin X, et al. Adipose-specific knockdown of Sirt1 results in obesity and insulin resistance by promoting exosomes release. *Cell Cycle*. 2019;18:2067-2082.
669. Song M, Han L, Chen F-f, et al. Adipocyte-derived exosomes carrying sonic hedgehog mediate M1 macrophage polarization-induced insulin resistance via Ptch and PI3K pathways. *Cell Physiol Biochem*. 2018;48:1416-1432.
670. Chen J, Chen J, Cheng Y, et al. Mesenchymal stem cell-derived exosomes protect beta cells against hypoxia-induced apoptosis via miR-21 by alleviating ER stress and inhibiting p38 MAPK phosphorylation. *Stem Cell Res Ther*. 2020;11:97.
671. Qian B, Yang Y, Tang N, et al. M1 macrophage-derived exosomes impair beta cell insulin secretion via miR-212-5p by targeting SIRT2 and inhibiting Akt/GSK-3 β / β -catenin pathway in mice. *Diabetologia*. 2021;64:2037-2051.
672. He Q, Wang L, Zhao R, et al. Mesenchymal stem cell-derived exosomes exert ameliorative effects in type 2 diabetes by improving

- hepatic glucose and lipid metabolism via enhancing autophagy. *Stem Cell Res Ther*. 2020;11:223.
673. Salazar JJ, Ennis WJ, Koh TJ. Diabetes medications: impact on inflammation and wound healing. *J Diabetes Complicat*. 2016;30:746-752.
 674. Eming SA, Martin P, Tomic-Canic M. Wound repair and regeneration: mechanisms, signaling, and translation. *Sci Transl Med*. 2014;6:265sr6.
 675. Mirza RE, Fang MM, Weinheimer-Haus EM, Ennis WJ, Koh TJ. Sustained inflammasome activity in macrophages impairs wound healing in type 2 diabetic humans and mice. *Diabetes*. 2014;63:1103-1114.
 676. Kalantari K, Mostafaei A, Afifi AM, et al. Wound dressings functionalized with silver nanoparticles: promises and pitfalls. *Nanoscale*. 2020;12:2268-2291.
 677. Falanga V. Wound healing and its impairment in the diabetic foot. *Lancet*. 2005;366:1736-1743.
 678. Reiber GE, Raugi GJ. Preventing foot ulcers and amputations in diabetes. *Lancet*. 2005;366:1676-1677.
 679. Bailey AJ, Li H, Kirkham AM, et al. MSC-derived extracellular vesicles to heal diabetic wounds: a systematic review and meta-analysis of preclinical animal studies. *Stem Cell Rev Rep*. 2022;18:968-979.
 680. Pop-Busui R, Boulton AJ, Feldman EL, et al. Diabetic neuropathy: a position statement by the American Diabetes Association. *Diabetes Care*. 2017;40:136-154.
 681. Gordo A, Scuffham P, Shearer A, Oglesby A, Tobian JA. The health care costs of diabetic peripheral neuropathy in the US. *Diabetes Care*. 2003;26:1790-1795.
 682. Yaribeygi H, Ashrafizadeh M, Henney NC, Sathyapalan T, Jamialahmadi T, Sahebkar A. Neuromodulatory effects of anti-diabetes medications: a mechanistic review. *Pharmacol Res*. 2020;152:104611.
 683. Zhu Y, Han S, Li X, et al. Paeoniflorin effect of Schwann cell-derived exosomes ameliorates dorsal root ganglion neurons apoptosis through IRE1 α pathway. *Evid Based Complement Alternat Med*. 2021;2021:6079305.
 684. Fan B, Chopp M, Zhang ZG, Liu XS. Treatment of diabetic peripheral neuropathy with engineered mesenchymal stromal cell-derived exosomes enriched with microRNA-146a provide amplified therapeutic efficacy. *Exp Neurol*. 2021;341:113694.
 685. Azushima K, Gurley SB, Coffman TM. Modelling diabetic nephropathy in mice. *Nat Rev Nephrol*. 2018;14:48-56.
 686. Nagib AM, Elsayed Matter Y, Gheith OA, Refaie AF, Othman NF, Al-Otaibi T. Diabetic nephropathy following posttransplant diabetes mellitus. *Exp Clin Transplant*. 2019;17:138-146.
 687. Cai X, Zou F, Xuan R, Lai X-Y. Exosomes from mesenchymal stem cells expressing microribonucleic acid-125b inhibit the progression of diabetic nephropathy via the tumour necrosis factor receptor-associated factor 6/Akt axis. *Endocr J*. 2021;68:817-828.
 688. Ashrafizadeh M, Zarrabi A, Orouei S, et al. MicroRNA-mediated autophagy regulation in cancer therapy: the role in chemoresistance/chemosensitivity. *Eur J Pharmacol*. 2021;892:173660.
 689. Ashrafizadeh M, Ahmadi Z, Farkhondeh T, Samarghandian S. Modulatory effects of statins on the autophagy: a therapeutic perspective. *J Cell Physiol*. 2020;235:3157-3168.
 690. Ashrafizadeh M, Tavakoli S, Ahmadi Z, Roomiani S, Mohammadinejad R, Samarghandian S. Therapeutic effects of kaempferol affecting autophagy and endoplasmic reticulum stress. *Phytother Res*. 2020;34:911-923.
 691. Mohammadinejad R, Ahmadi Z, Tavakoli S, Ashrafizadeh M. Berberine as a potential autophagy modulator. *J Cell Physiol*. 2019;234:14914-14926.
 692. Ashrafizadeh M, Ahmadi Z, Farkhondeh T, Samarghandian S. Autophagy as a molecular target of quercetin underlying its protective effects in human diseases. *Arch Physiol Biochem*. 2022;128:200-208.
 693. Wang D, He X, Wang D, et al. Quercetin suppresses apoptosis and attenuates intervertebral disc degeneration via the SIRT1-autophagy pathway. *Front Cell Dev Biol*. 2020;8:613006.
 694. Sultan SA, Liu W, Peng Y, Roberts W, Whitelaw D, Graham AM. The role of maternal gestational diabetes in inducing fetal endothelial dysfunction. *J Cell Physiol*. 2015;230:2695-2705.
 695. Sobrevia L, Salsoso R, Sáez T, Sanhueza C, Pardo F, Leiva A. Insulin therapy and fetoplacental vascular function in gestational diabetes mellitus. *Exp Physiol*. 2015;100:231-238.
 696. Leiva Mendoza AA, Fuenzalida Saavedra B, Barros Lamus ER, et al. Nitric oxide is a central common metabolite in vascular dysfunction associated with diseases of human pregnancy. *Curr Vasc Pharmacol*. 2016;14:237-259.
 697. Bai S, Yin Q, Dong T, et al. Endothelial progenitor cell-derived exosomes ameliorate endothelial dysfunction in a mouse model of diabetes. *Biomed Pharmacother*. 2020;131:110756.
 698. Giugliano F, Maiorino M, Bellastella G, Gicchino M, Giugliano D, Esposito K. Determinants of erectile dysfunction in type 2 diabetes. *Int J Impot Res*. 2010;22:204-209.
 699. Zhang H, Tong W-T, Zhang C-R, et al. Gross saponin of *Tribulus terrestris* improves erectile dysfunction in type 2 diabetic rats by repairing the endothelial function of the penile corpus cavernosum. *Diabetes Metab Syndr Obes*. 2019;12:1705-1716.
 700. Chen S, Zhu J, Wang M, et al. Comparison of the therapeutic effects of adipose-derived and bone marrow mesenchymal stem cells on erectile dysfunction in diabetic rats. *Int J Mol Med*. 2019;44:1006-1014.
 701. Defeudis G, Mazzilli R, Tenuta M, et al. Erectile dysfunction and diabetes: a melting pot of circumstances and treatments. *Diabetes Metab Res Rev*. 2022;38:e3494.
 702. Gandaglia G, Briganti A, Jackson G, et al. A systematic review of the association between erectile dysfunction and cardiovascular disease. *Eur Urol*. 2014;65:968-978.
 703. Chen F, Zhang H, Wang Z, et al. Adipose-derived stem cell-derived exosomes ameliorate erectile dysfunction in a rat model of type 2 diabetes. *J Sex Med*. 2017;14:1084-1094.
 704. Song J, Sun T, Tang Z, et al. Exosomes derived from smooth muscle cells ameliorate diabetes-induced erectile dysfunction by inhibiting fibrosis and modulating the NO/cGMP pathway. *J Cell Mol Med*. 2020;24:13289-13302.
 705. Di Bartolo BA, Cartland SP, Genner S, et al. HDL improves cholesterol and glucose homeostasis and reduces atherosclerosis in diabetes-associated atherosclerosis. *J Diabetes Res*. 2021;2021:6668506.
 706. Laddha AP, Kulkarni YA. Daidzein mitigates myocardial injury in streptozotocin-induced diabetes in rats. *Life Sci*. 2021;284:119664.
 707. Lin Y, Zhang F, Lian X-f, Peng W-q, Yin C-y. Mesenchymal stem cell-derived exosomes improve diabetes mellitus-induced myocardial injury and fibrosis via inhibition of TGF- β 1/Smad2 signaling pathway. *Cell Mol Biol*. 2019;65:123-126.
 708. Shamardan R, Mahmoud R, Amer E, Ragiae A, Sabry D, Safwat A. Adipose mesenchymal stem cells-derived exosomes attenuate retina degeneration of streptozotocin-induced diabetes in rabbits. *J Circ Biomark*. 2018;7:1849454418807827.
 709. Xiong J, Hu H, Guo R, Wang H, Jiang H. Mesenchymal stem cell exosomes as a new strategy for the treatment of diabetes complications. *Front Endocrinol*. 2021;12:646233.
 710. Ding J, Wang X, Chen B, Zhang J, Xu J. Exosomes derived from human bone marrow mesenchymal stem cells stimulated by deferroxamine accelerate cutaneous wound healing by promoting angiogenesis. *Biomed Res Int*. 2019;2019:9742765.
 711. Shafei S, Khanmohammadi M, Heidari R, et al. Exosome loaded alginate hydrogel promotes tissue regeneration in full-thickness skin wounds: an in vivo study. *J Biomed Mater Res A*. 2020;108:545-556.

712. Lv Q, Deng J, Chen Y, Wang Y, Liu B, Liu J. Engineered human adipose stem-cell-derived exosomes loaded with miR-21-5p to promote diabetic cutaneous wound healing. *Mol Pharm.* 2020;17:1723-1733.
713. Shabbir A, Cox A, Rodriguez-Menocal L, Salgado M, Badiavas EV. Mesenchymal stem cell exosomes induce proliferation and migration of normal and chronic wound fibroblasts, and enhance angiogenesis in vitro. *Stem Cells Dev.* 2015;24:1635-1647.
714. Li B, Luan S, Chen J, et al. The MSC-derived exosomal lncRNA H19 promotes wound healing in diabetic foot ulcers by upregulating PTEN via MicroRNA-152-3p. *Mol Ther Nucleic Acids.* 2020;19:814-826.
715. Shiekh PA, Singh A, Kumar A. Exosome laden oxygen releasing antioxidant and antibacterial cryogel wound dressing OxOBand alleviate diabetic and infectious wound healing. *Biomaterials.* 2020;249:120020.
716. Tao S-C, Guo S-C, Li M, Ke Q-F, Guo Y-P, Zhang C-Q. Chitosan wound dressings incorporating exosomes derived from microRNA-126-overexpressing synovium mesenchymal stem cells provide sustained release of exosomes and heal full-thickness skin defects in a diabetic rat model. *Stem Cells Transl Med.* 2017;6:736-747.
717. Chen C-Y, Rao S-S, Ren L, et al. Exosomal DMBT1 from human urine-derived stem cells facilitates diabetic wound repair by promoting angiogenesis. *Theranostics.* 2018;8:1607-1623.
718. Li X, Xie X, Lian W, et al. Exosomes from adipose-derived stem cells overexpressing Nrf2 accelerate cutaneous wound healing by promoting vascularization in a diabetic foot ulcer rat model. *Exp Mol Med.* 2018;50:1-14.
719. Kobayashi H, Ebisawa K, Kambe M, et al. Effects of exosomes derived from the induced pluripotent stem cells on skin wound healing. *Nagoya J Med Sci.* 2018;80:141.
720. Dalirfardouei R, Jamialahmadi K, Jafarian AH, Mahdipour E. Promising effects of exosomes isolated from menstrual blood-derived mesenchymal stem cell on wound-healing process in diabetic mouse model. *J Tissue Eng Regen Med.* 2019;13:555-568.
721. Wang C, Wang M, Xu T, et al. Engineering bioactive self-healing antibacterial exosomes hydrogel for promoting chronic diabetic wound healing and complete skin regeneration. *Theranostics.* 2019;9:65-76.
722. Wang J, Yi Y, Zhu Y, et al. Effects of adipose-derived stem cell released exosomes on wound healing in diabetic mice. *Zhongguo Xiu Fu Chong Jian Wai Ke Za Zhi.* 2020;34:124-131.
723. Shi R, Jin Y, Hu W, et al. Exosomes derived from mmu_circ_0000250-modified adipose-derived mesenchymal stem cells promote wound healing in diabetic mice by inducing miR-128-3p/SIRT1-mediated autophagy. *Am J Physiol Cell Physiol.* 2020;318:C848-C856.
724. Liu W, Yu M, Xie D, et al. Melatonin-stimulated MSC-derived exosomes improve diabetic wound healing through regulating macrophage M1 and M2 polarization by targeting the PTEN/AKT pathway. *Stem Cell Res Ther.* 2020;11:259.
725. Yu M, Liu W, Li J, et al. Exosomes derived from atorvastatin-pretreated MSC accelerate diabetic wound repair by enhancing angiogenesis via AKT/eNOS pathway. *Stem Cell Res Ther.* 2020;11:350.
726. Yang J, Chen Z, Pan D, Li H, Shen J. Umbilical cord-derived mesenchymal stem cell-derived exosomes combined pluronic F127 hydrogel promote chronic diabetic wound healing and complete skin regeneration. *Int J Nanomedicine.* 2020;15:5911-5926.
727. Jin J, Wang Y, Zhao L, Zou W, Tan M, He Q. Exosomal miRNA-215-5p derived from adipose-derived stem cells attenuates epithelial-mesenchymal transition of podocytes by inhibiting ZEB2. *Biomed Res Int.* 2020;2020:2685305.
728. Jiang Z-z, Liu Y-m, Niu X, et al. Exosomes secreted by human urine-derived stem cells could prevent kidney complications from type I diabetes in rats. *Stem Cell Res Ther.* 2016;7:24.
729. Jin J, Shi Y, Gong J, et al. Exosome secreted from adipose-derived stem cells attenuates diabetic nephropathy by promoting autophagy flux and inhibiting apoptosis in podocyte. *Stem Cell Res Ther.* 2019;10:95.
730. Duan YR, Chen BP, Chen F, et al. Exosomal microRNA-16-5p from human urine-derived stem cells ameliorates diabetic nephropathy through protection of podocyte. *J Cell Mol Med.* 2021;25:10798-10813.
731. Li H, Rong P, Ma X, et al. Mouse umbilical cord mesenchymal stem cell paracrine alleviates renal fibrosis in diabetic nephropathy by reducing myofibroblast transdifferentiation and cell proliferation and upregulating MMPs in mesangial cells. *J Diabetes Res.* 2020;2020.
732. Xiang E, Han B, Zhang Q, et al. Human umbilical cord-derived mesenchymal stem cells prevent the progression of early diabetic nephropathy through inhibiting inflammation and fibrosis. *Stem Cell Res Ther.* 2020;11:336.
733. Wang J, Mi Y, Wu S, et al. Exosomes from adipose-derived stem cells protect against high glucose-induced erectile dysfunction by delivery of corin in a streptozotocin-induced diabetic rat model. *Regen Ther.* 2020;14:227-233.
734. Huo W, Li Y, Zhang Y, Li H. Mesenchymal stem cells-derived exosomal microRNA-21-5p downregulates PDCD4 and ameliorates erectile dysfunction in a rat model of diabetes mellitus. *FASEB J.* 2020;34:13345-13360.
735. Zhao W, Zhang H, Yan J, Ma X. An experimental study on the treatment of diabetes-induced cognitive disorder mice model with exosomes deriving from mesenchymal stem cells (MSCs). *Pak J Pharm Sci.* 2019;32:2599-2603.
736. Kubota K, Nakano M, Kobayashi E, et al. An enriched environment prevents diabetes-induced cognitive impairment in rats by enhancing exosomal miR-146a secretion from endogenous bone marrow-derived mesenchymal stem cells. *PLoS One.* 2018;13:e0204252.
737. Venkat P, Zacharek A, Landschoot-Ward J, et al. Exosomes derived from bone marrow mesenchymal stem cells harvested from type two diabetes rats promotes neurorestorative effects after stroke in type two diabetes rats. *Exp Neurol.* 2020;334:113456.
738. Cui C, Ye X, Chopp M, et al. miR-145 regulates diabetes-bone marrow stromal cell-induced neurorestorative effects in diabetes stroke rats. *Stem Cells Transl Med.* 2016;5:1656-1667.
739. AbuBakr N, Haggag T, Sabry D, Salem ZA. Functional and histological evaluation of bone marrow stem cell-derived exosomes therapy on the submandibular salivary gland of diabetic albino rats through TGFβ/Smad3 signaling pathway. *Heliyon.* 2020;6:e03789.
740. Kordelas L, Rebmann V, Ludwig A, et al. MSC-derived exosomes: a novel tool to treat therapy-refractory graft-versus-host disease. *Leukemia.* 2014;28:970-973.
741. De Veirman K, Wang J, Xu S, et al. Induction of miR-146a by multiple myeloma cells in mesenchymal stromal cells stimulates their pro-tumoral activity. *Cancer Lett.* 2016;377:17-24.
742. Barrera-Ramirez J, Lavoie JR, Maganti HB, et al. Micro-RNA profiling of exosomes from marrow-derived mesenchymal stromal cells in patients with acute myeloid leukemia: implications in leukemogenesis. *Stem Cell Rev Rep.* 2017;13:817-825.
743. Zhang X, Yang Y, Yang Y, Chen H, Tu H, Li J. Exosomes from bone marrow microenvironment-derived mesenchymal stem cells affect CML cells growth and promote drug resistance to tyrosine kinase inhibitors. *Stem Cells Int.* 2020;2020:8890201.
744. Crompton E, Van Damme M, Pieters K, et al. Extracellular vesicles of bone marrow stromal cells rescue chronic lymphocytic leukemia B cells from apoptosis, enhance their migration and induce gene expression modifications. *Haematologica.* 2017;102:1594-1604.
745. Tosetti F, Venè R, Camodeca C, et al. Specific ADAM10 inhibitors localize in exosome-like vesicles released by Hodgkin lymphoma

- and stromal cells and prevent sheddase activity carried to bystander cells. *Onco Targets Ther.* 2018;7:e1421889.
746. Muntion S, Ramos TL, Diez-Campelo M, et al. Microvesicles from mesenchymal stromal cells are involved in HPC-microenvironment crosstalk in myelodysplastic patients. *PLoS One.* 2016;11:e0146722.
 747. Shen M, Chen T. Mesenchymal stem cell-derived exosomes and their potential agents in hematological diseases. *Oxidative Med Cell Longev.* 2021;2021:4539453.
 748. Zhang B, Yeo RWY, Lai RC, Sim EWK, Chin KC, Lim SK. Mesenchymal stromal cell exosome-enhanced regulatory T-cell production through an antigen-presenting cell-mediated pathway. *Cytotherapy.* 2018;20:687-696.
 749. Wang L, Gu Z, Zhao X, et al. Extracellular vesicles released from human umbilical cord-derived mesenchymal stromal cells prevent life-threatening acute graft-versus-host disease in a mouse model of allogeneic hematopoietic stem cell transplantation. *Stem Cells Dev.* 2016;25:1874-1883.
 750. Zhang Q, Fu L, Liang Y, et al. Exosomes originating from MSCs stimulated with TGF- β and IFN- γ promote Treg differentiation. *J Cell Physiol.* 2018;233:6832-6840.
 751. Lai P, Chen X, Guo L, et al. A potent immunomodulatory role of exosomes derived from mesenchymal stromal cells in preventing cGVHD. *J Hematol Oncol.* 2018;11:135.
 752. Gladkova N, Umez T, Imanishi S, Kawana C, Ohyashiki JH, Ohyashiki K. Effect of the extracellular component of bone marrow mesenchymal stromal cells from healthy donors on hematologic neoplasms and their angiogenesis. *Hum Cell.* 2020;33:599-609.
 753. Roccaro AM, Sacco A, Maiso P, et al. BM mesenchymal stromal cell-derived exosomes facilitate multiple myeloma progression. *J Clin Invest.* 2013;123:1542-1555.
 754. Wang J, Hendrix A, Hernot S, et al. Bone marrow stromal cell-derived exosomes as communicators in drug resistance in multiple myeloma cells. *Blood.* 2014;124:555-566.
 755. Dabbah M, Attar-Schneider O, Tartakover Matalon S, et al. Microvesicles derived from normal and multiple myeloma bone marrow mesenchymal stem cells differentially modulate myeloma cells' phenotype and translation initiation. *Carcinogenesis.* 2017;38:708-716.
 756. Umez T, Imanishi S, Azuma K, et al. Replenishing exosomes from older bone marrow stromal cells with miR-340 inhibits myeloma-related angiogenesis. *Blood Adv.* 2017;1:812-823.
 757. Deng M, Yuan H, Liu S, Hu Z, Xiao H. Exosome-transmitted LINC00461 promotes multiple myeloma cell proliferation and suppresses apoptosis by modulating microRNA/BCL-2 expression. *Cytotherapy.* 2019;21:96-106.
 758. Xu H, Han H, Song S, et al. Exosome-transmitted PSMA3 and PSMA3-AS1 promote proteasome inhibitor resistance in multiple myeloma. *Clin Cancer Res.* 2019;25:1923-1935.
 759. Lyu T, Wang Y, Li D, et al. Exosomes from BM-MSCs promote acute myeloid leukemia cell proliferation, invasion and chemoresistance via upregulation of S100A4. *Exp Hematol Oncol.* 2021;10:24.
 760. Viola S, Traer E, Huan J, et al. Alterations in acute myeloid leukaemia bone marrow stromal cell exosome content coincide with gains in tyrosine kinase inhibitor resistance. *Br J Haematol.* 2016;172:983-986.
 761. Liu Y, Song B, Wei Y, et al. Exosomes from mesenchymal stromal cells enhance imatinib-induced apoptosis in human leukemia cells via activation of caspase signaling pathway. *Cytotherapy.* 2018;20:181-188.
 762. Youssef El Baradie KB, Hamrick MW. Therapeutic application of extracellular vesicles for musculoskeletal repair & regeneration. *Connect Tissue Res.* 2021;62:99-114.
 763. Li S, Shao L, Xu T, Jiang X, Yang G, Dong L. An indispensable tool: exosomes play a role in therapy for radiation damage. *Biomed Pharmacother.* 2021;137:111401.
 764. Jeremic B, Fidarova E, Sharma V, et al. The International Atomic Energy Agency (IAEA) randomized trial of palliative treatment of incurable locally advanced non small cell lung cancer (NSCLC) using radiotherapy (RT) and chemotherapy (CHT) in limited resource setting. *Radiother Oncol.* 2015;116:21-26.
 765. Hu Y, Zhang R, Chen G. Exosome and secretion: action on? *Regulation of Cancer Immune Checkpoints: Molecular and Cellular Mechanisms and Therapy*; 2020:455-483.
 766. Rasini V, Dominici M, Kluba T, et al. Mesenchymal stromal/stem cells markers in the human bone marrow. *Cytotherapy.* 2013;15:292-306.
 767. Pu X, Ma S, Gao Y, Xu T, Chang P, Dong L. Mesenchymal stem cell-derived exosomes: biological function and their therapeutic potential in radiation damage. *Cells.* 2020;10:42.
 768. Jacobs SA, Roobrouck VD, Verfaillie CM, Van Gool SW. Immunological characteristics of human mesenchymal stem cells and multipotent adult progenitor cells. *Immunol Cell Biol.* 2013;91:32-39.
 769. Zhang Y, Bi J, Huang J, Tang Y, Du S, Li P. Exosome: a review of its classification, isolation techniques, storage, diagnostic and targeted therapy applications. *Int J Nanomedicine.* 2020;15:6917-6934.
 770. Singh A, Singh A, Sen D. Mesenchymal stem cells in cardiac regeneration: a detailed progress report of the last 6 years (2010-2015). *Stem Cell Res Ther.* 2016;7:82.
 771. Xue J, Li X, Lu Y, et al. Gene-modified mesenchymal stem cells protect against radiation-induced lung injury. *Mol Ther.* 2013;21:456-465.
 772. Li Y, Wang X, Pang Y, Wang S, Luo M, Huang B. The potential therapeutic role of mesenchymal stem cells-derived exosomes in osteoradionecrosis. *J Oncol.* 2021;2021:4758364.
 773. Xu N, He D, Shao Y, et al. Lung-derived exosomes in phosgene-induced acute lung injury regulate the functions of mesenchymal stem cells partially via miR-28-5p. *Biomed Pharmacother.* 2020;121:109603.
 774. Zhang B, Yin Y, Lai RC, Lim SK. Immunotherapeutic potential of extracellular vesicles. *Front Immunol.* 2014;5:518.
 775. Zhu Y-g, Feng X-m, Abbott J, et al. Human mesenchymal stem cell microvesicles for treatment of *Escherichia coli* endotoxin-induced acute lung injury in mice. *Stem Cells.* 2014;32:116-125.
 776. Mahoney JE, Mori M, Szymaniak AD, Varelas X, Cardoso WV. The hippo pathway effector yap controls patterning and differentiation of airway epithelial progenitors. *Dev Cell.* 2014;30:137-150.
 777. Liu X, Li Q, Niu X, et al. Exosomes secreted from human-induced pluripotent stem cell-derived mesenchymal stem cells prevent osteonecrosis of the femoral head by promoting angiogenesis. *Int J Biol Sci.* 2017;13:232-244.
 778. Blazquez R, Sanchez-Margallo FM, de la Rosa O, et al. Immunomodulatory potential of human adipose mesenchymal stem cells derived exosomes on in vitro stimulated T cells. *Front Immunol.* 2014;5:556.
 779. Hu L, Wang J, Zhou X, et al. Exosomes derived from human adipose mesenchymal stem cells accelerates cutaneous wound healing via optimizing the characteristics of fibroblasts. *Sci Rep.* 2016;6:32993.
 780. Ross RJ, Shayan R, Mutimer KL, Ashton MW. Autologous fat grafting: current state of the art and critical review. *Ann Plast Surg.* 2014;73:352-357.
 781. Shukla L, Morrison WA, Shayan R. Adipose-derived stem cells in radiotherapy injury: a new frontier. *Front Surg.* 2015;2:1.
 782. Huang S-P, Huang C-H, Shyu J-F, et al. Promotion of wound healing using adipose-derived stem cells in radiation ulcer of a rat model. *J Biomed Sci.* 2013;20:51.
 783. Forcheron F, Agay D, Scherthan H, et al. Autologous adipocyte derived stem cells favour healing in a minipig model of cutaneous radiation syndrome. *PLoS One.* 2012;7:e31694.

784. Fang S, Xu C, Zhang Y, et al. Umbilical cord-derived mesenchymal stem cell-derived exosomal microRNAs suppress myofibroblast differentiation by inhibiting the transforming growth factor- β /SMAD2 pathway during wound healing. *Stem Cells Transl Med*. 2016;5:1425-1439.
785. Qian X, Xu C, Fang S, et al. Exosomal microRNAs derived from umbilical mesenchymal stem cells inhibit hepatitis C virus infection. *Stem Cells Transl Med*. 2016;5:1190-1203.
786. Tang MK, Yue PY, Ip PP, et al. Soluble E-cadherin promotes tumor angiogenesis and localizes to exosome surface. *Nat Commun*. 2018;9:2270.
787. Ma X, Becker Buscaglia LE, Barker JR, Li Y. MicroRNAs in NF- κ B signaling. *J Mol Cell Biol*. 2011;3:159-166.
788. Sun L, Zhu M, Feng W, et al. Exosomal miRNA Let-7 from menstrual blood-derived endometrial stem cells alleviates pulmonary fibrosis through regulating mitochondrial DNA damage. *Oxidative Med Cell Longev*. 2019;2019:4506303.
789. Geiger A, Walker A, Nissen E. Human fibrocyte-derived exosomes accelerate wound healing in genetically diabetic mice. *Biochem Biophys Res Commun*. 2015;467:303-309.
790. Saha S, Bhanja P, Kabarriti R, Liu L, Alfieri AA, Guha C. Bone marrow stromal cell transplantation mitigates radiation-induced gastrointestinal syndrome in mice. *PLoS One*. 2011;6:e24072.
791. Xu W, Chen J, Liu X, Li H, Qi X, Guo X. Autologous bone marrow stromal cell transplantation as a treatment for acute radiation enteritis induced by a moderate dose of radiation in dogs. *Transl Res*. 2016;171:38-51.
792. Linard C, Busson E, Holler V, et al. Repeated autologous bone marrow-derived mesenchymal stem cell injections improve radiation-induced proctitis in pigs. *Stem Cells Transl Med*. 2013;2:916-927.
793. Liu S, Liu D, Chen C, et al. MSC transplantation improves osteopenia via epigenetic regulation of notch signaling in lupus. *Cell Metab*. 2015;22:606-618.
794. Zuo R, Liu M, Wang Y, et al. BM-MSC-derived exosomes alleviate radiation-induced bone loss by restoring the function of recipient BM-MSCs and activating Wnt/ β -catenin signaling. *Stem Cell Res Ther*. 2019;10:30.
795. Biancone L, Bruno S, Deregis MC, Tetta C, Camussi G. Therapeutic potential of mesenchymal stem cell-derived microvesicles. *Nephrol Dial Transplant*. 2012;27:3037-3042.
796. Xie M-Y, Hou L-J, Sun J-J, et al. Porcine milk exosome MiRNAs attenuate LPS-induced apoptosis through inhibiting TLR4/NF- κ B and p53 pathways in intestinal epithelial cells. *J Agric Food Chem*. 2019;67:9477-9491.
797. Gao R, Zhang R, Qian T, et al. A comparison of exosomes derived from different periods breast milk on protecting against intestinal organoid injury. *Pediatr Surg Int*. 2019;35:1363-1368.
798. Rakici SY, Guzel AI, Tumkaya L, Sevim Nalkiran H, Mercantepe T. Pelvic radiation-induced testicular damage: an experimental study at 1 gray. *Syst Biol Reprod Med*. 2020;66:89-98.
799. Lester SN, Li K. Toll-like receptors in antiviral innate immunity. *J Mol Biol*. 2014;426:1246-1264.
800. Zhang J, Han X, Zhao Y, Xue X, Fan S. Mouse serum protects against total body irradiation-induced hematopoietic system injury by improving the systemic environment after radiation. *Free Radic Biol Med*. 2019;131:382-392.
801. Zhang J, Wang X, Vikash V, et al. ROS and ROS-mediated cellular signaling. *Oxidative Med Cell Longev*. 2016;2016:4350965.
802. Ashraf A, Pervaiz S. Hippo circuitry and the redox modulation of hippo components in cancer cell fate decisions. *Int J Biochem Cell Biol*. 2015;69:20-28.
803. Ma Y, Liu X, Long Y, Chen Y. Emerging therapeutic potential of mesenchymal stem cell-derived extracellular vesicles in chronic respiratory diseases: an overview of recent progress. *Front Bioeng Biotechnol*. 2022;10:845042.
804. Zargar MJ, Kaviani S, Vasei M, Soufi Zomorrod M, Heidari Keshel S, Soleimani M. Therapeutic role of mesenchymal stem cell-derived exosomes in respiratory disease. *Stem Cell Res Ther*. 2022;13:194.
805. Purghe B, Manfredi M, Ragnoli B, Baldanzi G, Malerba M. Exosomes in chronic respiratory diseases. *Biomed Pharmacother*. 2021;144:112270.
806. Putcha N, Puhon MA, Hansel NN, Drummond MB, Boyd CM. Impact of co-morbidities on self-rated health in self-reported COPD: an analysis of NHANES 2001-2008. *COPD*. 2013;10:324-332.
807. Chen Y, Wang H, Luo G, Dai X. SIRT4 inhibits cigarette smoke extracts-induced mononuclear cell adhesion to human pulmonary microvascular endothelial cells via regulating NF- κ B activity. *Toxicol Lett*. 2014;226:320-327.
808. Sundar IK, Li D, Rahman I. Small RNA-sequence analysis of plasma-derived extracellular vesicle miRNAs in smokers and patients with chronic obstructive pulmonary disease as circulating biomarkers. *J Extracell Vesicles*. 2019;8:1684816.
809. Salimian J, Mirzaei H, Moridikia A, Harchegani AB, Sahebkar A, Salehi H. Chronic obstructive pulmonary disease: MicroRNAs and exosomes as new diagnostic and therapeutic biomarkers. *J Res Med Sci*. 2018;23:27.
810. Chen Y, Luo H, Kang N, et al. Beraprost sodium attenuates cigarette smoke extract-induced apoptosis in vascular endothelial cells. *Mol Biol Rep*. 2012;39:10447-10457.
811. Collaborators GCRD. Global, regional, and national deaths, prevalence, disability-adjusted life years, and years lived with disability for chronic obstructive pulmonary disease and asthma, 1990-2015: a systematic analysis for the global burden of disease study 2015. *Lancet Respir Med*. 2017;5:691-706.
812. Hari Krishnan S, Jeemon P, Mini G, Thankappan K, Sylaja P, GBD 2017 Causes of Death Collaborators. Global, regional, and national age-sex-specific mortality for 282 causes of death in 195 countries and territories, 1980-2017: a systematic analysis for the global burden of disease study 2017. *Lancet*. 2018;392:1736-1788.
813. Pouwels SD, Heijink IH, Ten Hacken NH, et al. DAMPs activating innate and adaptive immune responses in COPD. *Mucosal Immunol*. 2014;7:215-226.
814. Kumar M, Seeger W, Voswinkel R. Senescence-associated secretory phenotype and its possible role in chronic obstructive pulmonary disease. *Am J Respir Cell Mol Biol*. 2014;51:323-333.
815. Yao H, Rahman I. Current concepts on oxidative/carbonyl stress, inflammation and epigenetics in pathogenesis of chronic obstructive pulmonary disease. *Toxicol Appl Pharmacol*. 2011;254:72-85.
816. Mei D, Tan WD, Tay Y, Mukhopadhyay A, Wong WF. Therapeutic RNA strategies for chronic obstructive pulmonary disease. *Trends Pharmacol Sci*. 2020;41:475-486.
817. Sun Y, An N, Li J, et al. miRNA-206 regulates human pulmonary microvascular endothelial cell apoptosis via targeting in chronic obstructive pulmonary disease. *J Cell Biochem*. 2019;120:6223-6236.
818. Shen W, Liu J, Fan M, et al. MiR-3202 protects smokers from chronic obstructive pulmonary disease through inhibiting FAIM2: an in vivo and in vitro study. *Exp Cell Res*. 2018;362:370-377.
819. Gomez N, James V, Onion D, Fairclough LC. Extracellular vesicles and chronic obstructive pulmonary disease (COPD): a systematic review. *Respir Res*. 2022;23:82.
820. Miethe S, Guarino M, Alhamdan F, et al. Effects of obesity on asthma: immunometabolic links. *Pol Arch Intern Med*. 2018;128:469-477.
821. Ray A, Oriss TB, Wenzel SE. Emerging molecular phenotypes of asthma. *Am J Physiol Lung Cell Mol Physiol*. 2015;308:L130-L140.
822. Potaczek DP, Miethe S, Schindler V, Alhamdan F, Garn H. Role of airway epithelial cells in the development of different asthma phenotypes. *Cell Signal*. 2020;69:109523.
823. Tost J. A translational perspective on epigenetics in allergic diseases. *J Allergy Clin Immunol*. 2018;142:715-726.

824. Quirt J, Gagnon R, Ellis A, Kim H. CSACI position statement: prescribing sublingual immunotherapy tablets for aeroallergens. *Allergy Asthma Clin Immunol.* 2018;14:1.
825. Bush A. Pathophysiological mechanisms of asthma. *Front Pediatr.* 2019;7:68.
826. Wenzel SE. Asthma phenotypes: the evolution from clinical to molecular approaches. *Nat Med.* 2012;18:716-725.
827. Alashkar Alhamwe B, Miethe S, Pogge von Strandmann E, Potaczek DP, Garn H. Epigenetic regulation of airway epithelium immune functions in asthma. *Front Immunol.* 2020;11:1747.
828. Alashkar Alhamwe B, Potaczek DP, Miethe S, et al. Extracellular vesicles and asthma—more than just a co-existence. *Int J Mol Sci.* 2021;22:4984.
829. Zucca E, Corsini E, Galbiati V, Lange-Consiglio A, Ferrucci F. Evaluation of amniotic mesenchymal cell derivatives on cytokine production in equine alveolar macrophages: an in vitro approach to lung inflammation. *Stem Cell Res Ther.* 2016;7:137.
830. Du Y-m, Zhuansun Y-x, Chen R, Lin L, Lin Y, Li J-g. Mesenchymal stem cell exosomes promote immunosuppression of regulatory T cells in asthma. *Exp Cell Res.* 2018;363:114-120.
831. Cruz FF, Borg ZD, Goodwin M, et al. Systemic administration of human bone marrow-derived mesenchymal stromal cell extracellular vesicles ameliorates aspergillus hyphal extract-induced allergic airway inflammation in immunocompetent mice. *Stem Cells Transl Med.* 2015;4:1302-1316.
832. Shahir M, Mahmoud Hashemi S, Asadirad A, et al. Effect of mesenchymal stem cell-derived exosomes on the induction of mouse tolerogenic dendritic cells. *J Cell Physiol.* 2020;235:7043-7055.
833. de Castro LL, Xisto DG, Kitoko JZ, et al. Human adipose tissue mesenchymal stromal cells and their extracellular vesicles act differentially on lung mechanics and inflammation in experimental allergic asthma. *Stem Cell Res Ther.* 2017;8:151.
834. Shang Y, Sun Y, Xu J, et al. Exosomes from mmu_circ_0001359-modified ADSCs attenuate airway remodeling by enhancing FoxO1 signaling-mediated M2-like macrophage activation. *Mol Ther Nucleic Acids.* 2020;19:951-960.
835. Oh S, Jung J-H, Ahn K-J, et al. Stem cell and exosome therapy in pulmonary hypertension. *Korean Circ J.* 2022;52:110-122.
836. Simonneau G, Montani D, Celermajer DS, et al. Haemodynamic definitions and updated clinical classification of pulmonary hypertension. *Eur Respir J.* 2019;53:1801913.
837. Hoepfer MM, Humbert M, Souza R, et al. A global view of pulmonary hypertension. *Lancet Respir Med.* 2016;4:306-322.
838. Leber L, Beaudet A, Muller A. Epidemiology of pulmonary arterial hypertension and chronic thromboembolic pulmonary hypertension: identification of the most accurate estimates from a systematic literature review. *Pulm Circ.* 2021;11:2045894020977300.
839. Humbert M, Guignabert C, Bonnet S, et al. Pathology and pathobiology of pulmonary hypertension: state of the art and research perspectives. *Eur Respir J.* 2019;53:1801887.
840. Guignabert C, Tu L, Girerd B, et al. New molecular targets of pulmonary vascular remodeling in pulmonary arterial hypertension: importance of endothelial communication. *Chest.* 2015;147:529-537.
841. Frid MG, Thurman JM, Hansen KC, Maron BA, Stenmark KR. Inflammation, immunity, and vascular remodeling in pulmonary hypertension; evidence for complement involvement? *Global Cardiol Sci Pract.* 2020;2020:e202001.
842. Yeo Y, Yi ES, Kim J-M, et al. FGF12 (fibroblast growth factor 12) inhibits vascular smooth muscle cell remodeling in pulmonary arterial hypertension. *Hypertension.* 2020;76:1778-1786.
843. Ahn KJ, Jang AY, Park SJ, Chung W-J. 15 years journey of idiopathic pulmonary arterial hypertension with BMPR2 mutation. *Clin Hypertens.* 2019;25:22.
844. Jang AY, Kim B-G, Kwon S, et al. Prevalence and clinical features of bone morphogenetic protein receptor type 2 mutation in Korean idiopathic pulmonary arterial hypertension patients: the PILGRIM explorative cohort. *PLoS One.* 2020;15:e0238698.
845. Lee H, Yeom A, Kim KC, Hong YM. Effect of ambrisentan therapy on the expression of endothelin receptor, endothelial nitric oxide synthase and NADPH oxidase 4 in monocrotaline-induced pulmonary arterial hypertension rat model. *Korean Circ J.* 2019;49:866-876.
846. Jang AY, Kim S, Park SJ, et al. A nationwide multicenter registry and biobank program for deep phenotyping of idiopathic and hereditary pulmonary arterial hypertension in Korea: the PAH platform for deep phenotyping in Korean subjects (PHOENIX) cohort. *Clin Hypertens.* 2019;25:21.
847. Glembofski CC. Expanding the paracrine hypothesis of stem cell-mediated repair in the heart: when the unconventional becomes conventional. *Circ Res.* 2017;120:772-774.
848. Lopatina T, Bruno S, Tetta C, Kalinina N, Porta M, Camussi G. Platelet-derived growth factor regulates the secretion of extracellular vesicles by adipose mesenchymal stem cells and enhances their angiogenic potential. *Cell Commun Signal.* 2014;12:26.
849. Zhang C, Wang P, Mohammed A, et al. Function of adipose-derived mesenchymal stem cells in monocrotaline-induced pulmonary arterial hypertension through miR-191 via regulation of BMPR2. *Biomed Res Int.* 2019;2019:2858750.
850. Klinger JR, Pereira M, Del Tatto M, et al. Mesenchymal stem cell extracellular vesicles reverse sugen/hypoxia pulmonary hypertension in rats. *Am J Respir Cell Mol Biol.* 2020;62:577-587.
851. Aliotta JM, Pereira M, Wen S, et al. Exosomes induce and reverse monocrotaline-induced pulmonary hypertension in mice. *Cardiovasc Res.* 2016;110:319-330.
852. Lee C, Mitsialis SA, Aslam M, et al. Exosomes mediate the cytoprotective action of mesenchymal stromal cells on hypoxia-induced pulmonary hypertension. *Circulation.* 2012;126:2601-2611.
853. Komaki M, Numata Y, Morioka C, et al. Exosomes of human placenta-derived mesenchymal stem cells stimulate angiogenesis. *Stem Cell Res Ther.* 2017;8:219.
854. Salomon C, Ryan J, Sobrevia L, et al. Exosomal signaling during hypoxia mediates microvascular endothelial cell migration and vasculogenesis. *PLoS One.* 2013;8:e68451.
855. Zhang Z, Ge L, Zhang S, et al. The protective effects of MSC-EXO against pulmonary hypertension through regulating Wnt5a/BMP signalling pathway. *J Cell Mol Med.* 2020;24:13938-13948.
856. Willis GR, Fernandez-Gonzalez A, Reis M, et al. Mesenchymal stromal cell-derived small extracellular vesicles restore lung architecture and improve exercise capacity in a model of neonatal hyperoxia-induced lung injury. *J Extracell Vesicles.* 2020;9:1790874.
857. Ibrahim A, Ibrahim A, Parimon T. Diagnostic and therapeutic applications of extracellular vesicles in interstitial lung diseases. *Diagnostics.* 2021;11:87.
858. Lederer DJ, Martinez FJ. Idiopathic pulmonary fibrosis. *N Engl J Med.* 2018;378:1811-1823.
859. Antin-Ozerkis D, Hinchcliff M. Connective tissue disease-associated interstitial lung disease: evaluation and management. *Clin Chest Med.* 2019;40:617-636.
860. Ryerson CJ, Urbania TH, Richeldi L, et al. Prevalence and prognosis of unclassifiable interstitial lung disease. *Eur Respir J.* 2013;42:750-757.
861. Khor YH, Ng Y, Barnes H, Goh NS, McDonald CF, Holland AE. Prognosis of idiopathic pulmonary fibrosis without anti-fibrotic therapy: a systematic review. *Eur Respir Rev.* 2020;29:190158.
862. Ryerson CJ, Corte TJ, Lee JS, et al. A standardized diagnostic ontology for fibrotic interstitial lung disease. An international working group perspective. *Am J Respir Crit Care Med.* 2017;196:1249-1254.
863. Dhooria S, Agarwal R, Sehgal IS, et al. Spectrum of interstitial lung diseases at a tertiary center in a developing country: a study of 803 subjects. *PLoS One.* 2018;13:e0191938.

864. Distler O, Highland KB, Gahlemann M, et al. Nintedanib for systemic sclerosis-associated interstitial lung disease. *N Engl J Med*. 2019;380:2518-2528.
865. Wells AU, Flaherty KR, Brown KK, et al. Nintedanib in patients with progressive fibrosing interstitial lung diseases—subgroup analyses by interstitial lung disease diagnosis in the INBUILD trial: a randomised, double-blind, placebo-controlled, parallel-group trial. *Lancet Respir Med*. 2020;8:453-460.
866. De Sadeleer LJ, Verleden SE, Vos R, Van Raemdonck D, Verleden GM. Advances in lung transplantation for interstitial lung diseases. *Curr Opin Pulm Med*. 2020;26:518-525.
867. Sakamoto S, Kataoka K, Kondoh Y, et al. Pirfenidone plus inhaled N-acetylcysteine for idiopathic pulmonary fibrosis: a randomised trial. *Eur Respir J*. 2021;57:2000348.
868. Flaherty KR, Wells AU, Cottin V, et al. Nintedanib in progressive fibrosing interstitial lung diseases. *N Engl J Med*. 2019;381:1718-1727.
869. Nathan SD, Albera C, Bradford WZ, et al. Effect of pirfenidone on mortality: pooled analyses and meta-analyses of clinical trials in idiopathic pulmonary fibrosis. *Lancet Respir Med*. 2017;5:33-41.
870. Murphy DE, de Jong OG, Brouwer M, et al. Extracellular vesicle-based therapeutics: natural versus engineered targeting and trafficking. *Exp Mol Med*. 2019;51:1-12.
871. Wiklander OP, Brennan MÁ, Lötvald J, Breakefield XO, El Andaloussi S. Advances in therapeutic applications of extracellular vesicles. *Sci Transl Med*. 2019;11:eaav8521.
872. Yang D, Zhang W, Zhang H, et al. Progress, opportunity, and perspective on exosome isolation-efforts for efficient exosome-based theranostics. *Theranostics*. 2020;10:3684-3707.
873. Worthington EN, Hagood JS. Therapeutic use of extracellular vesicles for acute and chronic lung disease. *Int J Mol Sci*. 2020;21:2318.
874. Lener T, Gimona M, Aigner L, et al. Applying extracellular vesicles based therapeutics in clinical trials – an ISEV position paper. *J Extracell Vesicles*. 2015;4:30087.
875. Wan X, Chen S, Fang Y, Zuo W, Cui J, Xie S. Mesenchymal stem cell-derived extracellular vesicles suppress the fibroblast proliferation by downregulating FZD6 expression in fibroblasts via miRNA-29b-3p in idiopathic pulmonary fibrosis. *J Cell Physiol*. 2020;235:8613-8625.
876. Dinh P-UC, Paudel D, Brochu H, et al. Inhalation of lung spheroid cell secretome and exosomes promotes lung repair in pulmonary fibrosis. *Nat Commun*. 2020;11:1064.
877. Lananna BV, Si I. Friends and foes: extracellular vesicles in aging and rejuvenation. *FASEB Bioadv*. 2021;3:787-801.
878. Prattichizzo F, Giuliani A, Sabbatinelli J, et al. Extracellular vesicles circulating in young organisms promote healthy longevity. *J Extracell Vesicles*. 2019;8:1656044.
879. Mahmoudi S, Xu L, Brunet A. Turning back time with emerging rejuvenation strategies. *Nat Cell Biol*. 2019;21:32-43.
880. Takasugi M. Emerging roles of extracellular vesicles in cellular senescence and aging. *Aging Cell*. 2018;17:e12734.
881. Shah R, Patel T, Freedman JE. Circulating extracellular vesicles in human disease. *N Engl J Med*. 2018;379:958-966.
882. Robbins PD. Extracellular vesicles and aging. *Stem Cell Investig*. 2017;4.
883. Forest A, Pautas E, Ray P, et al. Circulating microparticles and pro-coagulant activity in elderly patients. *J Gerontol A Biol Sci Med Sci*. 2010;65:414-420.
884. Villeda SA, Plambeck KE, Middeldorp J, et al. Young blood reverses age-related impairments in cognitive function and synaptic plasticity in mice. *Nat Med*. 2014;20:659-663.
885. Castellano JM, Mosher KI, Abbey RJ, et al. Human umbilical cord plasma proteins revitalize hippocampal function in aged mice. *Nature*. 2017;544:488-492.
886. Horowitz AM, Fan X, Bieri G, et al. Blood factors transfer beneficial effects of exercise on neurogenesis and cognition to the aged brain. *Science*. 2020;369:167-173.
887. Conboy IM, Conboy MJ, Wagers AJ, Girma ER, Weissman IL, Rando TA. Rejuvenation of aged progenitor cells by exposure to a young systemic environment. *Nature*. 2005;433:760-764.
888. Brahmer A, Neuberger E, Esch-Heisser L, et al. Platelets, endothelial cells and leukocytes contribute to the exercise-triggered release of extracellular vesicles into the circulation. *J Extracell Vesicles*. 2019;8:1615820.
889. Sharon JS, Deutsch GK, Tian L, et al. Safety, tolerability, and feasibility of young plasma infusion in the plasma for Alzheimer symptom amelioration study: a randomized clinical trial. *JAMA Neurol*. 2019;76:35-40.
890. Zhang Y, Kim MS, Jia B, et al. Hypothalamic stem cells control ageing speed partly through exosomal miRNAs. *Nature*. 2017;548:52-57.
891. Satoh A, Brace CS, Rensing N, et al. Sirt1 extends life span and delays aging in mice through the regulation of Nk2 homeobox 1 in the DMH and LH. *Cell Metab*. 2013;18:416-430.
892. Xiao Y-Z, Yang M, Xiao Y, et al. Reducing hypothalamic stem cell senescence protects against aging-associated physiological decline. *Cell Metab*. 2020;31:534-548.e5.
893. Yoshida M, Satoh A, Lin JB, et al. Extracellular vesicle-contained eNAMPT delays aging and extends lifespan in mice. *Cell Metab*. 2019;30:329-342.e5.
894. Yoon MJ, Yoshida M, Johnson S, et al. SIRT1-mediated eNAMPT secretion from adipose tissue regulates hypothalamic NAD⁺ and function in mice. *Cell Metab*. 2015;21:706-717.
895. Imai S-i. The NAD world 2.0: the importance of the inter-tissue communication mediated by NAMPT/NAD⁺/SIRT1 in mammalian aging and longevity control. *NPJ Syst Biol Appl*. 2016;2:16018.
896. Xu R, Rai A, Chen M, Suwakulsiri W, Greening DW, Simpson RJ. Extracellular vesicles in cancer—implications for future improvements in cancer care. *Nat Rev Clin Oncol*. 2018;15:617-638.
897. Robbins PD, Dorronsoro A, Booker CN. Regulation of chronic inflammatory and immune processes by extracellular vesicles. *J Clin Invest*. 2016;126:1173-1180.
898. Théry C, Ostrowski M, Segura E. Membrane vesicles as conveyors of immune responses. *Nat Rev Immunol*. 2009;9:581-593.
899. Kitai Y, Kawasaki T, Sueyoshi T, et al. DNA-containing exosomes derived from cancer cells treated with topotecan activate a STING-dependent pathway and reinforce antitumor immunity. *J Immunol*. 2017;198:1649-1659.
900. Al-Mayah AH, Bright SJ, Bowler DA, Slijepcevic P, Goodwin E, Kadhim MA. Exosome-mediated telomere instability in human breast epithelial cancer cells after X irradiation. *Radiat Res*. 2017;187:98-106.
901. Fafián-Labora JA, Rodríguez-Navarro JA, O'Loughlin A. Small extracellular vesicles have GST activity and ameliorate senescence-related tissue damage. *Cell Metab*. 2020;32:71-86.e5.
902. DeLise AM, Fischer L, Tuan R. Cellular interactions and signaling in cartilage development. *Osteoarthritis Cartil*. 2000;8:309-334.
903. Findlay DM, Kuliwaba JS. Bone–cartilage crosstalk: a conversation for understanding osteoarthritis. *Bone Res*. 2016;4:16028.
904. Zioupos P, Currey J, Hamer A. The role of collagen in the declining mechanical properties of aging human cortical bone. *J Biomed Mater Res*. 1999;45:108-116.
905. Sivaraj KK, Adams RH. Blood vessel formation and function in bone. *Development*. 2016;143:2706-2715.
906. Doorn J, Moll G, Le Blanc K, Van Blitterswijk C, de Boer J. Therapeutic applications of mesenchymal stromal cells: paracrine effects and potential improvements. *Tissue Eng Part B Rev*. 2012;18:101-115.
907. Gronthos S, Mankani M, Brahimi J, Robey PG, Shi S. Postnatal human dental pulp stem cells (DPSCs) in vitro and in vivo. *Proc Natl Acad Sci*. 2000;97:13625-13630.
908. Seo B-M, Miura M, Gronthos S, et al. Investigation of multipotent postnatal stem cells from human periodontal ligament. *Lancet*. 2004;364:149-155.

909. Grawish ME. Gingival-derived mesenchymal stem cells: an endless resource for regenerative dentistry. *World J Stem Cells*. 2018;10:116-118.
910. Liu H, Li D, Zhang Y, Li M. Inflammation, mesenchymal stem cells and bone regeneration. *Histochem Cell Biol*. 2018;149:393-404.
911. Naji A, Favier B, Deschaseaux F, Rouas-Freiss N, Eitoku M, Suganuma N. Mesenchymal stem/stromal cell function in modulating cell death. *Stem Cell Res Ther*. 2019;10:56.
912. Wu T, Liu Y, Wang B, Li G. The roles of mesenchymal stem cells in tissue repair and disease modification. *Curr Stem Cell Res Ther*. 2014;9:424-431.
913. Dominici M, Le Blanc K, Mueller I, et al. Minimal criteria for defining multipotent mesenchymal stromal cells. The International Society for Cellular Therapy position statement. *Cytotherapy*. 2006;8:315-317.
914. Jackson L, Jones D, Scotting P, Sottile V. Adult mesenchymal stem cells: differentiation potential and therapeutic applications. *J Postgrad Med*. 2007;53:121-127.
915. Turner L, Knoepfler P. Selling stem cells in the USA: assessing the direct-to-consumer industry. *Cell Stem Cell*. 2016;19:154-157.
916. Chen S, Tang Y, Liu Y, et al. Exosomes derived from miR-375-overexpressing human adipose mesenchymal stem cells promote bone regeneration. *Cell Prolif*. 2019;52:e12669.
917. Takeuchi R, Katagiri W, Endo S, Kobayashi T. Exosomes from conditioned media of bone marrow-derived mesenchymal stem cells promote bone regeneration by enhancing angiogenesis. *PLoS One*. 2019;14:e0225472.
918. Morrell AE, Brown GN, Robinson ST, et al. Mechanically induced Ca^{2+} oscillations in osteocytes release extracellular vesicles and enhance bone formation. *Bone Res*. 2018;6:6.
919. Furuta T, Miyaki S, Ishitobi H, et al. Mesenchymal stem cell-derived exosomes promote fracture healing in a mouse model. *Stem Cells Transl Med*. 2016;5:1620-1630.
920. Zhang Y, Hao Z, Wang P, et al. Exosomes from human umbilical cord mesenchymal stem cells enhance fracture healing through HIF-1 α -mediated promotion of angiogenesis in a rat model of stabilized fracture. *Cell Prolif*. 2019;52:e12570.
921. Liu W, Li L, Rong Y, et al. Hypoxic mesenchymal stem cell-derived exosomes promote bone fracture healing by the transfer of miR-126. *Acta Biomater*. 2020;103:196-212.
922. Qi X, Zhang J, Yuan H, et al. Exosomes secreted by human-induced pluripotent stem cell-derived mesenchymal stem cells repair critical-sized bone defects through enhanced angiogenesis and osteogenesis in osteoporotic rats. *Int J Biol Sci*. 2016;12:836-849.
923. Zhang J, Liu X, Li H, et al. Exosomes/tricalcium phosphate combination scaffolds can enhance bone regeneration by activating the PI3K/Akt signaling pathway. *Stem Cell Res Ther*. 2016;7:136.
924. Pizzicannella J, Diomedea F, Gugliandolo A, et al. 3D printing PLA/gingival stem cells/EVs upregulate miR-2861 and-210 during osteoangiogenesis commitment. *Int J Mol Sci*. 2019;20:3256.
925. Li W, Liu Y, Zhang P, et al. Tissue-engineered bone immobilized with human adipose stem cells-derived exosomes promotes bone regeneration. *ACS Appl Mater Interfaces*. 2018;10:5240-5254.
926. Diomedea F, Gugliandolo A, Cardelli P, et al. Three-dimensional printed PLA scaffold and human gingival stem cell-derived extracellular vesicles: a new tool for bone defect repair. *Stem Cell Res Ther*. 2018;9:104.
927. Pizzicannella J, Gugliandolo A, Orsini T, et al. Engineered extracellular vesicles from human periodontal-ligament stem cells increase VEGF/VEGFR2 expression during bone regeneration. *Front Physiol*. 2019;10:512.
928. Otsuru S, Desbordes L, Guess AJ, et al. Extracellular vesicles released from mesenchymal stromal cells stimulate bone growth in osteogenesis imperfecta. *Cytotherapy*. 2018;20:62-73.
929. Li H, Liu D, Li C, et al. Exosomes secreted from mutant-HIF-1 α -modified bone-marrow-derived mesenchymal stem cells attenuate early steroid-induced avascular necrosis of femoral head in rabbit. *Cell Biol Int*. 2017;41:1379-1390.
930. Zhang S, Chu W, Lai R, Lim S, Hui J, Toh W. Exosomes derived from human embryonic mesenchymal stem cells promote osteochondral regeneration. *Osteoarthritis Cartil*. 2016;24:2135-2140.
931. Zhang S, Chuah SJ, Lai RC, Hui JHP, Lim SK, Toh WS. MSC exosomes mediate cartilage repair by enhancing proliferation, attenuating apoptosis and modulating immune reactivity. *Biomaterials*. 2018;156:16-27.
932. Mohammed E, Khalil E, Sabry D. Effect of adipose-derived stem cells and their exo as adjunctive therapy to nonsurgical periodontal treatment: a histologic and histomorphometric study in rats. *Biomolecules*. 2018;8:167.
933. Chew JRJ, Chuah SJ, Teo KYW, et al. Mesenchymal stem cell exosomes enhance periodontal ligament cell functions and promote periodontal regeneration. *Acta Biomater*. 2019;89:252-264.
934. Wang X, Thomsen P. Mesenchymal stem cell-derived small extracellular vesicles and bone regeneration. *Basic Clin Pharmacol Toxicol*. 2021;128:18-36.
935. Xie H, Wang Z, Zhang L, et al. Extracellular vesicle-functionalized decalcified bone matrix scaffolds with enhanced pro-angiogenic and pro-bone regeneration activities. *Sci Rep*. 2017;7:45622.
936. Wang X, Omar O, Vazirani F, Thomsen P, Ekström K. Mesenchymal stem cell-derived exosomes have altered microRNA profiles and induce osteogenic differentiation depending on the stage of differentiation. *PLoS One*. 2018;13:e0193059.
937. Zhu Y, Jia Y, Wang Y, Xu J, Chai Y. Impaired bone regenerative effect of exosomes derived from bone marrow mesenchymal stem cells in type 1 diabetes. *Stem Cells Transl Med*. 2019;8:593-605.
938. Omar O, Lennerås M, Svensson S, et al. Integrin and chemokine receptor gene expression in implant-adherent cells during early osseointegration. *J Mater Sci Mater Med*. 2010;21:969-980.
939. Palmquist A, Omar OM, Esposito M, Lausmaa J, Thomsen P. Titanium oral implants: surface characteristics, interface biology and clinical outcome. *J R Soc Interface*. 2010;7:S515-S527.
940. Wang X, Shah FA, Vazirani F, et al. Exosomes influence the behavior of human mesenchymal stem cells on titanium surfaces. *Biomaterials*. 2020;230:119571.
941. Lu Z, Chen Y, Dunstan C, Roohani-Esfahani S, Zreiqat H. Priming adipose stem cells with tumor necrosis factor- α preconditioning potentiates their exosome efficacy for bone regeneration. *Tissue Eng Part A*. 2017;23:1212-1220.
942. Fujita T, Azuma Y, Fukuyama R, et al. Runx2 induces osteoblast and chondrocyte differentiation and enhances their migration by coupling with PI3K-Akt signaling. *J Cell Biol*. 2004;166:85-95.
943. Ghosh-Choudhury N, Abboud SL, Nishimura R, Celeste A, Mahimainathan L, Choudhury GG. Requirement of BMP-2-induced phosphatidylinositol 3-kinase and Akt serine/threonine kinase in osteoblast differentiation and Smad-dependent BMP-2 gene transcription. *J Biol Chem*. 2002;277:33361-33368.
944. Cui A, Li H, Wang D, Zhong J, Chen Y, Lu H. Global, regional prevalence, incidence and risk factors of knee osteoarthritis in population-based studies. *EClinicalMedicine*. 2020;29:100587.
945. Mobasheri A, Batt M. An update on the pathophysiology of osteoarthritis. *Ann Phys Rehabil Med*. 2016;59:333-339.
946. Abramoff B, Caldera FE. Osteoarthritis: pathology, diagnosis, and treatment options. *Med Clin*. 2020;104:293-311.
947. Buckwalter J, Mankin H. Articular cartilage: degeneration and osteoarthritis, repair, regeneration, and transplantation. *Instr Course Lect*. 1998;47:487-504.
948. Re'em T, Witte F, Willbold E, Ruvinov E, Cohen S. Simultaneous regeneration of articular cartilage and subchondral bone induced by spatially presented TGF- β and BMP-4 in a bilayer affinity binding system. *Acta Biomater*. 2012;8:3283-3293.

949. Liao LL, Hassan MNF, Tang YL, Ng MH, Law JX. Feasibility of human platelet lysate as an alternative to foetal bovine serum for in vitro expansion of chondrocytes. *Int J Mol Sci*. 2021;22:1269.
950. Evans CH, Kraus VB, Setton LA. Progress in intra-articular therapy. *Nat Rev Rheumatol*. 2014;10:11-22.
951. Pinheiro A, Silva AM, Teixeira JH, et al. Extracellular vesicles: intelligent delivery strategies for therapeutic applications. *J Control Release*. 2018;289:56-69.
952. Patil M, Henderson J, Luong H, Annamalai D, Sreejit G, Krishnamurthy P. The art of intercellular wireless communications: exosomes in heart disease and therapy. *Front Cell Dev Biol*. 2019;7:315.
953. Quesenberry PJ, Aliotta J, Deregis MC, Camussi G. Role of extracellular RNA-carrying vesicles in cell differentiation and reprogramming. *Stem Cell Res Ther*. 2015;6:153.
954. Ng CY, Chai JY, Foo JB, et al. Potential of exosomes as cell-free therapy in articular cartilage regeneration: a review. *Int J Nanomedicine*. 2021;16:6749-6781.
955. Liu C, Li Y, Yang Z, Zhou Z, Lou Z, Zhang Q. Kartogenin enhances the therapeutic effect of bone marrow mesenchymal stem cells derived exosomes in cartilage repair. *Nanomedicine*. 2020;15:273-288.
956. Liu X, Wang L, Ma C, Wang G, Zhang Y, Sun S. Exosomes derived from platelet-rich plasma present a novel potential in alleviating knee osteoarthritis by promoting proliferation and inhibiting apoptosis of chondrocyte via Wnt/ β -catenin signaling pathway. *J Orthop Surg Res*. 2019;14:470.
957. Qi H, Liu D-P, Xiao D-W, Tian D-C, Su Y-W, Jin S-F. Exosomes derived from mesenchymal stem cells inhibit mitochondrial dysfunction-induced apoptosis of chondrocytes via p38, ERK, and Akt pathways. *In Vitro Cell Dev Biol Anim*. 2019;55:203-210.
958. Zhao S, Xiu G, Wang J, et al. Engineering exosomes derived from subcutaneous fat MSCs specially promote cartilage repair as miR-199a-3p delivery vehicles in osteoarthritis. *J Nanobiotechnol*. 2023;21:341.
959. Zheng L, Wang Y, Qiu P, et al. Primary chondrocyte exosomes mediate osteoarthritis progression by regulating mitochondrion and immune reactivity. *Nanomedicine*. 2019;14:3193-3212.
960. Jin Z, Ren J, Qi S. Exosomal miR-9-5p secreted by bone marrow-derived mesenchymal stem cells alleviates osteoarthritis by inhibiting syndecan-1. *Cell Tissue Res*. 2020;381:99-114.
961. Liang Y, Xu X, Li X, et al. Chondrocyte-targeted microRNA delivery by engineered exosomes toward a cell-free osteoarthritis therapy. *ACS Appl Mater Interfaces*. 2020;12:36938-36947.
962. Wong KL, Zhang S, Wang M, et al. Intra-articular injections of mesenchymal stem cell exosomes and hyaluronic acid improve structural and mechanical properties of repaired cartilage in a rabbit model. *Arthroscopy*. 2020;36:2215-2228.e2.
963. Yan L, Wu X. Exosomes produced from 3D cultures of umbilical cord mesenchymal stem cells in a hollow-fiber bioreactor show improved osteochondral regeneration activity. *Cell Biol Toxicol*. 2020;36:165-178.
964. Zavatti M, Beretti F, Casciaro F, Bertucci E, Maraldi T. Comparison of the therapeutic effect of amniotic fluid stem cells and their exosomes on monoiodoacetate-induced animal model of osteoarthritis. *Biofactors*. 2020;46:106-117.
965. Wang Y, He SH, Liang X, Zhang XX, Li SS, Li TF. ATF4-modified serum exosomes derived from osteoarthritic mice inhibit osteoarthritis by inducing autophagy. *IUBMB Life*. 2021;73:146-158.
966. Yan L, Liu G, Wu X. Exosomes derived from umbilical cord mesenchymal stem cells in mechanical environment show improved osteochondral activity via upregulation of lncRNA H19. *J Orthop Translat*. 2021;26:111-120.
967. Lui PPY. Mesenchymal stem cell-derived extracellular vesicles for the promotion of tendon repair—an update of literature. *Stem Cell Rev Rep*. 2021;17:379-389.
968. Lim WL, Liao LL, Ng MH, Chowdhury SR, Law JX. Current progress in tendon and ligament tissue engineering. *Tissue Eng Regen Med*. 2019;16:549-571.
969. Gissi C, Radeghieri A, Antonetti Lamorgese Passeri C, et al. Extracellular vesicles from rat-bone-marrow mesenchymal stromal/stem cells improve tendon repair in rat Achilles tendon injury model in dose-dependent manner: a pilot study. *PLoS One*. 2020;15:e0229914.
970. Madhi MI, Yausep OE, Khamdan K, Trigkilidas D. The use of PRP in treatment of Achilles tendinopathy: a systematic review of literature. Study design: systematic review of literature. *Ann Med Surg*. 2020;55:320-326.
971. Lin M-T, Wei K-C, Wu C-H. Effectiveness of platelet-rich plasma injection in rotator cuff tendinopathy: a systematic review and meta-analysis of randomized controlled trials. *Diagnostics*. 2020;10:189.
972. Cruciani M, Franchini M, Mengoli C, et al. Platelet-rich plasma for sports-related muscle, tendon and ligament injuries: an umbrella review. *Blood Transfus*. 2019;17:465.
973. Kia C, Baldino J, Bell R, Ramji A, Uyeki C, Mazzocca A. Platelet-rich plasma: review of current literature on its use for tendon and ligament pathology. *Curr Rev Musculoskelet Med*. 2018;11:566-572.
974. Shi Z, Wang Q, Jiang D. Extracellular vesicles from bone marrow-derived multipotent mesenchymal stromal cells regulate inflammation and enhance tendon healing. *J Transl Med*. 2019;17:211.
975. Yu H, Cheng J, Shi W, et al. Bone marrow mesenchymal stem cell-derived exosomes promote tendon regeneration by facilitating the proliferation and migration of endogenous tendon stem/progenitor cells. *Acta Biomater*. 2020;106:328-341.
976. Wang Y, He G, Guo Y, et al. Exosomes from tendon stem cells promote injury tendon healing through balancing synthesis and degradation of the tendon extracellular matrix. *J Cell Mol Med*. 2019;23:5475-5485.
977. Shen H, Korpakis I, Havlioglu N, et al. The effect of mesenchymal stromal cell sheets on the inflammatory stage of flexor tendon healing. *Stem Cell Res Ther*. 2016;7:144.
978. Gelberman RH, Linderman SW, Jayaram R, et al. Combined administration of ASCs and BMP-12 promotes an M2 macrophage phenotype and enhances tendon healing. *Clin Orthop Relat Res*. 2017;475:2318-2331.
979. Manning CN, Martel C, Sakiyama-Elbert SE, et al. Adipose-derived mesenchymal stromal cells modulate tendon fibroblast responses to macrophage-induced inflammation in vitro. *Stem Cell Res Ther*. 2015;6:74.
980. Chamberlain CS, Clements AE, Kink JA, et al. Extracellular vesicle-educated macrophages promote early achilles tendon healing. *Stem Cells*. 2019;37:652-662.
981. Kornicka-Garbowska K, Pędziwiatr R, Woźniak P, Kucharczyk K, Marycz K. Microvesicles isolated from 5-azacytidine-and-resve ratrot-treated mesenchymal stem cells for the treatment of suspensory ligament injury in horse—a case report. *Stem Cell Res Ther*. 2019;10:394.
982. Shen H, Yoneda S, Abu-Amer Y, Guilak F, Gelberman RH. Stem cell-derived extracellular vesicles attenuate the early inflammatory response after tendon injury and repair. *J Orthop Res*. 2020;38:117-127.
983. Shi Y, Kang X, Wang Y, et al. Exosomes derived from bone marrow stromal cells (BMSCs) enhance tendon-bone healing by regulating macrophage polarization. *Med Sci Monit*. 2020;26:e923328.
984. Wang C, Hu Q, Song W, Yu W, He Y. Adipose stem cell-derived exosomes decrease fatty infiltration and enhance rotator cuff healing in a rabbit model of chronic tears. *Am J Sports Med*. 2020;48:1456-1464.
985. Marofi F, Alexandrovna KI, Margiana R, et al. MSCs and their exosomes: a rapidly evolving approach in the context of cutaneous wounds therapy. *Stem Cell Res Ther*. 2021;12:597.

986. Weiliang Z, Lili G. Research advances in the application of adipose-derived stem cells derived exosomes in cutaneous wound healing. *Ann Dermatol*. 2021;33:309-317.
987. An Y, Lin S, Tan X, et al. Exosomes from adipose-derived stem cells and application to skin wound healing. *Cell Prolif*. 2021;54:e12993.
988. Manchon E, Hirt N, Bouaziz J-D, Jabrane-Ferrat N, Al-Daccak R. Stem cells-derived extracellular vesicles: potential therapeutics for wound healing in chronic inflammatory skin diseases. *Int J Mol Sci*. 2021;22:3130.
989. Wan R, Hussain A, Behfar A, Moran SL, Zhao C. The therapeutic potential of exosomes in soft tissue repair and regeneration. *Int J Mol Sci*. 2022;23:3869.
990. Zhao M, Shi J, Cai W, et al. Advances on graphene-based nanomaterials and mesenchymal stem cell-derived exosomes applied in cutaneous wound healing. *Int J Nanomedicine*. 2021;16:2647-2665.
991. Sampath R, Rudaityte A, Stašyte G, Aparicio Calvente M, Jekabsone A. Extracellular vesicles in skin wound healing. *Pharmaceutics*. 2021;13:1543.
992. Bray ER, Oropallo AR, Grande DA, Kirsner RS, Badiavas EV. Extracellular vesicles as therapeutic tools for the treatment of chronic wounds. *Pharmaceutics*. 2021;13:1543.
993. Vu NB, Nguyen HT, Palumbo R, Pellicano R, Fagoonee S, Pham PV. Stem cell-derived exosomes for wound healing: current status and promising directions. *Minerva Med*. 2020;112:384-400.
994. Huang J, Zhang J, Xiong J, et al. Stem cell-derived nanovesicles: a novel cell-free therapy for wound healing. *Stem Cells Int*. 2021;2021:1285087.
995. Zeng Q-L, Liu D-W. Mesenchymal stem cell-derived exosomes: an emerging therapeutic strategy for normal and chronic wound healing. *World J Clin Cases*. 2021;9:6218-6233.
996. Liew FF, Chew BC, Ooi DJ. Wound healing properties of exosomes—A review and modelling of combinatorial analysis strategies. *Curr Mol Med*. 2022;22:165-191.
997. Silachev DN, Goryunov KV, Shpilyuk MA, et al. Effect of MSCs and MSC-derived extracellular vesicles on human blood coagulation. *Cells*. 2019;8:258.
998. He X, Dong Z, Cao Y, et al. MSC-derived exosome promotes M2 polarization and enhances cutaneous wound healing. *Stem Cells Int*. 2019;2019:7132708.
999. Wang L, Hu L, Zhou X, et al. Exosomes secreted by human adipose mesenchymal stem cells promote scarless cutaneous repair by regulating extracellular matrix remodelling. *Sci Rep*. 2017;7:13321.
1000. Jiang T, Wang Z, Sun J. Human bone marrow mesenchymal stem cell-derived exosomes stimulate cutaneous wound healing mediates through TGF- β /Smad signaling pathway. *Stem Cell Res Ther*. 2020;11:198.
1001. Wang X, Jiao Y, Pan Y, et al. Fetal dermal mesenchymal stem cell-derived exosomes accelerate cutaneous wound healing by activating notch signaling. *Stem Cells Int*. 2019;2019:2402916.
1002. Zhang B, Wang M, Gong A, et al. HucMSC-exosome mediated-Wnt4 signaling is required for cutaneous wound healing. *Stem Cells*. 2015;33:2158-2168.
1003. Kim S, Lee SK, Kim H, Kim TM. Exosomes secreted from induced pluripotent stem cell-derived mesenchymal stem cells accelerate skin cell proliferation. *Int J Mol Sci*. 2018;19:3119.
1004. Zhang Y, Han F, Gu L, et al. Adipose mesenchymal stem cell exosomes promote wound healing through accelerated keratinocyte migration and proliferation by activating the AKT/HIF-1 α axis. *J Mol Histol*. 2020;51:375-383.
1005. Zhao G, Liu F, Liu Z, et al. MSC-derived exosomes attenuate cell death through suppressing AIF nucleus translocation and enhance cutaneous wound healing. *Stem Cell Res Ther*. 2020;11:174.
1006. Liang X, Zhang L, Wang S, Han Q, Zhao RC. Exosomes secreted by mesenchymal stem cells promote endothelial cell angiogenesis by transferring miR-125a. *J Cell Sci*. 2016;129:2182-2189.
1007. Fan B, Li C, Szalad A, et al. Mesenchymal stromal cell-derived exosomes ameliorate peripheral neuropathy in a mouse model of diabetes. *Diabetologia*. 2020;63:431-443.
1008. Alzahrani FA. Melatonin improves therapeutic potential of mesenchymal stem cells-derived exosomes against renal ischemia-reperfusion injury in rats. *Am J Transl Res*. 2019;11:2887-2907.
1009. Ha DH, Kim H-k, Lee J, et al. Mesenchymal stem/stromal cell-derived exosomes for immunomodulatory therapeutics and skin regeneration. *Cells*. 2020;9:1157.
1010. Liu J, Yan Z, Yang F, et al. Exosomes derived from human umbilical cord mesenchymal stem cells accelerate cutaneous wound healing by enhancing angiogenesis through delivering angiopoietin-2. *Stem Cell Rev Rep*. 2021;17:305-317.
1011. Sung DK, Chang YS, Sung SI, Ahn SY, Park WS. Thrombin preconditioning of extracellular vesicles derived from mesenchymal stem cells accelerates cutaneous wound healing by boosting their biogenesis and enriching cargo content. *J Clin Med*. 2019;8:533.
1012. Xie K, Liu L, Chen J, Liu F. Exosomes derived from human umbilical cord blood mesenchymal stem cells improve hepatic ischemia reperfusion injury via delivering miR-1246. *Cell Cycle*. 2019;18:3491-3501.
1013. Shi Q, Qian Z, Liu D, et al. GMSC-derived exosomes combined with a chitosan/silk hydrogel sponge accelerates wound healing in a diabetic rat skin defect model. *Front Physiol*. 2017;8:904.
1014. Zhang W, Bai X, Zhao B, et al. Cell-free therapy based on adipose tissue stem cell-derived exosomes promotes wound healing via the PI3K/Akt signaling pathway. *Exp Cell Res*. 2018;370:333-342.
1015. Ren S, Chen J, Duscher D, et al. Microvesicles from human adipose stem cells promote wound healing by optimizing cellular functions via AKT and ERK signaling pathways. *Stem Cell Res Ther*. 2019;10:47.
1016. Cheng S, Xi Z, Chen G, Liu K, Ma R, Zhou C. Extracellular vesicle-carried microRNA-27b derived from mesenchymal stem cells accelerates cutaneous wound healing via E3 ubiquitin ligase ITCH. *J Cell Mol Med*. 2020;24:11254-11271.
1017. Jiang L, Zhang Y, Liu T, et al. Exosomes derived from TSG-6 modified mesenchymal stromal cells attenuate scar formation during wound healing. *Biochimie*. 2020;177:40-49.
1018. Liu J, Qiu X, Lv Y, et al. Apoptotic bodies derived from mesenchymal stem cells promote cutaneous wound healing via regulating the functions of macrophages. *Stem Cell Res Ther*. 2020;11:507.
1019. Qiu X, Liu J, Zheng C, et al. Exosomes released from educated mesenchymal stem cells accelerate cutaneous wound healing via promoting angiogenesis. *Cell Prolif*. 2020;53:e12830.
1020. Li Y, Zhang J, Shi J, et al. Exosomes derived from human adipose mesenchymal stem cells attenuate hypertrophic scar fibrosis by miR-192-5p/IL-17RA/Smad axis. *Stem Cell Res Ther*. 2021;12:221.
1021. Pomatto M, Gai C, Negro F, et al. Differential therapeutic effect of extracellular vesicles derived by bone marrow and adipose mesenchymal stem cells on wound healing of diabetic ulcers and correlation to their cargoes. *Int J Mol Sci*. 2021;22:3851.
1022. Zhang J, Guan J, Niu X, et al. Exosomes released from human induced pluripotent stem cells-derived MSCs facilitate cutaneous wound healing by promoting collagen synthesis and angiogenesis. *J Transl Med*. 2015;13:49.
1023. Narauskaitė D, Vydmantaitė G, Rusteikaitė J, et al. Extracellular vesicles in skin wound healing. *Pharmaceutics*. 2021;14(8):811.
1024. Daltro SRT, Meira CS, Santos IP, Ribeiro dos Santos R, Soares MBP. Mesenchymal stem cells and atopic dermatitis: a review. *Front Cell Dev Biol*. 2020;8:326.
1025. Drucker AM, Ellis AG, Bohdanowicz M, et al. Systemic immunomodulatory treatments for patients with atopic dermatitis: a systematic review and network meta-analysis. *JAMA Dermatol*. 2020;156:659-667.
1026. Quiñones-Vico MI, Sanabria-De La Torre R, Sánchez-Díaz M, et al. The role of exosomes derived from mesenchymal stromal cells in dermatology. *Front Cell Dev Biol*. 2021;9:647012.

1027. Leung DY. Atopic dermatitis: new insights and opportunities for therapeutic intervention. *J Allergy Clin Immunol*. 2000;105:860-876.
1028. Armstrong AW, Read C. Pathophysiology, clinical presentation, and treatment of psoriasis: a review. *JAMA*. 2020;323:1945-1960.
1029. Shao S, Fang H, Li Q, Wang G. Extracellular vesicles in inflammatory skin disorders: from pathophysiology to treatment. *Theranostics*. 2020;10:9937-9955.
1030. Lowes M, Suarez-Fariñas M, Krueger JG. Immunology of psoriasis. *Annu Rev Immunol*. 2014;32:227-255.
1031. Zhang Y, Yan J, Li Z, Zheng J, Sun Q. Exosomes derived from human umbilical cord mesenchymal stem cells alleviate psoriasis-like skin inflammation. *J Interf Cytokine Res*. 2022;42:8-18.
1032. Shen X, Song S, Chen N, Liao J, Zeng L. Stem cell-derived exosomes: a supernova in cosmetic dermatology. *J Cosmet Dermatol*. 2021;20:3812-3817.
1033. McCabe MC, Hill RC, Calderone K, et al. Alterations in extracellular matrix composition during aging and photoaging of the skin. *Matrix Biology Plus*. 2020;8:100041.
1034. Oh M, Lee J, Kim YJ, Rhee WJ, Park JH. Exosomes derived from human induced pluripotent stem cells ameliorate the aging of skin fibroblasts. *Int J Mol Sci*. 2018;19:1715.
1035. Zhong Q-Y, Lin B, Chen Y-T, et al. Gender differences in UV-induced skin inflammation, skin carcinogenesis and systemic damage. *Environ Toxicol Pharmacol*. 2021;81:103512.
1036. Varani J, Dame MK, Rittie L, et al. Decreased collagen production in chronologically aged skin: roles of age-dependent alteration in fibroblast function and defective mechanical stimulation. *Am J Pathol*. 2006;168:1861-1868.
1037. Byron A, Humphries JD, Humphries MJ. Defining the extracellular matrix using proteomics. *Int J Exp Pathol*. 2013;94:75-92.
1038. Hu S, Li Z, Cores J, et al. Needle-free injection of exosomes derived from human dermal fibroblast spheroids ameliorates skin photoaging. *ACS Nano*. 2019;13:11273-11282.
1039. Li L, Ngo HT, Hwang E, et al. Conditioned medium from human adipose-derived mesenchymal stem cell culture prevents UVB-induced skin aging in human keratinocytes and dermal fibroblasts. *Int J Mol Sci*. 2020;21:49.
1040. Tallant C, Marrero A, Gomis-Rüth FX. Matrix metalloproteinases: fold and function of their catalytic domains. *Biochim Biophys Acta*. 2010;1803:20-28.
1041. Juhl P, Bondesen S, Hawkins CL, et al. Dermal fibroblasts have different extracellular matrix profiles induced by TGF- β , PDGF and IL-6 in a model for skin fibrosis. *Sci Rep*. 2020;10:17300.
1042. Kim YJ, Yoo SM, Park HH, et al. Exosomes derived from human umbilical cord blood mesenchymal stem cells stimulates rejuvenation of human skin. *Biochem Biophys Res Commun*. 2017;493:1102-1108.
1043. Shi H, Wang M, Sun Y, Yang D, Xu W, Qian H. Exosomes: emerging cell-free based therapeutics in dermatologic diseases. *Front Cell Dev Biol*. 2021;9:736022.
1044. Zhang J, Chen C, Hu B, et al. Exosomes derived from human endothelial progenitor cells accelerate cutaneous wound healing by promoting angiogenesis through Erk1/2 signaling. *Int J Biol Sci*. 2016;12:1472-1487.
1045. Hu Y, Rao S-S, Wang Z-X, et al. Exosomes from human umbilical cord blood accelerate cutaneous wound healing through miR-21-3p-mediated promotion of angiogenesis and fibroblast function. *Theranostics*. 2018;8:169-184.
1046. Sideek MA, Teia A, Kopecki Z, Cowin AJ, Gibson MA. Co-localization of LTBP-2 with FGF-2 in fibrotic human keloid and hypertrophic scar. *J Mol Histol*. 2016;47:35-45.
1047. Zhao B, Zhang Y, Han S, et al. Exosomes derived from human amniotic epithelial cells accelerate wound healing and inhibit scar formation. *J Mol Histol*. 2017;48:121-132.
1048. Chen J, Zhou R, Liang Y, Fu X, Wang D, Wang C. Blockade of lncRNA-ASLNC5088-enriched exosome generation in M2 macrophages by GW4869 dampens the effect of M2 macrophages on orchestrating fibroblast activation. *FASEB J*. 2019;33:12200-12212.
1049. Xiong M, Zhang Q, Hu W, et al. The novel mechanisms and applications of exosomes in dermatology and cutaneous medical aesthetics. *Pharmacol Res*. 2021;166:105490.
1050. Gugliandolo A, Mazzone E. Dental mesenchymal stem cell secretome: an intriguing approach for neuroprotection and neuroregeneration. *Int J Mol Sci*. 2022;23:456.
1051. Ivica A, Zehnder M, Weber F. Therapeutic potential of mesenchymal stem cell-derived extracellular vesicles in regenerative endodontics. *Eur Cells Mater*. 2021;41:233-244.
1052. Mai Z, Chen H, Ye Y, et al. Translational and clinical applications of dental stem cell-derived exosomes. *Front Genet*. 2021;12:750990.
1053. Jafari N, Khoradmehr A, Moghiminasr R, Seyed Habashi M. Mesenchymal stromal/stem cells-derived exosomes as an antimicrobial weapon for Orofacial infections. *Front Microbiol*. 2022;12:4113.
1054. Nik Mohamed Kamal NNS, Shahidan WNS. Salivary exosomes: from waste to promising periodontitis treatment. *Front Physiol*. 2022;12:2336.
1055. Santonocito S, Polizzi A, Palazzo G, Isola G. The emerging role of microRNA in periodontitis: pathophysiology, clinical potential and future molecular perspectives. *Int J Mol Sci*. 2021;22:5456.
1056. Iwasaki K, Peng Y, Kanda R, Umeda M, Ishikawa I. Stem cell transplantation and cell-free treatment for periodontal regeneration. *Int J Mol Sci*. 2022;23:1011.
1057. Turksen K. *Cell Biology and Translational Medicine, Volume 12: Stem Cells in Development and Disease*. Springer Nature; 2021.
1058. Novello S, Pellen-Mussi P, Jeanne S. Mesenchymal stem cell-derived small extracellular vesicles as cell-free therapy: perspectives in periodontal regeneration. *J Periodontol Res*. 2021;56:433-442.
1059. Hua S, Bartold PM, Gulati K, Moran CS, Ivanovski S, Han P. Periodontal and dental pulp cell-derived small extracellular vesicles: a review of the current status. *Nanomaterials*. 2021;11:1858.
1060. Yin J-Y, Luo X-H, Feng W-Q, et al. Multidifferentiation potential of dental-derived stem cells. *World J Stem Cells*. 2021;13:342-365.
1061. Khodamoradi K, Golan R, Dullea A, Ramasamy R. Exosomes as potential biomarkers for erectile dysfunction, varicocele, and testicular injury. *Sex Med Rev*. 2022;10:311-322.
1062. Lötvall J, Hill AF, Hochberg F, et al. Minimal experimental requirements for definition of extracellular vesicles and their functions: a position statement from the International Society for Extracellular Vesicles. *J Extracell Vesicles*. 2014;3:26913.
1063. Kost Y, Muskat A, Mhaimeed N, Nazarian RS, Kobets K. Exosome therapy in hair regeneration: a literature review of the evidence, challenges, and future opportunities. *J Cosmet Dermatol*. 2022;21:3226-3231.
1064. Yuan A-R, Bian Q, Gao J-Q. Current advances in stem cell-based therapies for hair regeneration. *Eur J Pharmacol*. 2020;881:173197.
1065. Stevens J, Khetarpal S. Platelet-rich plasma for androgenetic alopecia: a review of the literature and proposed treatment protocol. *Int J Women's Dermatol*. 2019;5:46-51.
1066. Matsuzaki T, Yoshizato K. Role of hair papilla cells on induction and regeneration processes of hair follicles. *Wound Repair Regen*. 1998;6:524-530.
1067. Zhou L, Wang H, Jing J, Yu L, Wu X, Lu Z. Regulation of hair follicle development by exosomes derived from dermal papilla cells. *Biochem Biophys Res Commun*. 2018;500:325-332.
1068. Kwack MH, Seo CH, Gangadaran P, et al. Exosomes derived from human dermal papilla cells promote hair growth in cultured human hair follicles and augment the hair-inductive capacity of cultured dermal papilla spheres. *Exp Dermatol*. 2019;28:854-857.
1069. Deng W, Zhang Y, Wang W, et al. Hair follicle-derived mesenchymal stem cells decrease alopecia areata mouse hair loss and reduce inflammation around the hair follicle. *Stem Cell Res Ther*. 2021;12:548.

1070. Driskell RR, Clavel C, Rendl M, Watt FM. Hair follicle dermal papilla cells at a glance. *J Cell Sci.* 2011;124:1179-1182.
1071. Yan H, Gao Y, Ding Q, et al. Exosomal micro RNAs derived from dermal papilla cells mediate hair follicle stem cell proliferation and differentiation. *Int J Biol Sci.* 2019;15:1368-1382.
1072. Kazi T, Nagata A, Nakagawa T, Matsuzaki T, Inui S. Dermal papilla cell-derived extracellular vesicles increase hair inductive gene expression in adipose stem cells via β -catenin activation. *Cells.* 2022;11:202.
1073. Nilforoushzadeh MA, Aghdami N, Taghiabadi E. Human hair outer root sheath cells and platelet-lysis exosomes promote hair inductivity of dermal papilla cell. *Tissue Eng Regen Med.* 2020;17:525-536.
1074. Zöller M, Zhao K, Kutlu N, et al. Immunoregulatory effects of myeloid-derived suppressor cell exosomes in mouse model of autoimmune alopecia areata. *Front Immunol.* 2018;9:1279.
1075. Rajendran RL, Gangadaran P, Seo CH, et al. Macrophage-derived extracellular vesicle promotes hair growth. *Cells.* 2020;9:856.
1076. Yoon YC, Ahn BH, Min JW, Lee KR, Park SH, Kang HC. Stimulatory effects of extracellular vesicles derived from *Leuconostoc holzapfelii* that exists in human scalp on hair growth in human follicle dermal papilla cells. *Curr Issues Mol Biol.* 2022;44:845-866.
1077. Riche A, Aberdam E, Marchand L, et al. Extracellular vesicles from activated dermal fibroblasts stimulate hair follicle growth through dermal papilla-secreted norrin. *Stem Cells.* 2019;37:1166-1175.
1078. Aziz I, Che Ramli MD, Mohd Zain NS, Sanusi J. Behavioral and histopathological study of changes in spinal cord injured rats supplemented with *Spirulina platensis*. *Evid Based Complement Alternat Med.* 2014;2014:871657.
1079. Sykova E, Cizkova D, Kubinova S. Mesenchymal stem cells in treatment of spinal cord injury and amyotrophic lateral sclerosis. *Front Cell Dev Biol.* 2021;9:695900.
1080. Kim G-U, Sung S-E, Kang K-K, et al. Therapeutic potential of mesenchymal stem cells (MSCs) and MSC-derived extracellular vesicles for the treatment of spinal cord injury. *Int J Mol Sci.* 2021;22:13672.
1081. Bhat IA, Sivanarayanan TB, Somal A, et al. An allogenic therapeutic strategy for canine spinal cord injury using mesenchymal stem cells. *J Cell Physiol.* 2019;234:2705-2718.
1082. Vismara I, Papa S, Rossi F, Forloni G, Veglianese P. Current options for cell therapy in spinal cord injury. *Trends Mol Med.* 2017;23:831-849.
1083. Liao LL, Looi QH, Chia WC, Subramaniam T, Ng MH, Law JX. Treatment of spinal cord injury with mesenchymal stem cells. *Cell Biosci.* 2020;10:112.
1084. Singh A, Tetreault L, Kalsi-Ryan S, Nouri A, Fehlings MG. Global prevalence and incidence of traumatic spinal cord injury. *Clin Epidemiol.* 2014;6:309-331.
1085. Kumar R, Lim J, Mekary RA, et al. Traumatic spinal injury: global epidemiology and worldwide volume. *World Neurosurg.* 2018;113:e345-e363.
1086. Witiw CD, Fehlings MG. Acute spinal cord injury. *J Spinal Disord Tech.* 2015;28:202-210.
1087. Morales I-I, Toscano-Tejeda D, Ibarra A. Non pharmacological strategies to promote spinal cord regeneration: a view on some individual or combined approaches. *Curr Pharm Des.* 2016;22:720-727.
1088. Tyler JY, Xu X-M, Cheng J-X. Nanomedicine for treating spinal cord injury. *Nanoscale.* 2013;5:8821-8836.
1089. Wang J, Pearse DD. Therapeutic hypothermia in spinal cord injury: the status of its use and open questions. *Int J Mol Sci.* 2015;16:16848-16879.
1090. Ramer L, Ramer M, Steeves J. Setting the stage for functional repair of spinal cord injuries: a cast of thousands. *Spinal Cord.* 2005;43:134-161.
1091. Rosenzweig ES, McDonald JW. Rodent models for treatment of spinal cord injury: research trends and progress toward useful repair. *Curr Opin Neurol.* 2004;17:121-131.
1092. Kumar H, Ropper AE, Lee S-H, Han I. Propitious therapeutic modulators to prevent blood-spinal cord barrier disruption in spinal cord injury. *Mol Neurobiol.* 2017;54:3578-3590.
1093. Carlson GD, Gorden C. Current developments in spinal cord injury research. *Spine J.* 2002;2:116-128.
1094. Hayta E, Elden H. Acute spinal cord injury: a review of pathophysiology and potential of non-steroidal anti-inflammatory drugs for pharmacological intervention. *J Chem Neuroanat.* 2018;87:25-31.
1095. Tator CH, Fehlings MG. Review of the secondary injury theory of acute spinal cord trauma with emphasis on vascular mechanisms. *J Neurosurg.* 1991;75:15-26.
1096. Fehlings MG, Hawryluk GW. Scarring after spinal cord injury. *J Neurosurg Spine.* 2010;13:165-167.
1097. Kwiecien JM, Dabrowski W, Dąbrowska-Bouta B, et al. Prolonged inflammation leads to ongoing damage after spinal cord injury. *PLoS One.* 2020;15:e0226584.
1098. Wilson JR, Forgione N, Fehlings MG. Emerging therapies for acute traumatic spinal cord injury. *CMAJ.* 2013;185:485-492.
1099. Dvorak MF, Noonan VK, Fallah N, et al. The influence of time from injury to surgery on motor recovery and length of hospital stay in acute traumatic spinal cord injury: an observational Canadian cohort study. *J Neurotrauma.* 2015;32:645-654.
1100. Burke JF, Yue JK, Ngwenya LB, et al. Ultra-early (<12 hours) surgery correlates with higher rate of American spinal injury association impairment scale conversion after cervical spinal cord injury. *Neurosurgery.* 2019;85:199-203.
1101. Fehlings MG, Vaccaro A, Wilson JR, et al. Early versus delayed decompression for traumatic cervical spinal cord injury: results of the surgical timing in acute spinal cord injury study (STASCIS). *PLoS One.* 2012;7:e32037.
1102. Wutte C, Klein B, Becker J, et al. Earlier decompression (<8 hours) results in better neurological and functional outcome after traumatic thoracolumbar spinal cord injury. *J Neurotrauma.* 2019;36:2020-2027.
1103. Rath N, Balain B. Spinal cord injury—the role of surgical treatment for neurological improvement. *J Clin Orthop Trauma.* 2017;8:99-102.
1104. Ahuja CS, Wilson JR, Nori S, et al. Traumatic spinal cord injury. *Nat Rev Dis Primers.* 2017;3:17018.
1105. Streijger F, So K, Manouchehri N, et al. A direct comparison between norepinephrine and phenylephrine for augmenting spinal cord perfusion in a porcine model of spinal cord injury. *J Neurotrauma.* 2018;35:1345-1357.
1106. Hawryluk G, Whetstone W, Saigal R, et al. Mean arterial blood pressure correlates with neurological recovery after human spinal cord injury: analysis of high frequency physiologic data. *J Neurotrauma.* 2015;32:1958-1967.
1107. Ryken TC, Hurlbert RJ, Hadley MN, et al. The acute cardiopulmonary management of patients with cervical spinal cord injuries. *Neurosurgery.* 2013;72:84-92.
1108. Seki T, Fehlings MG. Mechanistic insights into posttraumatic syringomyelia based on a novel in vivo animal model. *J Neurosurg Spine.* 2008;8:365-375.
1109. Kwiecien JM, Jarosz B, Oakden W, et al. An in vivo model of anti-inflammatory activity of subdural dexamethasone following the spinal cord injury. *Neurol Neurochir Pol.* 2016;50:7-15.
1110. Kwiecien JM, Jarosz B, Urdzikova LM, Rola R, Dabrowski W. Subdural infusion of dexamethasone inhibits leukomyelitis after acute spinal cord injury in a rat model. *Folia Neuropathol.* 2015;53:41-51.
1111. Hu R, Zhou J, Luo C, et al. Glial scar and neuroregeneration: histological, functional, and magnetic resonance imaging analysis in chronic spinal cord injury. *J Neurosurg Spine.* 2010;13:169-180.
1112. Sofroniew MV. Molecular dissection of reactive astrogliosis and glial scar formation. *Trends Neurosci.* 2009;32:638-647.
1113. Kwiecien JM. Cellular mechanisms of white matter regeneration in an adult dysmyelinated rat model. *Folia Neuropathol.* 2013;51:189-202.

1114. Jain NB, Ayers GD, Peterson EN, et al. Traumatic spinal cord injury in the United States, 1993–2012. *JAMA*. 2015;313:2236–2243.
1115. Alizadeh A, Dyck SM, Karimi-Abdolrezaee S. Traumatic spinal cord injury: an overview of pathophysiology, models and acute injury mechanisms. *Front Neurol*. 2019;10:282.
1116. Liddelow SA, Barres BA. Reactive astrocytes: production, function, and therapeutic potential. *Immunity*. 2017;46:957–967.
1117. Martinez FO, Gordon S. The M1 and M2 paradigm of macrophage activation: time for reassessment. *F1000prime Rep*. 2014;6:13.
1118. Sofroniew M. Astrogliosis. *Cold Spring Harb Perspect Biol*. 2014;7:a020420.
1119. Anderson MA, Burda JE, Ren Y, et al. Astrocyte scar formation aids central nervous system axon regeneration. *Nature*. 2016;532:195–200.
1120. Sofroniew MV, Vinters HV. Astrocytes: biology and pathology. *Acta Neuropathol*. 2010;119:7–35.
1121. Liddelow SA, Guttenplan KA, Clarke LE, et al. Neurotoxic reactive astrocytes are induced by activated microglia. *Nature*. 2017;541:481–487.
1122. Herbert FJ, Bharathi D, Suresh S, David E, Kumar S. Regenerative potential of stem cell-derived extracellular vesicles in spinal cord injury (SCI). *Curr Stem Cell Res Ther*. 2022;17:280–293.
1123. Zhao C, Zhou X, Qiu J, et al. Exosomes derived from bone marrow mesenchymal stem cells inhibit complement activation in rats with spinal cord injury. *Drug Des Devel Ther*. 2019;13:3693–3704.
1124. Lee J-R, Kyung JW, Kumar H, et al. Targeted delivery of mesenchymal stem cell-derived nanovesicles for spinal cord injury treatment. *Int J Mol Sci*. 2020;21:4185.
1125. Li C, Jiao G, Wu W, et al. Exosomes from bone marrow mesenchymal stem cells inhibit neuronal apoptosis and promote motor function recovery via the Wnt/ β -catenin signaling pathway. *Cell Transplant*. 2019;28:1373–1383.
1126. Rong Y, Liu W, Wang J, et al. Neural stem cell-derived small extracellular vesicles attenuate apoptosis and neuroinflammation after traumatic spinal cord injury by activating autophagy. *Cell Death Dis*. 2019;10:340.
1127. Yeng C-H, Chen P-J, Chang H-K, et al. Attenuating spinal cord injury by conditioned medium from human umbilical cord blood-derived CD34+ cells in rats. *Taiwan J Obstet Gynecol*. 2016;55:85–93.
1128. Ijaz S, Mohammed I, Gholaminejad M, Mokhtari T, Akbari M, Hassanzadeh G. Modulating pro-inflammatory cytokines, tissue damage magnitude, and motor deficit in spinal cord injury with subventricular zone-derived extracellular vesicles. *J Mol Neurosci*. 2020;70:458–466.
1129. Wang L, Pei S, Han L, et al. Mesenchymal stem cell-derived exosomes reduce A1 astrocytes via downregulation of phosphorylated NF κ B P65 subunit in spinal cord injury. *Cell Physiol Biochem*. 2018;50:1535–1559.
1130. Luo Y, Xu T, Liu W, et al. Exosomes derived from GIT1-overexpressing bone marrow mesenchymal stem cells promote traumatic spinal cord injury recovery in a rat model. *Int J Neurosci*. 2021;131:170–182.
1131. Huang J-H, Fu C-H, Xu Y, Yin X-M, Cao Y, Lin F-Y. Extracellular vesicles derived from epidural fat-mesenchymal stem cells attenuate NLRP3 inflammasome activation and improve functional recovery after spinal cord injury. *Neurochem Res*. 2020;45:760–771.
1132. de Munter J, Beugels J, Munter S, et al. Standardized human bone marrow-derived stem cells infusion improves survival and recovery in a rat model of spinal cord injury. *J Neurol Sci*. 2019;402:16–29.
1133. Lu Y, Zhou Y, Zhang R, et al. Bone mesenchymal stem cell-derived extracellular vesicles promote recovery following spinal cord injury via improvement of the integrity of the blood-spinal cord barrier. *Front Neurosci*. 2019;13:209.
1134. Ma K, Xu H, Zhang J, et al. Insulin-like growth factor-1 enhances neuroprotective effects of neural stem cell exosomes after spinal cord injury via an miR-219a-2-3p/YY1 mechanism. *Aging (Albany NY)*. 2019;11:12278–12294.
1135. Xu G, Ao R, Zhi Z, Jia J, Yu B. miR-21 and miR-19b delivered by hMSC-derived EVs regulate the apoptosis and differentiation of neurons in patients with spinal cord injury. *J Cell Physiol*. 2019;234:10205–10217.
1136. Zhou X, Chu X, Yuan H, et al. Mesenchymal stem cell derived EVs mediate neuroprotection after spinal cord injury in rats via the microRNA-21-5p/FasL gene axis. *Biomed Pharmacother*. 2019;115:108818.
1137. Gu J, Jin ZS, Wang CM, Yan XF, Mao YQ, Chen S. Bone marrow mesenchymal stem cell-derived exosomes improves spinal cord function after injury in rats by activating autophagy. *Drug Des Devel Ther*. 2020;14:1621–1631.
1138. Li C, Li X, Zhao B, Wang C. Exosomes derived from miR-544-modified mesenchymal stem cells promote recovery after spinal cord injury. *Arch Physiol Biochem*. 2020;126:369–375.
1139. Li D, Zhang P, Yao X, et al. Exosomes derived from miR-133b-modified mesenchymal stem cells promote recovery after spinal cord injury. *Front Neurosci*. 2018;12:845.
1140. Chudickova M, Vackova I, Machova Urdzikova L, et al. The effect of Wharton jelly-derived mesenchymal stromal cells and their conditioned media in the treatment of a rat spinal cord injury. *Int J Mol Sci*. 2019;20:4516.
1141. Li L, Zhang Y, Mu J, et al. Transplantation of human mesenchymal stem-cell-derived exosomes immobilized in an adhesive hydrogel for effective treatment of spinal cord injury. *Nano Lett*. 2020;20:4298–4305.
1142. Liu W, Rong Y, Wang J, et al. Exosome-shuttled miR-216a-5p from hypoxic preconditioned mesenchymal stem cells repair traumatic spinal cord injury by shifting microglial M1/M2 polarization. *J Neuroinflammation*. 2020;17:47.
1143. Kim D-k, Nishida H, An SY, Shetty AK, Bartosh TJ, Prockop DJ. Chromatographically isolated CD63+ CD81+ extracellular vesicles from mesenchymal stromal cells rescue cognitive impairments after TBI. *Proc Natl Acad Sci*. 2016;113:170–175.
1144. Yu T, Zhao C, Hou S, Zhou W, Wang B, Chen Y. Exosomes secreted from miRNA-29b-modified mesenchymal stem cells repaired spinal cord injury in rats. *Braz J Med Biol Res*. 2019;52:e8735.
1145. Niu Y, Xia X, Song P, et al. Bone mesenchymal stem cell-conditioned medium attenuates the effect of oxidative stress injury on NSCs by inhibiting the Notch1 signaling pathway. *Cell Biol Int*. 2019;43:1267–1275.
1146. Sun G, Li G, Li D, et al. hucMSC derived exosomes promote functional recovery in spinal cord injury mice via attenuating inflammation. *Mater Sci Eng C*. 2018;89:194–204.
1147. Huang J-H, Xu Y, Yin X-M, Lin F-Y. Exosomes derived from miR-126-modified MSCs promote angiogenesis and neurogenesis and attenuate apoptosis after spinal cord injury in rats. *Neuroscience*. 2020;424:133–145.
1148. Guo S, Redenski I, Levenberg S. Spinal cord repair: from cells and tissue engineering to extracellular vesicles. *Cells*. 2021;10:1872.
1149. Huang J-H, Yin X-M, Xu Y, et al. Systemic administration of exosomes released from mesenchymal stromal cells attenuates apoptosis, inflammation, and promotes angiogenesis after spinal cord injury in rats. *J Neurotrauma*. 2017;34:3388–3396.
1150. Liu W, Wang Y, Gong F, et al. Exosomes derived from bone mesenchymal stem cells repair traumatic spinal cord injury by suppressing the activation of A1 neurotoxic reactive astrocytes. *J Neurotrauma*. 2019;36:469–484.
1151. Ruppert KA, Nguyen TT, Prabhakara KS, et al. Human mesenchymal stromal cell-derived extracellular vesicles modify microglial response and improve clinical outcomes in experimental spinal cord injury. *Sci Rep*. 2018;8:480.
1152. Mohammed I, Ijaz S, Mokhtari T, et al. Subventricular zone-derived extracellular vesicles promote functional recovery in rat model of spinal cord injury by inhibition of NLRP3 inflammasome complex formation. *Metab Brain Dis*. 2020;35:809–818.

1153. Rong Y, Liu W, Lv C, et al. Neural stem cell small extracellular vesicle-based delivery of 14-3-3t reduces apoptosis and neuroinflammation following traumatic spinal cord injury by enhancing autophagy by targeting Beclin-1. *Aging (Albany NY)*. 2019;11:7723-7745.
1154. Yuan X, Wu Q, Wang P, et al. Exosomes derived from pericytes improve microcirculation and protect blood-spinal cord barrier after spinal cord injury in mice. *Front Neurosci*. 2019;13:319.
1155. Guo S, Perets N, Betzer O, et al. Intranasal delivery of mesenchymal stem cell derived exosomes loaded with phosphatase and tensin homolog siRNA repairs complete spinal cord injury. *ACS Nano*. 2019;13:10015-10028.
1156. Liu W-z, Ma Z-j, Li J-r, Kang X-w. Mesenchymal stem cell-derived exosomes: therapeutic opportunities and challenges for spinal cord injury. *Stem Cell Res Ther*. 2021;12:102.
1157. Katz JN. Lumbar disc disorders and low-back pain: socioeconomic factors and consequences. *JBJS*. 2006;88:21-24.
1158. Gague E, Turci F, Newman E, et al. 3D assessment of intervertebral disc degeneration in zebrafish identifies changes in bone density that prime disc disease. *Bone Res*. 2021;9:39.
1159. Park J-H, Choi K-E, Kim S-G, et al. Long-term follow-up of inpatients with failed back surgery syndrome who received integrative Korean medicine treatment: a retrospective analysis and questionnaire survey study. *J Clin Med*. 2021;10:1703.
1160. Deyo RA, Weinstein JN. Low back pain affects men and women equally, with onset most often between the ages of 30 and 50 years. It is the most common cause of work-related disability in people under 45 years of age and the most expensive. *N Engl J Med*. 2001;344:363-370.
1161. Balagué F, Mannion AF, Pellisé F, Cedraschi C. Non-specific low back pain. *Lancet*. 2012;379:482-491.
1162. Binch AL, Fitzgerald JC, Growney EA, Barry F. Cell-based strategies for IVD repair: clinical progress and translational obstacles. *Nat Rev Rheumatol*. 2021;17:158-175.
1163. Liao Z, Luo R, Li G, et al. Exosomes from mesenchymal stem cells modulate endoplasmic reticulum stress to protect against nucleus pulposus cell death and ameliorate intervertebral disc degeneration in vivo. *Theranostics*. 2019;9:4084-4100.
1164. Au TY, Lam TK, Peng Y, et al. Transformation of resident notochord-descendent nucleus pulposus cells in mouse injury-induced fibrotic intervertebral discs. *Aging Cell*. 2020;19:e13254.
1165. Croft AS, Illien-Jünger S, Grad S, Guerrero J, Wangler S, Gantenbein B. The application of mesenchymal stromal cells and their homing capabilities to regenerate the intervertebral disc. *Int J Mol Sci*. 2021;22:3519.
1166. DiStefano TJ, Vaso K, Danias G, Chionuma HN, Weiser JR, Iatridis JC. Extracellular vesicles as an emerging treatment option for intervertebral disc degeneration: therapeutic potential, translational pathways, and regulatory considerations. *Adv Healthc Mater*. 2022;11:2100596.
1167. Melrose J. Strategies in regenerative medicine for intervertebral disc repair using mesenchymal stem cells and bioscaffolds. *Regen Med*. 2016;11:705-724.
1168. Richardson SM, Kalamegam G, Pushparaj PN, et al. Mesenchymal stem cells in regenerative medicine: focus on articular cartilage and intervertebral disc regeneration. *Methods*. 2016;99:69-80.
1169. Vadalà G, Russo F, Ambrosio L, Papalia R, Denaro V. Mesenchymal stem cells for intervertebral disc regeneration. *J Biol Regul Homeost Agents*. 2016;30:173-179.
1170. Loibl M, Wuertz-Kozak K, Vadala G, Lang S, Fairbank J, Urban JP. Controversies in regenerative medicine: should intervertebral disc degeneration be treated with mesenchymal stem cells? *JOR Spine*. 2019;2:e1043.
1171. Mohd Noor NA, Abdullah Nurul A, Ahmad Mohd Zain MR, Wan Nor Aduni WK, Azlan M. Extracellular vesicles from mesenchymal stem cells as potential treatments for osteoarthritis. *Cells*. 2021;10:1287.
1172. Lu L, Xu A, Gao F, et al. Mesenchymal stem cell-derived exosomes as a novel strategy for the treatment of intervertebral disc degeneration. *Front Cell Dev Biol*. 2022;9:770510.
1173. Qi L, Wang R, Shi Q, Yuan M, Jin M, Li D. Umbilical cord mesenchymal stem cell conditioned medium restored the expression of collagen II and aggrecan in nucleus pulposus mesenchymal stem cells exposed to high glucose. *J Bone Miner Metab*. 2019;37:455-466.
1174. Li M, Li R, Yang S, et al. Exosomes derived from bone marrow mesenchymal stem cells prevent acidic pH-induced damage in human nucleus pulposus cells. *Med Sci Monit*. 2020;26:e922928.
1175. Li Z-q, Kong L, Liu C, Xu H-G. Human bone marrow mesenchymal stem cell-derived exosomes attenuate IL-1 β -induced annulus fibrosus cell damage. *Am J Med Sci*. 2020;360:693-700.
1176. Yuan Q, Wang X, Liu L, et al. Exosomes derived from human placental mesenchymal stromal cells carrying antagomiR-4450 alleviate intervertebral disc degeneration through upregulation of ZNF121. *Stem Cells Dev*. 2020;29:1038-1058.
1177. Zhu G, Yang X, Peng C, Yu L, Hao Y. Exosomal miR-532-5p from bone marrow mesenchymal stem cells reduce intervertebral disc degeneration by targeting RASSF5. *Exp Cell Res*. 2020;393:112109.
1178. Wen T, Wang H, Li Y, et al. Bone mesenchymal stem cell-derived extracellular vesicles promote the repair of intervertebral disc degeneration by transferring microRNA-199a. *Cell Cycle*. 2021;20:256-270.
1179. Zhu L, Shi Y, Liu L, Wang H, Shen P, Yang H. Mesenchymal stem cells-derived exosomes ameliorate nucleus pulposus cells apoptosis via delivering miR-142-3p: therapeutic potential for intervertebral disc degenerative diseases. *Cell Cycle*. 2020;19:1727-1739.
1180. Cheng X, Zhang G, Zhang L, et al. Mesenchymal stem cells deliver exogenous miR-21 via exosomes to inhibit nucleus pulposus cell apoptosis and reduce intervertebral disc degeneration. *J Cell Mol Med*. 2018;22:261-276.
1181. Zhang J, Zhang J, Zhang Y, et al. Mesenchymal stem cells-derived exosomes ameliorate intervertebral disc degeneration through inhibiting pyroptosis. *J Cell Mol Med*. 2020;24:11742-11754.
1182. Hingert D, Ekström K, Aldridge J, Crescitelli R, Brisby H. Extracellular vesicles from human mesenchymal stem cells expedite chondrogenesis in 3D human degenerative disc cell cultures. *Stem Cell Res Ther*. 2020;11:323.
1183. Sun Y, Zhang W, Li X. Induced pluripotent stem cell-derived mesenchymal stem cells deliver exogenous miR-105-5p via small extracellular vesicles to rejuvenate senescent nucleus pulposus cells and attenuate intervertebral disc degeneration. *Stem Cell Res Ther*. 2021;12:286.
1184. Lu K, Li H-y, Yang K, et al. Exosomes as potential alternatives to stem cell therapy for intervertebral disc degeneration: in vitro study on exosomes in interaction of nucleus pulposus cells and bone marrow mesenchymal stem cells. *Stem Cell Res Ther*. 2017;8:108.
1185. Xia C, Zeng Z, Fang B, et al. Mesenchymal stem cell-derived exosomes ameliorate intervertebral disc degeneration via anti-oxidant and anti-inflammatory effects. *Free Radic Biol Med*. 2019;143:1-15.
1186. Bari E, Perteghella S, Di Silvestre D, et al. Pilot production of mesenchymal stem/stromal freeze-dried secretome for cell-free regenerative nanomedicine: a validated GMP-compliant process. *Cells*. 2018;7:190.
1187. Xie L, Chen Z, Liu M, et al. MSC-derived exosomes protect vertebral endplate chondrocytes against apoptosis and calcification via the miR-31-5p/ATF6 axis. *Mol Ther Nucleic Acids*. 2020;22:601-614.
1188. Tracy EP, Stielberg V, Rowe G, et al. State of the field: cellular and exosomal therapeutic approaches in vascular regeneration. *Am J Physiol Heart Circ Physiol*. 2022;322:H647-H680.
1189. Benton RL, Maddie MA, Gruenthal MJ, Hagg T, Whittemore SR. Neutralizing endogenous VEGF following traumatic spinal cord

- injury modulates microvascular plasticity but not tissue sparing or functional recovery. *Curr Neurovasc Res.* 2009;6:124-131.
1190. Fassbender JM, Whittmore SR, Hagg T. Targeting microvasculature for neuroprotection after SCI. *Neurotherapeutics.* 2011;8:240-251.
 1191. Weyand CM, Goronzy JJ. Medium-and large-vessel vasculitis. *N Engl J Med.* 2003;349:160-169.
 1192. Seo P, Stone JH. Small-vessel and medium-vessel vasculitis. *Arthritis Care Res.* 2007;57:1552-1559.
 1193. Langford C, Vasculitis J. View PDF view article view in Scopus. *Allergy Clin Immunol.* 2010;125(2 Suppl 2):S216-S225.
 1194. Basu N, Karabayas M, Pusey C. Prognosis and future developments in vasculitis. *Best Pract Res Clin Rheumatol.* 2018;32:148-165.
 1195. Yang N, Zhao Y, Wu X, et al. Recent advances in extracellular vesicles and their involvements in vasculitis. *Free Radic Biol Med.* 2021;171:203-218.
 1196. Shi Y, Du L, Lv D, et al. Emerging role and therapeutic application of exosome in hepatitis virus infection and associated diseases. *J Gastroenterol.* 2021;56:336-349.
 1197. Sharma A, Ahmad Farouk I, Lal SK. COVID-19: a review on the novel coronavirus disease evolution, transmission, detection, control and prevention. *Viruses.* 2021;13:202.
 1198. You J, Fu Z, Zou L. Mechanism and potential of extracellular vesicles derived from mesenchymal stem cells for the treatment of infectious diseases. *Front Microbiol.* 2021;12:761338.
 1199. Fauci AS, Lane HC, Redfield RR. Covid-19—Navigating the Uncharted. *N Engl J Med.* 2020;382:1268-1269.
 1200. Velavan T, Meyer C. The COVID-19 epidemic. *Trop Med Int Health.* 2020;25(3):278-280.
 1201. Thanh Le T, Andreadakis Z, Kumar A, et al. The COVID-19 vaccine development landscape. *Nat Rev Drug Discov.* 2020;19:305-306.
 1202. Cao X. COVID-19: immunopathology and its implications for therapy. *Nat Rev Immunol.* 2020;20:269-270.
 1203. Babaei G, Zare N, Mihanfar A, Ansari MHK. Exosomes and COVID-19: challenges and opportunities. *Comp Clin Pathol.* 2022;31:347-354.
 1204. Hu J, Stojanović J, Yasamineh S, et al. The potential use of microRNAs as a therapeutic strategy for SARS-CoV-2 infection. *Arch Virol.* 2021;166:2649-2672.
 1205. Hosseini NF, Dalirfardouei R, Aliramaei MR, Najafi R. Stem cells or their exosomes: which is preferred in COVID-19 treatment? *Biotechnol Lett.* 2022;44:159-177.
 1206. Gupta S, Krishnakumar V, Sharma Y, Dinda AK, Mohanty S. Mesenchymal stem cell derived exosomes: a nano platform for therapeutics and drug delivery in combating COVID-19. *Stem Cell Rev Rep.* 2021;17:33-43.
 1207. Golchin A. Cell-based therapy for severe COVID-19 patients: clinical trials and cost-utility. *Stem Cell Rev Rep.* 2021;17:56-62.
 1208. Jamshidi E, Babajani A, Soltani P, Niknejad H. Proposed mechanisms of targeting COVID-19 by delivering mesenchymal stem cells and their exosomes to damaged organs. *Stem Cell Rev Rep.* 2021;17:176-192.
 1209. Jamalkhah M, Asaadi Y, Azangou-Khyavy M, et al. MSC-derived exosomes carrying a cocktail of exogenous interfering RNAs an unprecedented therapy in era of COVID-19 outbreak. *J Transl Med.* 2021;19:164.
 1210. Wang W, Lei W, Jiang L, et al. Therapeutic mechanisms of mesenchymal stem cells in acute respiratory distress syndrome reveal potentials for Covid-19 treatment. *J Transl Med.* 2021;19:198.
 1211. Abdelgawad M, Bakry NS, Farghali AA, Abdel-Latif A, Lotfy A. Mesenchymal stem cell-based therapy and exosomes in COVID-19: current trends and prospects. *Stem Cell Rev Rep.* 2021;12:469.
 1212. Krishnan A, Muthusamy S, Fernandez FB, Kasoju N. Mesenchymal stem cell-derived extracellular vesicles in the management of COVID19-associated lung injury: a review on publications, clinical trials and patent landscape. *Tissue Eng Regen Med.* 2022;19:659-673.
 1213. Karn V, Ahmed S, Tsai L-W, et al. Extracellular vesicle-based therapy for COVID-19: promises, challenges and future prospects. *Biomedicine.* 2021;9:1373.
 1214. Raghav A, Khan ZA, Upadhyay VK, et al. Mesenchymal stem cell-derived exosomes exhibit promising potential for treating SARS-CoV-2-infected patients. *Cells.* 2021;10:587.
 1215. Borowiec BM, Angelova Volponi A, Mozdziak P, Kempisty B, Dyszkiewicz-Konwińska M. Small extracellular vesicles and COVID19—using the “Trojan horse” to tackle the giant. *Cells.* 2021;10:3383.
 1216. Georgescu A, Simionescu M. Extracellular vesicles: versatile nanomediators, potential biomarkers and therapeutic agents in atherosclerosis and COVID-19-related thrombosis. *Int J Mol Sci.* 2021;22:5967.
 1217. Kiaie N, Ghanavati SPM, Miremadi SS, Hadipour A, Aghdam RM. Mesenchymal stem cell-derived exosomes for COVID-19 therapy, preclinical and clinical evidence. *Int J Stem Cells.* 2021;14:252-261.
 1218. Dauletova M, Hafsan H, Mahhngam N, Zekiy AO, Ahmadi M, Siahmansouri H. Mesenchymal stem cell alongside exosomes as a novel cell-based therapy for COVID-19: a review study. *Clin Immunol.* 2021;226:108712.
 1219. Dubey A, Lobo CL, Ravi G, Shetty A, Hebbar S, El-Zahaby SA. Exosomes: emerging implementation of nanotechnology for detecting and managing novel corona virus-SARS-CoV-2. *Asian J Pharm Sci.* 2022;17:20-34.
 1220. Farkhad NK, Mahmoudi A, Mahdipour E. Regenerative therapy by using mesenchymal stem cells-derived exosomes in COVID-19 treatment. The potential role and underlying mechanisms. *Regen Ther.* 2022;20:61-71.
 1221. Yousefi Dehbid M, Goodarzi N, Azhdari MH, Doroudian M. Mesenchymal stem cells and their derived exosomes to combat Covid-19. *Rev Med Virol.* 2022;32:e2281.
 1222. Popowski KD, Dinh PUC, George A, Lutz H, Cheng K. Exosome therapeutics for COVID-19 and respiratory viruses. *Viewpoints.* 2021;2:20200186.
 1223. Kuate S, Cinatl J, Doerr HW, Überla K. Exosomal vaccines containing the S protein of the SARS coronavirus induce high levels of neutralizing antibodies. *Virology.* 2007;362:26-37.
 1224. Sengupta V, Sengupta S, Lazo A, Woods P, Nolan A, Bremer N. Exosomes derived from bone marrow mesenchymal stem cells as treatment for severe COVID-19. *Stem Cells Dev.* 2020;29:747-754.
 1225. Rodriguez A-M, Nakhle J, Griessinger E, Vignais M-L. Intercellular mitochondria trafficking highlighting the dual role of mesenchymal stem cells as both sensors and rescuers of tissue injury. *Cell Cycle.* 2018;17:712-721.
 1226. Park J, Kim S, Lim H, et al. Therapeutic effects of human mesenchymal stem cell microvesicles in an ex vivo perfused human lung injured with severe *E. coli* pneumonia. *Thorax.* 2019;74:43-50.
 1227. Vats D, Mukundan L, Odegaard JI, et al. Oxidative metabolism and PGC-1 β attenuate macrophage-mediated inflammation. *Cell Metab.* 2006;4:13-24.
 1228. Phinney DG, Di Giuseppe M, Njah J, et al. Mesenchymal stem cells use extracellular vesicles to outsource mitophagy and shuttle microRNAs. *Nat Commun.* 2015;6:8472.
 1229. Morrison TJ, Jackson MV, Cunningham EK, et al. Mesenchymal stromal cells modulate macrophages in clinically relevant lung injury models by extracellular vesicle mitochondrial transfer. *Am J Respir Crit Care Med.* 2017;196:1275-1286.
 1230. Wei Li J, Wei L, Han Z, Chen Z. Mesenchymal stromal cells-derived exosomes alleviate ischemia/reperfusion injury in mouse lung by transporting anti-apoptotic miR-21-5p. *Eur J Pharmacol.* 2019;852:68-76.
 1231. Bari E, Ferrarotti I, Di Silvestre D, et al. Adipose mesenchymal extracellular vesicles as alpha-1-antitrypsin physiological delivery systems for lung regeneration. *Cells.* 2019;8:965.

1232. Li G, Fan Y, Lai Y, et al. Coronavirus infections and immune responses. *J Med Virol*. 2020;92:424-432.
1233. Pedersen SF, Ho Y-C. SARS-CoV-2: a storm is raging. *J Clin Invest*. 2020;130:2202-2205.
1234. Henderson LA, Canna SW, Schulert GS, et al. On the alert for cytokine storm: immunopathology in COVID-19. *Arthritis Rheumatol*. 2020;72:1059-1063.
1235. Mehta P, McAuley DF, Brown M, Sanchez E, Tattersall RS, Manson JJ. COVID-19: consider cytokine storm syndromes and immunosuppression. *Lancet*. 2020;395:1033-1034.
1236. Piper C, Drobyski WR. Inflammatory cytokine networks in gastrointestinal tract graft vs. host disease. *Front Immunol*. 2019;10:163.
1237. Kamel AM, Elsharkawy NM, Abdelfattah EK, et al. IL12 and IFN γ secretion by donor mononuclear cells in response to host antigens may predict acute GVHD after HSCT. *Immunobiology*. 2019;224:659-665.
1238. Kim J, Yang YL, Jeong Y, Jang Y-S. Middle East respiratory syndrome-coronavirus infection into established hDPP4-transgenic mice accelerates lung damage via activation of the pro-inflammatory response and pulmonary fibrosis. *J Microbiol Biotechnol*. 2020;30:427.
1239. Zeng SL, Wang LH, Li P, Wang W, Yang J. Mesenchymal stem cells abrogate experimental asthma by altering dendritic cell function. *Mol Med Rep*. 2015;12:2511-2520.
1240. Herrero R, Sanchez G, Lorente JA. New insights into the mechanisms of pulmonary edema in acute lung injury. *Ann Transl Med*. 2018;6:32.
1241. Gennai S, Monsel A, Hao Q, Park J, Matthay M, Lee J. Microvesicles derived from human mesenchymal stem cells restore alveolar fluid clearance in human lungs rejected for transplantation. *Am J Transplant*. 2015;15:2404-2412.
1242. Pinky GS, Krishnakumar V, Sharma Y, Dinda AK, Mohanty S. Mesenchymal stem cell derived exosomes: a nano platform for therapeutics and drug delivery in combating COVID-19. *Stem Cell Rev Rep*. 2021;17:33-43.
1243. Jalalian SH, Ramezani M, Jalalian SA, Abnous K, Taghdisi SM. Exosomes, new biomarkers in early cancer detection. *Anal Biochem*. 2019;571:1-13.
1244. Kumar DN, Chaudhuri A, Aqil F, et al. Exosomes as emerging drug delivery and diagnostic modality for breast cancer: recent advances in isolation and application. *Cancer*. 2022;14:1435.
1245. Xi Y, Xu P. Global colorectal cancer burden in 2020 and projections to 2040. *Transl Oncol*. 2021;14:101174.
1246. Alzahrani SM, Al Doghaither HA, Al-Ghafari AB. General insight into cancer: an overview of colorectal cancer. *Mol Clin Oncol*. 2021;15:271.
1247. Yang J, Zhang L. The roles and therapeutic approaches of MSC-derived exosomes in colorectal cancer. *Clin Transl Oncol*. 2022;24:959-967.
1248. Guo G, Tan Z, Liu Y, Shi F, She J. The therapeutic potential of stem cell-derived exosomes in the ulcerative colitis and colorectal cancer. *Stem Cell Res Ther*. 2022;13:138.
1249. Arnold M, Abnet CC, Neale RE, et al. Global burden of 5 major types of gastrointestinal cancer. *Gastroenterology*. 2020;159:335-349.e15.
1250. Lu L, Fang S, Zhang Y, Jin L, Xu W, Liang Z. Exosomes and exosomal circRNAs: the rising stars in the progression, diagnosis and prognosis of gastric cancer. *Cancer Manage Res*. 2021;13:8121-8129.
1251. Xie M, Yu T, Jing X, et al. Exosomal circSHKBP1 promotes gastric cancer progression via regulating the miR-582-3p/HUR/VEGF axis and suppressing HSP90 degradation. *Mol Cancer*. 2020;19:112.
1252. Kahroba H, Hejazi MS, Samadi N. Exosomes: from carcinogenesis and metastasis to diagnosis and treatment of gastric cancer. *Cell Mol Life Sci*. 2019;76:1747-1758.
1253. Yang L, Huang X, Guo H, et al. Exosomes as efficient nanocarriers in osteosarcoma: biological functions and potential clinical applications. *Front Cell Dev Biol*. 2021;9:737314.
1254. Baglio SR, Lagerweij T, Pérez-Lanzón M, et al. Blocking tumor-educated MSC paracrine activity halts osteosarcoma progression. *Clin Cancer Res*. 2017;23:3721-3733.
1255. Notaro A, Emanuele S, Geraci F, et al. WIN55, 212-2-induced expression of Mir-29b1 favours the suppression of osteosarcoma cell migration in a SPARC-independent manner. *Int J Mol Sci*. 2019;20:5235.
1256. Zhang H, Wang J, Ren T, et al. Bone marrow mesenchymal stem cell-derived exosomal miR-206 inhibits osteosarcoma progression by targeting TRA2B. *Cancer Lett*. 2020;490:54-65.
1257. Zhang K, Dong C, Chen M, et al. Extracellular vesicle-mediated delivery of miR-101 inhibits lung metastasis in osteosarcoma. *Theranostics*. 2020;10:411-425.
1258. Ye Z, Zheng Z, Peng L. MicroRNA profiling of serum exosomes in patients with osteosarcoma by high-throughput sequencing. *J Invest Med*. 2020;68:893-901.
1259. Wang J-W, Wu X-F, Gu X-J, Jiang X-H. Exosomal miR-1228 from cancer-associated fibroblasts promotes cell migration and invasion of osteosarcoma by directly targeting SCAI. *Oncol Res*. 2019;27:979-986.
1260. Santos A, Domingues C, Jarak I, Veiga F, Figueiras A. Osteosarcoma from the unknown to the use of exosomes as a versatile and dynamic therapeutic approach. *Eur J Pharm Biopharm*. 2022;170:91-111.
1261. Izadi M, Dehghan Marvast L, Rezvani ME, et al. Mesenchymal stem-cell derived exosome therapy as a potential future approach for treatment of male infertility caused by chlamydia infection. *Front Microbiol*. 2022;12:4116.
1262. Mobarak H, Heidarpour M, Rahbarghazi R, Nouri M, Mahdipour M. Amniotic fluid-derived exosomes improved spermatogenesis in a rat model of azoospermia. *Life Sci*. 2021;274:119336.
1263. Mahiddine FY, Qamar AY, Kim MJ. Canine amniotic membrane derived mesenchymal stem cells exosomes addition in canine sperm freezing medium. *J Anim Reproduct Biotechnol*. 2020;35:268-272.
1264. Qamar AY, Fang X, Kim MJ, Cho J. Improved post-thaw quality of canine semen after treatment with exosomes from conditioned medium of adipose-derived mesenchymal stem cells. *Animals*. 2019;9:865.
1265. Mokarizadeh A, Rezvanfar M-A, Dorostkar K, Abdollahi M. Mesenchymal stem cell derived microvesicles: trophic shuttles for enhancement of sperm quality parameters. *Reprod Toxicol*. 2013;42:78-84.
1266. Herrmann IK, Wood MJA, Fuhrmann G. Extracellular vesicles as a next-generation drug delivery platform. *Nat Nanotechnol*. 2021;16:748-759.
1267. Kumar V, Kiran S, Kumar S, Singh UP. Extracellular vesicles in obesity and its associated inflammation. *Int Rev Immunol*. 2022;41:30-44.
1268. Ellulu MS, Patimah I, Khaza'i H, Rahmat A, Abed Y. Obesity and inflammation: the linking mechanism and the complications. *Arch Med Sci*. 2017;13:851-863.
1269. Engin A. The definition and prevalence of obesity and metabolic syndrome. In: Engin AB, Engin A, eds. *Obesity and Lipotoxicity Advances in Experimental Medicine and Biology*. Vol. 960. Springer; 2017:1-17.
1270. Khalyfa A, Sanz-Rubio D. The mystery of red blood cells extracellular vesicles in sleep apnea with metabolic dysfunction. *Int J Mol Sci*. 2021;22:4301.
1271. Waldman LT, Parthasarathy S, Villa KF, Bron M, Bujanover S, Brod M. Understanding the burden of illness of excessive daytime sleepiness associated with obstructive sleep apnea: a qualitative study. *Health Qual Life Outcomes*. 2020;18:128.
1272. He H, Tang J, Liu T, Hao W, Liao Y. Gender differences in sleep problems among drug users. *Front Psych*. 2020;11:808.
1273. Joo HS, Suh JH, Lee HJ, Bang ES, Lee JM. Current knowledge and future perspectives on mesenchymal stem cell-derived exosomes as a new therapeutic agent. *Int J Mol Sci*. 2020;21:727.

1274. Jafari D, Malih S, Eslami SS, et al. The relationship between molecular content of mesenchymal stem cells derived exosomes and their potentials: opening the way for exosomes based therapeutics. *Biochimie*. 2019;165:76-89.
1275. Phinney DG, Pittenger MF. Concise review: MSC-derived exosomes for cell-free therapy. *Stem Cells*. 2017;35:851-858.
1276. Del Fattore A, Luciano R, Pascucci L, et al. Immunoregulatory effects of mesenchymal stem cell-derived extracellular vesicles on T lymphocytes. *Cell Transplant*. 2015;24:2615-2627.
1277. Aryani A, Denecke B. Exosomes as a nanodelivery system: a key to the future of neuromedicine? *Mol Neurobiol*. 2016;53:818-834.
1278. Ye Q, Wang B, Mao J. Cytokine storm in COVID-19 and treatment. *J Infect*. 2020;80:607-613.
1279. Whitford W, Guterstam P. Exosome manufacturing status. *Future Med Chem*. 2019;11:1225-1236.
1280. Muraca M, Piccoli M, Franzin C, et al. Diverging concepts and novel perspectives in regenerative medicine. *Int J Mol Sci*. 2017;18:1021.
1281. Armstrong JP, Holme MN, Stevens MM. Re-engineering extracellular vesicles as smart nanoscale therapeutics. *ACS Nano*. 2017;11:69-83.
1282. Riazifar M, Pone EJ, Lötvall J, Zhao W. Stem cell extracellular vesicles: extended messages of regeneration. *Annu Rev Pharmacol Toxicol*. 2017;57:125-154.

How to cite this article: Miron RJ, Estrin NE, Sculean A, Zhang Y. Understanding exosomes: Part 2—Emerging leaders in regenerative medicine. *Periodontol 2000*. 2024;94:257-414. doi:[10.1111/prd.12561](https://doi.org/10.1111/prd.12561)